Tab. 1b Adjusted Odds Ratios from a multivariable <u>logistic regression</u> for self-reported current STI (diagnosed positive)

Current STI	Odds Ratio	p-value	95% CI	
Age group				
18-20	ref.			
21-30	0.39	0.192	0.10	1.60
31-40	0.89	0.866	0.24	3.37
41-50	3.73	0.052	0.99	14.05
>50	1.18	0.862	0.18	7.78
Number of sexual male partners in the last 12 months				
1	ref.			
2-4	0.59	0.399	0.18	2.00
5-12	2.26	0.126	0.80	6.41
>12	1.79	0.297	0.60	5.38
Frequency of visiting gay clubs, bars, discos to meet sexual partners				
Many times	ref.			
Sometimes	3.13	0.150	0.66	14.79
Rarely	4.38	0.062	0.93	20.72
Doesn't visit	6.06	0.030	1.20	30.76
Condom use in last sex intercourse				
No	ref.			
Yes	1.72	0.260	0.67	4.44
Popper use before or during last sex intercourse				
No	ref.			
Yes	2.45	0.081	0.89	6.71
Alcohol use before or during last sex intercourse				
No	ref.			
Yes	0.98	0.955	0.44	2.15

Tab. 2b Adjusted Odds Ratios from a multivariable <u>MIXED EFFECT</u> logistic regression for self-reported current STI (diagnosed positive) with age group as a random effect

Current STI	Odds Ratio	p-value	95% CI					
FIXED EFFECT								
Number of sexual male partners in the last 12 months								
1	ref.							
2-4	0.59	0.391	0.18	1.97				
5-12	2.25	0.125	0.80	6.36				
>12	1.85	0.269	0.62	5.52				
Frequency of visiting gay clubs, bars, discos to meet sexual partners								
Many times	ref.							
Sometimes	3.05	0.158	0.65	14.29				
Rarely	4.25	0.066	0.91	19.92				
Doesn't visit	6.05	0.029	1.20	30.39				
Condom use in last sex intercourse								
No	ref.							
Yes	1.70	0.270	0.66	4.37				
Popper use before or during last sex intercourse								
No	ref.							
Yes	2.32	0.100	0.85	6.30				
Alcohol use before or during last sex intercourse								
No	ref.							
Yes	0.97	0.936	0.44	2.12				
RANDOM EFFECT								
AGE GROUP (var)	0.50		0.10	2.43				

Tab. 3b Adjusted Odds Ratios from a multivariable <u>MIXED EFFECT</u> logistic regression for self-reported current STI (diagnosed positive) with number of sexual male partners in the last 12 months as a random effect

Current STI	Odds Ratio	p-value	95% CI			
	FIXED EFFECT			_		
Age group						
18-20	ref.					
21-30	0.39	0.185	0.10	1.57		
31-40	0.93	0.909	0.25	3.45		
41-50	3.83	0.045	1.03	14.18		
>50	1.18	0.864	0.18	7.63		
Frequency of visiting gay clubs, bars, disco	s					
to meet sexual partners						
Many times	ref.					
Sometimes	3.00	0.165	0.64	14.16		
Rarely	3.84	0.090	0.81	18.27		
Doesn't visit	4.78	0.060	0.94	24.35		
Condom use in last sex intercourse						
No	ref.					
Yes	1.90	0.177	0.75	4.83		
Popper use before or during last sex intercourse						
No	ref.					
Yes	2.62	0.057	0.97	7.06		
Alcohol use before or during last sex intercourse						
No	ref.					
Yes	0.97	0.946	0.44	2.13		
RANDOM EFFECT						
Number of sexual male partners in the last 12 months (var)	0.10		0.00	2.90		