








# BMJ Open Study protocol for a prospective cohort study identifying risk factors for sport injury in adolescent female football players: the Karolinska football Injury Cohort (KIC)

Ulrika Tranaeus <sup>1,2</sup>, Nathan Weiss <sup>2,3</sup>, Victor Lyberg,<sup>3</sup> Martin Hagglund <sup>4,5,6</sup>, Markus Waldén <sup>5,6,7,8</sup>, Urban Johnson <sup>9</sup>, Martin Asker <sup>2,3,10</sup>, Eva Skillgate <sup>2,3</sup>

**To cite:** Tranaeus U, Weiss N, Lyberg V, *et al.* Study protocol for a prospective cohort study identifying risk factors for sport injury in adolescent female football players: the Karolinska football Injury Cohort (KIC). *BMJ Open* 2022;**12**:e055063. doi:10.1136/bmjopen-2021-055063

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-055063>).

Received 29 July 2021

Accepted 17 December 2021



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Ulrika Tranaeus;  
[ulrika.tranaeus@gih.se](mailto:ulrika.tranaeus@gih.se)

## ABSTRACT

**Introduction** Football is a popular sport among young females worldwide, but studies concerning injuries in female players are scarce compared with male players. The aim of this study is to identify risk factors for injury in adolescent female football players.

**Methods and analysis** The Karolinska football Injury Cohort (KIC) is an ongoing longitudinal study that will include approximately 400 female football academy players 12–19 years old in Sweden. A detailed questionnaire regarding demographics, health status, lifestyle, stress, socioeconomic factors, psychosocial factors and various football-related factors are completed at baseline and after 1 year. Clinical tests measuring strength, mobility, neuromuscular control of the lower extremity, trunk and neck are carried out at baseline. Players are followed prospectively with weekly emails regarding exposure to football and other physical activity, health issues (such as stress, recovery, etc), pain, performance and injuries via the Oslo Sports Trauma Research Center Overuse Injury Questionnaire (OSTRC-O). Players who report a substantial injury in the OSTRC-O, that is, not being able to participate in football activities, or have reduced their training volume performance to a moderate or major degree, are contacted for full injury documentation. In addition to player data, academy coaches also complete a baseline questionnaire regarding coach experience and education.

**Ethics and dissemination** The study was approved by the Regional Ethical Review Authority at Karolinska Institutet, Stockholm, Sweden (2016/1251-31/4). All participating players and their legal guardians give their written informed consent. The study will be reported in accordance with the Strengthening the Reporting of Observational studies in Epidemiology. The results will be published in peer-reviewed academic journals and disseminated to the Swedish football movement through stakeholders and media.

## Strengths and limitations of this study

- A strength is the bio-psychosocial and multi-professional perspective of the risk of injuries in young female football players and factors of importance for not being injured, even though the bio-psychosocial factors are not equal included.
- Strengths are also the large sample size and the robust data collection of exposures, potential confounding factors, potential effect measure modifiers and outcomes.
- A potential limitation is the risk of misclassification of time varying exposures and outcomes in the weekly self-reported data collection.
- Using emails and SMS for weekly reports might decrease the response rates and thereby increase the risk of selection bias in the results. If the response turns out to be low, there is a risk of selection bias in the risk analyses.

## INTRODUCTION

Four million females worldwide are registered football players, of which 2.5 million are under 17 years old according to Fédération Internationale de Football Association.<sup>1</sup> Studies regarding injuries in female football players are fewer compared with the number of studies in male football players.<sup>2–4</sup> In brief, these studies show that common injuries in female football players are joint and ligament injuries to the knee and ankle joints as well as muscle and tendon injuries of the thigh. In addition, there is a particular concern for concussions and anterior cruciate ligament injuries in female players.<sup>3 5–8</sup>

Female football players have more absence days from football due to injuries compared with male players,<sup>8</sup> and long-term consequences of injuries might be considerable

for young football players.<sup>9</sup> For players with a history of injury, the risk of osteoarthritis in lower extremity joints are high and greater than in the general population.<sup>10 11</sup> Injuries may also lead to premature career ending,<sup>12</sup> and mental health problems.<sup>13</sup> Identifying risk factors for injury is, therefore, an important step towards reduction of injury risk.<sup>14</sup> To identify possible risk factors, well-designed prospective cohort studies are needed.<sup>15 16</sup> Specifically, the suggested risk factors in this setting can be classified as bio-psychosocial factors (see Wiese-Bjornstal for bio-psychosocial view on a sport injury risk profile).<sup>17</sup> Biological risk factors for injury in female players are previous injury,<sup>7 18–20</sup> a hamstring/quadriceps ratio of less than 55%, increased body mass index, as well as results of plyometric tests, for example, poor performance in drop jump landing test is associated with increased risk of ankle injury.<sup>21</sup> Other biological risk factors associated with an increased risk of injury during the season are young age,<sup>6 18</sup> physical complaints at the beginning of the season,<sup>18</sup> familial disposition such as a parent/sibling<sup>18</sup> or a twin<sup>22</sup> with knee injury, and lower level of preseason aerobic fitness.<sup>23 24</sup> Findings regarding joint hypermobility as a risk factor in female players are inconclusive in older studies,<sup>23 25</sup> although no association was shown in more recently published studies.<sup>26 27</sup> Risk factors for back pain in adolescents include rapid growth rate, and tight muscle imbalance,<sup>28</sup> but risk factors for football related back/neck injuries in young females are not known. Psychological risk factors reported includes somatic trait anxiety, mistrust, ineffective coping,<sup>29</sup> life event stress<sup>30</sup> and perceived mastery climate.<sup>20</sup> Social factors that influenced the risk for injury in female athletes are coaches' and player's education regarding injury prevention strategies,<sup>31</sup> stress from teammates and coaches,<sup>20 29 32</sup> and for back pain in adolescents; smoking.<sup>28</sup> In football, an identified situational specific risk factor is the playing positions defenders and strikers.<sup>19</sup>

In summary, most knowledge about risk factors for injuries in adolescent female football players consists of isolated factors, and lack of using multidisciplinary and bio-psychosocial perspectives. Hence, the overall aim of the Karolinska football Injury Cohort study (KIC) is to identify risk factors for injuries in adolescent female football players from a bio-psychosocial perspective. Specific aims are to determine the incidence of injuries in young female football players and to identify modifiable risk factors for such injuries. Finally, our aims include to describe changes in muscle strength and range of motion (ROM) over a year, trajectories of pain and to identify important factors for not being injured over a year.

## METHODS AND ANALYSIS

This is a prospective observational cohort study designed in agreement with Strengthening the Reporting of Observational studies in Epidemiology (STROBE) guidelines.<sup>33</sup>

## Study setting and participants

Football clubs with adolescent female academy players aged 12–19 years, participating in Swedish divisions 1–2 for girls in the largest regions, are eligible to participate in the study. Most players will be recruited in Stockholm. The district of Stockholm consists of 140 teams and approximate 2520 female players, 13–19 years old. Clubs which meet the inclusion criteria are contacted and invited to participate and are given oral and written information. Clubs which choose to take part in the study are provided with a more detailed oral and written information in the presence of players, legal guardians and coaches.

A cohort of approximately 400 adolescent academy players will be recruited. An internal pilot study of 63 football players has been conducted to test the infrastructure and the implementation of the study, with satisfactory results (unpublished data).

## Baseline measurements

### Questionnaires

The baseline questionnaire covers potential risk factors for the aetiology of sport injuries as well as information about players' general health status. Players are surveyed in various areas, including *health*: health problems (eg, illness), medication, age at menarche, amenorrhoea; *lifestyle*: sleep patterns, eating habits, food supplements, tobacco as smoking or Swedish snus (snuff) and alcohol; and *socioeconomic* factors: guardians' education. Included *football-related factors* are: training and match play exposure, playing position, dominant limb, years of experience, other sports participation, injury preventive strategies (eg, the Swedish injury prevention warmup programme Knee Control)<sup>34</sup> and type of turf at the home facilities (artificial or natural grass) according to guidelines for football studies.<sup>35–37</sup> *Psychosocial factors* are surveyed using: the modified General Health Questionnaire-12 consisting of 12 items regarding self-reported general psychological health using a four-point Likert scale,<sup>38</sup> coping assessed by a 28 item self-report questionnaire that measure effective and ineffective strategies to cope with stressful events using a four-point Likert scale (Brief COPE),<sup>39</sup> player's passion to sport measured in harmonious and obsessive passions using a 14 item questionnaire with a seven-point Likert scale (Passion scale),<sup>40</sup> education in sport psychology, regularly seeing a sport psychologist/mental coach and perceived stress (single item question).<sup>41</sup> *Injury history*: injuries occurring within the previous 6 months prior to inclusion are captured using a modified Swedish version of the validated psychometric instrument Oslo Sports Trauma Research Center Overuse Injury Questionnaire (OSTRC-O).<sup>42 43</sup> *Back and neck pain* is covering the frequency, intensity, disability of low back pain (LBP) and upper back pain/neck pain (UBNP) and corresponding longitudinal trajectories the preceding 6 months using modified versions of the Chronic Pain Questionnaire<sup>44</sup> and Visual Trajectories Questionnaire-Pain,<sup>45</sup> respectively.

In addition, coaches in the included teams are surveyed regarding their education, years of experience, the use

of warmup and stretching regime and implementation of injury prevention programmes.

### Physical test protocol

The physical test protocol includes several tests that are considered valid, reliable, and field friendly; performed in approximately 60 min/player. The protocol comprises measurements of strength, mobility and control of lower extremity, trunk and neck and also include anthropometric measurements (height, weight and leg length). The protocol is briefly outlined below, however, further details including visual presentations is available in the electronic supplementary file (online supplemental file 1).

All test procedures are conducted in indoors facilities during weekends. The physical tests are divided into nine test stations with 1–2 test leaders each (online supplemental file 1). Hitherto, 52 clinically experienced test leaders have been involved in data collection. They were trained by MA, VL, NW and the previous test leader in charge of the station to ensure consistent execution and reliability. Information and instructions given to the players regarding the tests are standardised, and test leaders refrain from coaching or encouraging the players in any way during the procedures.

A maximum of nine players are tested per session (ie, one at each station) and are informed to train and compete as usual prior to testing. Players are informed to refrain from certain tests that evoke pain, provoke ongoing injuries or other health-related issues. Prior to performing the physical tests, players complete a standardised 7 min warm-up programme comprising 4 min of jogging, 10×1 body weight squats, 10×1 body weight squat jumps and 10×1 unilateral body weight lunges. Following the warm-up session, players are randomly assigned to a starting test station and subsequently follow a predefined order.

### Calf heel raises

Ankle plantarflexion (PF) muscle endurance is investigated using unilateral weight bearing calf heel raises.<sup>46</sup> The player is instructed to perform maximum unilateral barefoot heel raises continuously to failure, guided by a metronome to standardise the pace (1 s concentric, 1 s eccentric contraction). The test leader registers the number of accomplished repetitions and discontinues when the player fails to reach the marked target height. The same procedure is then conducted on the opposite foot.

### Active PF mobility

Active PF ROM is measured with a universal goniometer in supine position using fibula and fifth metatarsal as reference marks.<sup>47 48</sup> The player is instructed to maintain extended knees throughout the movement, and to perform a sequence of six maximal active PF cycles from a neutral dorsiflexion (DF) position, of which the final three trials are registered.

### Weight bearing ankle DF mobility

Weight bearing ankle DF ROM is measured in a lunge position with the player's foot placed on a metric ruler 10 cm away from a wall.<sup>46 49</sup> The player is instructed to lunge forward, until contact with the wall is achieved without allowing the heel to lift off the ground. Three warm-up trials are performed from the 10 cm mark to familiarise the player with the test. Thereafter, the test leader measures the following three trials. From the 10 cm reference mark, the player progresses 1 cm away at a time from the wall until unable to perform a successful repetition. If unable to perform a successful repetition at the 10 cm reference mark, she is asked to progress 1 cm forward until able to complete a successful repetition. The maximal DF ROM is measured with a digital inclinometer (Clinometer, Plaincode, Stephanskirchen, Germany) and distance from the wall to the greater toe is measured in centim.

### Trunk mobility

Trunk rotation mobility is measured in a modified seated rotation test, and in a lunge position on a gym mat graded with 5° increments.<sup>50–52</sup> The player is instructed to maximally rotate alternating between right and left, in a cross-legged position and subsequently in a lunge position on the dominant, and non-dominant limb while the test leader measures the rotational degrees in the end range. Three repetitions are performed in each direction during the three separate positions, and the mean value for each position is later used for analysis.

### Trunk strength

Isometric trunk rotational strength is measured in a modified standing wood chopper test using a force gauge to evaluate force output (RS Pro Digital Force Gauge, RS Components, Corby, UK).<sup>53–55</sup> In this modified test, the player holds a handle attached to the force gauge in shoulder height in a standing position. The player is instructed to generate force through her trunk and rotate for 5 s while maintaining straight arms. Three consecutive repetitions are conducted in each direction and the maximal force output is later used for analysis.

### Deep neck flexor endurance

Deep neck flexor muscle endurance is assessed through a modified version of the cranio-cervical flexion test with a pressure sensor (Stabilizer Pressure Bio-Feedback, Chattanooga Group, Hixson, Tennessee, USA).<sup>51 56 57</sup> The test consists of a pretest and an endurance test. In the pretest the player is positioned in a supine position on an examination table and are instructed to slightly push the neck against the pressure sensor to increase the pressure and then maintain the pressure for 3×3 s, with a 3 s rest in between each contraction, at a specific target pressure (TP), starting at 20 mm Hg. If the player can perform this task, she is instructed to increase the pressure to 24 mm Hg and keep the pressure for another 3×3 s. This is repeated with a 2 mm Hg increase until the



player reaches 30 mm Hg. If the player can perform the pretest the endurance test is subsequently performed. During the endurance test, the same setup and procedure as in the pretest is carried out. However, the player is instructed to hold each contraction at the TP for 3×10 s with a 10 s rest in between contractions. The highest completed TP with a full set of 3×10 s contractions is later used for analysis.

### Hip and knee strength

Isometric hip flexion, extension, adduction and abduction strength as well as eccentric hip abduction and adduction strength are measured with a hand-held dynamometer (HHD) (MicroFet2, Hoggan Health Industries inc. West Jordan, Utah, USA).<sup>58 59</sup> Furthermore, isometric knee extension strength is measured with an HHD and the player in a seated position with the knee joint in 90° of flexion. Prior to executing the strength tests, two submaximal isometric contractions in each direction are performed to familiarise the player with the procedures. Three isometric contractions with gradually increasing power output for 5 s, and three maximally eccentric contractions for 3 s are performed in the isometric and eccentric tests, respectively, with a 10 s rest in between contractions. The maximal power output for each position is later used for analysis.

### Hip mobility

Measures of passive hip ROM in flexion and abduction in prone position and extension, internal and external rotation in supine position is obtained using a universal goniometer.<sup>60 61</sup> Three consecutive measurements for each position are performed for both the dominant and the non-dominant leg, and the mean value for each position is later used for analysis.

### Functional performance tests

To assess the player's unilateral jump performance, the One-leg Long Box Jump Test (OLLBJ) and square hop test are performed.<sup>62 63</sup> A 40×40 cm square is marked on the foundation and later used as a reference mark in both tests.

In the OLLBJ, the starting position are calculated by dividing the player's height (cm) with 1.6 (height/1.6). Thereafter, the player is instructed to stand on one leg on the starting position and then jump on one leg directed inside the boundaries of the square and maintain balance after landing. Three warm-up trials and five consecutive test trials are performed on each leg. The total number of approved trials are registered by the test leader.

During the square hop test, previously described in detail,<sup>62 63</sup> the player is instructed to jump on one leg in and out of the square as many times as possible for 15 s in a clockwise direction, timed with a stopwatch while the test leader registers the number of approved jumps. The player performs two warm up trials on each foot prior to executing the test.

### Ankle and knee stability

To assess stability of player's talocrural joints, a modified anterior drawer test is employed.<sup>64 65</sup> Furthermore, a modified version of Fairbank's apprehension test is used to evaluate the player's stability in the patellofemoral joint.<sup>66</sup> The tests are conducted on both the dominant and non-dominant foot and knee and are considered positive if the player experience any pain or discomfort during the examination, and/or an involuntary contraction of the quadriceps musculature occur during the Fairbank's apprehension test.

### Isometric back extensor endurance

Isometric back extensor endurance is assessed by the modified Sorensen test.<sup>67–69</sup> In this previously described modified test,<sup>67 68</sup> the player's lower body is supported by an examination table in prone position with three straps and the anterior–superior iliac spine is aligned with the edge of the table. The player is instructed to keep her arms folded across the chest throughout the procedure and isometric maintaining the upper body in a horizontal position until failure. The test leader registers the time elapsed until failure. A digital inclinometer (Clinometer, Plaincode, Stephanskirchen, Germany) is placed on a metric ruler at the level of the 5th vertebra in the thoracic spine to monitor sagittal plane movement. Prior to the assessment, the player completes a shorter warmup trial to orient the desired sagittal plane target angle.

### Follow-up measurement and outcome

Follow-up measurements are collected prospectively during 1 year from the baseline. In the weekly online questionnaire, the players are asked to answer questions regarding new and ongoing injuries, LBP and UBNP intensity, social support, perceived stress, recovery, and to be able to consider workload, number of training and match play hours/week.<sup>70</sup> To assess whether players sustain football related injuries throughout the follow-up period, the Swedish version of OSTRC-O is employed and included in the weekly online questionnaire.<sup>42 43 71</sup> Two study specific adaptations were made to the OSTRC-O. First, a question regarding absence/reduced participation in training/match due to reasons not related to injuries was added. Second, the option to specify injuries in different anatomical localisations in the lower and upper extremity, back, neck, head and abdomen was included.

Football related injuries reported with the OSTRC-O in the weekly online questionnaire leading to moderate or severe reductions in participation and/ or sports performance or complete inability to participate in sport are classified as a substantial injury in this study.<sup>42</sup> Players reporting new substantial injuries are contacted via telephone by a clinically experienced research assistant to answer a standardised interview with questions concerning the injury such as: injury mechanism, localisation, type, time-loss, re-injury, diagnosis and medical care. Injuries are divided into acute and gradual onset. An acute injury is defined as a result from a specific,

identifiable event, whereas injuries with gradual onset are defined as an injury without a single, identifiable event responsible for the injury.<sup>35</sup> Players receive an automated link to the online questionnaire sent by email each Sunday, with a reminder email the next day to players not responding. Furthermore, if no response is received, a text message reminder with the link is sent on Tuesdays. Finally, every other week representatives of the study visit participating football clubs to collect unanswered surveys for the previous 2-week period.

After 52 weeks of participation, a questionnaire with equivalent content as the baseline questionnaire (excluding OSTRC-O with 2 months and 3–6 months recall) are distributed to the players to evaluate possible changes from the baseline characteristics. The first 106 included players also underwent a secondary physical test protocol after 52 weeks of follow-up. In the 1-year follow-up questionnaire, different aspects of UBP and LBP, respectively, in the preceding 6 months are measured. ‘Have you had UBP/Have you had LBP’ (Yes/No)? If yes, has the pain hindered your daily activities (No, Yes to some

extent or Yes to a high degree)? If Yes, the ‘VTQ-P’ is used to capture the longitudinal state of a player’s pain experience of UBP and LBP and are retrospectively reported for the preceding 6-month period.<sup>45</sup> See [table 1](#) for an overview of the measurements during the different phases of the study.

### Sample size

The statistical power for the analyses will depend on the exact research question, the number of exposed players and whether the exposure is continuous or categorised. The sample size in the KIC project is based on the definition ‘a substantial injury’ as proposed by Clarsen *et al*<sup>42</sup> and back injuries in adolescent female players in a previously published study.<sup>7</sup> Based on a relative risk of 1.9 for a substantial injury in the back/neck, when 88 of the players are exposed, and with a power of 0.80, a significance level 5% and with potential 10% drop out and a follow-up time of 1 year to identify risk factors, 420 players will be included.

**Table 1** Summary of the included measurements during the different phases of the study

Phase	Measurements	Tests/tools
Baseline: players (consecutive during inclusion; 2016–ongoing)	Demographic information, general health status (history of pain, illness, medication, plagues, menstrual cycle, back and neck pain), lifestyle (sleep patterns, resilience, food supplements, use of tobacco or alcohol), stress, socioeconomic factors (guardians’ education), football related factors (position, years of experience, injury preventive strategies)	KIC Baseline players, The Chronic Pain Questionnaire (CPQ), <sup>44</sup> Visual Trajectories Questionnaire-Pain (VTQ-P) <sup>45</sup>
	Anthropometric measurements (height, weight, leg length), and measurement of strength, mobility and control of lower extremity, trunk, and neck	KIC test protocol
	History of injury and complaints	Modified OSTRC-O <sup>42 43</sup>
	Passion	Passion scale <sup>40</sup>
	General health	GHQ-12 <sup>38</sup>
Baseline: coaches (consecutive during inclusion; 2016–ongoing)	Coping strategies	Brief COPE <sup>39</sup>
	Education, years of experience, the use of warmup and stretching regime and implementation of injury prevention programmes	KIC Baseline coaches
	Weekly follow-up: players (September 2016–ongoing)	KIC weekly report
	Exposure to football training and match play	
	Exposure to other physical activity	
Weekly follow-up: players (September 2016–ongoing)	Health (eg, stress, recovery) and social support	
	Report on pain, injury performance, complaints	Modified OSTRC-O <sup>42 43</sup>
	Report on injury/complaint (type of injury, localisation, inciting event)	KIC medical report
	One-year follow-up: players (consecutive after 52 weeks participation: 2017–ongoing)	KIC 1-year questionnaire
	Football related factors (position, injury preventive strategies). Health status (pain in back or neck) lifestyle (sleep patterns, resilience, food supplements, use of tobacco or alcohol, physical activity), stress, coping and passion for sport	
One-year follow-up (consecutive after 52 weeks participation in the first 106 included players)	Anthropometric measurements (height, weight, leg length), and measurement of strength, mobility and control of lower extremity, trunk and neck	KIC test protocol

GHQ-12, General Health Questionnaire-12; KIC, Karolinska football Injury Cohort; OSTRC-O, Oslo Sports Trauma Research Center Overuse Injury Questionnaire.

## Statistical methods

The data in the KIC study will be used to answer several different research questions and therefore, different analyses methods and statistics will be used. Kaplan-Meier estimates will be used to describe incidence. Cox regression analyses or discrete time survival analyses will be used to measure the associations between exposure and outcome, and to adjust for confounding. Only players without substantial injuries the two preceding months (reported in the baseline questionnaire) will be considered in the risk analyses, and stratified analyses to examine effect measure modification will be performed when relevant. The development of injuries is likely complex. This justifies why we measure an extensive number of factors so that we can consider confounders, intermediators and effect measure modifier in these analyses. When identifying trajectories of time, and various factors generalised estimating equations will be used for these analyses to consider the covariance between repeated measurements.

## Time plan

Players will be recruited from 2016 and followed weekly for 1 year regarding injuries/complaints. Players will consecutively be invited and included from the year they turn 13 years old and play in a participating club. The inclusion of participants will continue until we reach over 400 players.

## Data statement

The dataset and statistical codes will be available on reasonable request when the data collection is completed.

## Patient and public involvement

No patient involved.

## ETHICS AND DISSEMINATION

The study was approved by the Regional Ethical Review Authority at Karolinska Institutet, Stockholm, Sweden (2016/1251-31/4). All participating players and their legal guardians receive written and oral information regarding the study and give their written informed consent when entering the study. Players under the age of 15 are required to have written informed consent from their legal guardians. The study will be performed in accordance with the recommendations guiding research involving human subjects adopted by the 18th World Medical Association General Assembly, Helsinki, Finland, June 1964, amended at the 64th World Medical Association General Assembly, Fortaleza, Brazil, October 2013. The study will be reported in accordance with the STROBE.<sup>33</sup> The results will be presented in scientific conferences and published in peer-reviewed academic journals as well as being disseminated to the Swedish football movement through stakeholders and media.

## Author affiliations

<sup>1</sup>Swedish School of Sport and Health Sciences, Stockholm, Sweden

<sup>2</sup>Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

<sup>3</sup>Department of Health Promotion Science, Sophiahemmet University, Stockholm, Sweden

<sup>4</sup>Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden

<sup>5</sup>Sport Without Injury Programme (SWIPE), Linköping University, Linköping, Sweden

<sup>6</sup>Football Research Group, Linköping, Sweden

<sup>7</sup>Department of Orthopaedics, Håssleholm-Kristianstad Hospitals, Håssleholm, Sweden

<sup>8</sup>Unit of Community Medicine, Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden

<sup>9</sup>Halmstad University, Halmstad, Sweden

<sup>10</sup>Scandinavian College of Naprapathic Manual Medicine, Stockholm, Sweden

**Twitter** Ulrika Tranaeus @UlrikaTranaeus, Nathan Weiss @nathnweiss, Martin Hagglund @MHggglund, Markus Waldén @MarkusWalden and Martin Asker @martinasker

**Acknowledgements** Martin Samuelsson, Lena Holm and Henrik Källberg have been involved in planning of study design, Joakim Bogren, Victor Ramirez Kristiansen, contributed to the first version of the draft. Students from the classes DN 47, 48, 49, 50, 51 and 53 at the Scandinavian college of Naprapathic Manual Medicine, Stockholm, Sweden are participating during inclusion of participants, the physical tests and the prospective data collection.

**Contributors** VL, ES, MA and UT initiated the study. All authors, UT, NW, VL, MH, MW, UJ and ES conceived the study and contributed to the development of the study protocol. ES is the study guarantor. UT and NW bi-drafted the manuscript which was critically revised in steps by all coauthors. The final manuscript was approved by all authors.

**Funding** This study is funded by grants from Swedish Research Council for Sport Science (grant number: P2019-0045, P2020-0100). The Swedish Naprapathic, and the Norwegian Naprapathic Associations, Active life foundation and Sophiahemmet foundation. There are no grant numbers for the latter funders (award/grant number: N/A).

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

## ORCID iDs

Ulrika Tranaeus <http://orcid.org/0000-0002-2102-6352>

Nathan Weiss <http://orcid.org/0000-0002-1814-020X>

Martin Hagglund <http://orcid.org/0000-0002-6883-1471>

Markus Waldén <http://orcid.org/0000-0002-6790-4042>

Urban Johnson <http://orcid.org/0000-0003-0990-4842>

Martin Asker <http://orcid.org/0000-0002-5387-3572>

Eva Skillgate <http://orcid.org/0000-0003-2096-1530>

## REFERENCES

- 1 Federation Internationale de Football Association. *Women's football survey*, 2014.
- 2 López-Valenciano A, Ruiz-Pérez I, García-Gómez A, *et al*. Epidemiology of injuries in professional football: a systematic review and meta-analysis. *Br J Sports Med* 2020;54:711–8.



- 3 Bennett P, Fawcett L. Trauma injuries sustained by female footballers. *Trauma* 2006;8:69–76.
- 4 Hägglund M, Waldén M, Ekstrand J. Injuries among male and female elite football players. *Scand J Med Sci Sports* 2009;19:819–27.
- 5 Faude O, Junge A, Kindermann W, et al. Injuries in female soccer players: a prospective study in the German national League. *Am J Sports Med* 2005;33:1694–700.
- 6 Le Gall F, Carling C, Reilly T. Injuries in young elite female soccer players. *Am J Sports Med* 2008;36:276–84.
- 7 Clausen MB, Zebis MK, Möller M, et al. High injury incidence in adolescent female soccer. *Am J Sports Med* 2014;42:2487–94.
- 8 Larruskain J, Lekue JA, Diaz N, et al. A comparison of injuries in elite male and female football players: a five-season prospective study. *Scand J Med Sci Sports* 2018;28:237–45.
- 9 Lohmander LS, Ostberg A, Englund M, et al. High prevalence of knee osteoarthritis, pain, and functional limitations in female soccer players twelve years after anterior cruciate ligament injury. *Arthritis Rheum* 2004;50:3145–52.
- 10 Prien A, Prinz B, Dvořák J, et al. Health problems in former elite female football players: prevalence and risk factors. *Scand J Med Sci Sports* 2017;27:1404–10.
- 11 Roos EM. Joint injury causes knee osteoarthritis in young adults. *Curr Opin Rheumatol* 2005;17:195–200.
- 12 Wylleman P, Reints A. A lifespan perspective on the career of talented and elite athletes: perspectives on high-intensity sports. *Scand J Med Sci Sports* 2010;20:88–94.
- 13 Putukian M. The psychological response to injury in student athletes: a narrative review with a focus on mental health. *Br J Sports Med* 2016;50:145–8.
- 14 van Mechelen W, Hlobil H, Kemper HC. Incidence, severity, aetiology and prevention of sports injuries. A review of concepts. *Sports Med* 1992;14:82–99.
- 15 Dvorak J, Junge A, Chomiak J, et al. Risk factor analysis for injuries in football players - Possibilities for a prevention program. *Am J Sports Med* 2000;28:S69–74.
- 16 Steffen K, Engebretsen L. More data needed on injury risk among young elite athletes. *Br J Sports Med* 2010;44:485–9.
- 17 Wiese-Bjornstal DM. Psychology and socioculture affect injury risk, response, and recovery in high-intensity athletes: a consensus statement. *Scand J Med Sci Sports* 2010;20:103–11.
- 18 Hägglund M, Waldén M. Risk factors for acute knee injury in female youth football. *Knee Surg Sports Traumatol Arthrosc* 2016;24:737–46.
- 19 Faude O, Junge A, Kindermann W, et al. Risk factors for injuries in elite female soccer players. *Br J Sports Med* 2006;40:785–90.
- 20 Steffen K, Pensgaard AM, Bahr R. Self-reported psychological characteristics as risk factors for injuries in female youth football. *Scand J Med Sci Sports* 2009;19:442–51.
- 21 Alahmad TA, Kearney P, Cahalan R. Injury in elite women's soccer: a systematic review. *Phys Sportsmed* 2020:1–7.
- 22 Magnusson K, Turkiewicz A, Hughes V, et al. High genetic contribution to anterior cruciate ligament rupture: heritability ~69. *Br J Sports Med* 2020;55:385–9.
- 23 Ostberg A, Roos H. Injury risk factors in female European football. A prospective study of 123 players during one season. *Scand J Med Sci Sports* 2000;10:279–85.
- 24 Watson A, Brickson S, Brooks MA, et al. Preseason aerobic fitness predicts in-season injury and illness in female youth athletes. *Orthop J Sports Med* 2017;5:232596711772697.
- 25 Söderman K, Alfredson H, Pietilä T, et al. Risk factors for leg injuries in female soccer players: a prospective investigation during one out-door season. *Knee Surg Sports Traumatol Arthrosc* 2001;9:313–21.
- 26 Nilstad A, Andersen TE, Bahr R, et al. Risk factors for lower extremity injuries in elite female soccer players. *Am J Sports Med* 2014;42:940–8.
- 27 Blokland D, Thijs KM, Backx FJG, et al. No effect of generalized joint hypermobility on injury risk in elite female soccer players: a prospective cohort study. *Am J Sports Med* 2017;45:286–93.
- 28 Ehrmann Feldman D, Shrier I, Rossignol M, et al. Risk factors for the development of neck and upper limb pain in adolescents. *Spine* 2002;27:523–8.
- 29 Johnson U, Ivarsson A. Psychological predictors of sport injuries among junior soccer players. *Scand J Med Sci Sports* 2011;21:129–36.
- 30 Ivarsson A, Johnson U, Lindwall M, et al. Psychosocial stress as a predictor of injury in elite junior soccer: a latent growth curve analysis. *J Sci Med Sport* 2014;17:366–70.
- 31 Junge A, Röscher D, Peterson L, et al. Prevention of soccer injuries: a prospective intervention study in youth amateur players. *Am J Sports Med* 2002;30:652–9.
- 32 Pensgaard AM, Ivarsson A, Nilstad A, et al. Psychosocial stress factors, including the relationship with the coach, and their influence on acute and overuse injury risk in elite female football players. *BMJ Open Sport Exerc Med* 2018;4:e000317.
- 33 von Elm E, Altman DG, Egger M, et al. The strengthening of reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Int J Surg* 2014;12:1495–9.
- 34 Waldén M, Atroshi I, Magnusson H, et al. Prevention of acute knee injuries in adolescent female football players: cluster randomised controlled trial. *BMJ* 2012;344:e3042.
- 35 Fuller CW, Ekstrand J, Junge A. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Br J Sports Med* 2006;40:193–201.
- 36 Hägglund M, Waldén M, Bahr R, et al. Methods for epidemiological study of injuries to professional football players: developing the UEFA model. *Br J Sports Med* 2005;39:340–6.
- 37 Emery CA, Meeuwisse WH, Hartmann SE. Evaluation of risk factors for injury in adolescent soccer: implementation and validation of an injury surveillance system. *Am J Sports Med* 2005;33:1882–91.
- 38 Hardy GE, Shapiro DA, Haynes CE, et al. Validation of the General Health Questionnaire-12: using a sample of employees from England's health care services. *Psychol Assess* 1999;11:159–65.
- 39 Carver CS. You want to measure coping but your protocol's too long: consider the Brief cope. *Int J Behav Med* 1997;4:92–100.
- 40 Vallerand RJ, Blanchard C, Mageau GA, et al. Les passions de l'âme: on obsessive and harmonious passion. *J Pers Soc Psychol* 2003;85:756–67.
- 41 Salminen S, Kouvonen A, Koskinen A, et al. Is a single item stress measure independently associated with subsequent severe injury: a prospective cohort study of 16,385 forest industry employees. *BMC Public Health* 2014;14:543.
- 42 Clarsen B, Myklebust G, Bahr R. Development and validation of a new method for the registration of overuse injuries in sports injury epidemiology: the Oslo Sports Trauma Research Centre (OSTRC) overuse injury questionnaire. *Br J Sports Med* 2013;47:495–502.
- 43 Ekman E, Frohm A, Ek P, et al. Swedish translation and validation of a web-based questionnaire for registration of overuse problems. *Scand J Med Sci Sports* 2015;25:104–9.
- 44 Von Korf M, Ormel J, Keefe FJ, et al. Grading the severity of chronic pain. *Pain* 1992;50:133–49.
- 45 Dunn KM, Campbell P, Jordan KP. Validity of the visual trajectories questionnaire for pain. *J Pain* 2017;18:1451–8.
- 46 Dennis RJ, Finch CF, Elliott BC, et al. The reliability of musculoskeletal screening tests used in cricket. *Phys Ther Sport* 2008;9:25–33.
- 47 Youdas JW, Bogard CL, Suman VJ. Reliability of goniometric measurements and visual estimates of ankle joint active range of motion obtained in a clinical setting. *Arch Phys Med Rehabil* 1993;74:1113–8.
- 48 Ness BM, Sudhagani RG, Tao H, et al. The reliability of a novel heel-rise test versus goniometry to assess plantarflexion range of motion. *Int J Sports Phys Ther* 2018;13:19–27.
- 49 Konor MM, Morton S, Eckerson JM, et al. Reliability of three measures of ankle dorsiflexion range of motion. *Int J Sports Phys Ther* 2012;7:279.
- 50 Frohm A, Heijne A, Kowalski J, et al. A nine-test screening battery for athletes: a reliability study. *Scand J Med Sci Sports* 2012;22:306–15.
- 51 Asker M, Waldén M, Källberg H, et al. A prospective cohort study identifying risk factors for shoulder injuries in adolescent elite handball players: the Karolinska Handball Study (KHAST) study protocol. *BMC Musculoskelet Disord* 2017;18:485.
- 52 Johnson KD, Kim K-M, Yu B-K, et al. Reliability of thoracic spine rotation range-of-motion measurements in healthy adults. *J Athl Train* 2012;47:52–60.
- 53 Andre MJ, Fry AC, Heyrman MA, et al. A reliable method for assessing rotational power. *J Strength Cond Res* 2012;26:720–4.
- 54 Palmer TG, Uhl TL. Interday reliability of peak muscular power outputs on an isotonic dynamometer and assessment of active trunk control using the CHOP and lift tests. *J Athl Train* 2011;46:150–9.
- 55 Zois J, P Sharp A, Talukdar K, et al. The reliability of a rotational power assessment of the core. *J Athl Enhanc* 2016;5.
- 56 Arumugam A, Mani R, Raja K. Interrater reliability of the craniocervical flexion test in asymptomatic individuals—a cross-sectional study. *J Manipulative Physiol Ther* 2011;34:247–53.
- 57 James G, Doe T. The craniocervical flexion test: intra-tester reliability in asymptomatic subjects. *Physiother Res Int* 2010;15:144–9.
- 58 Thorborg K, Bandholm T, Hölmich P. Hip- and knee-strength assessments using a hand-held dynamometer with external belt-fixation are inter-tester reliable. *Knee Surg Sports Traumatol Arthrosc* 2013;21:550–5.

- 59 Kelln BM, McKeon PO, Gontkof LM, *et al.* Hand-held dynamometry: reliability of lower extremity muscle testing in healthy, physically active, young adults. *J Sport Rehabil* 2008;17:160–70.
- 60 Prather H, Harris-Hayes M, Hunt DM, *et al.* Reliability and agreement of hip range of motion and provocative physical examination tests in asymptomatic volunteers. *PM R* 2010;2:888–95.
- 61 Nussbaumer S, Leunig M, Glatthorn JF, *et al.* Validity and test-retest reliability of manual goniometers for measuring passive hip range of motion in femoroacetabular impingement patients. *BMC Musculoskelet Disord* 2010;11:194.
- 62 Sharma N, Sharma A, Singh Sandhu J. Functional performance testing in athletes with functional ankle instability. *Asian J Sports Med* 2011;2:249.
- 63 Caffrey E, Docherty CL, Schrader J, *et al.* The ability of 4 single-limb hopping tests to detect functional performance deficits in individuals with functional ankle instability. *J Orthop Sports Phys Ther* 2009;39:799–806.
- 64 Vaseenon T, Gao Y, Phisitkul P. Comparison of two manual tests for ankle laxity due to rupture of the lateral ankle ligaments. *Iowa Orthop J* 2012;32:9–16.
- 65 de Vries JS, Kerkhoffs GMMJ, Blankevoort L, *et al.* Clinical evaluation of a dynamic test for lateral ankle ligament laxity. *Knee Surg Sports Traumatol Arthrosc* 2010;18:628–33.
- 66 Smith TO, Clark A, Neda S, *et al.* The intra- and inter-observer reliability of the physical examination methods used to assess patients with patellofemoral joint instability. *Knee* 2012;19:404–10.
- 67 Moreau CE, Green BN, Johnson CD, *et al.* Isometric back extension endurance tests: a review of the literature. *J Manipulative Physiol Ther* 2001;24:110–22.
- 68 Demoulin C, Vanderthommen M, Duysens C, *et al.* Spinal muscle evaluation using the Sorensen test: a critical appraisal of the literature. *Joint Bone Spine* 2006;73:43–50.
- 69 Latimer J, Maher CG, Refshauge K, *et al.* The reliability and validity of the Biering-Sorensen test in asymptomatic subjects and subjects reporting current or previous nonspecific low back pain. *Spine* 1999;24:2085–9.
- 70 Malone S, Owen A, Newton M, *et al.* The acute:chronic workload ratio in relation to injury risk in professional soccer. *J Sci Med Sport* 2017;20:561–5.
- 71 Nilstad A, Bahr R, Andersen TE. Text messaging as a new method for injury registration in sports: a methodological study in elite female football. *Scand J Med Sci Sports* 2014;24:243–9.



This is a supplementary document describing the included tests in Karolinska football Injury Cohort, KIC. Table S1 shows the test stations, number of test leaders and randomization of the tests. The persons in the images have given their consent that the images will be used in publications related to this study.

### **Calf heel raises**

Ankle plantarflexion (PF) muscle endurance is investigated using unilateral barefoot weight bearing calf heel raises<sup>46</sup>. Firstly, the player's maximal weight bearing PF range of motion (ROM) is obtained by painting a reference mark on the player's heel at floor level and registering the maximal height achieved during one calf heel raise with a metric ruler.


The player is thereafter instructed to perform repeated maximum unilateral heel raises until failure, guided by a metronome to standardize the pace (1 second concentric-, 1 second eccentric contraction). The player is allowed to have light contact with her fingers against a wall. A repetition is considered approved on the basis whether knee extension is maintained, and the reference mark on the player's heel levels with the registered maximal PF ROM height on the ruler. The test leader registers the total number of approved repetitions and discontinues the test when the player fails to reach the marked maximal height. The same procedure is then conducted on the opposite foot. The order of execution is randomized prior to the test.



**Figure S1.** Calf heel raises.

### **Active plantarflexion mobility**

Active PF ROM is measured with a clear plastic goniometer positioned at the lateral malleolus, utilizing fibula and fifth metatarsal as reference marks<sup>47 48</sup>. The player is positioned in supine on a treatment table, with feet off the edge of the table. The player is instructed to perform a sequence of six maximal active PF cycles starting from a neutral dorsiflexion (DF) position, whilst maintaining extended knees throughout the movement. The test leader measures and registers the maximal PF ROM in the final three cycles.



**Figure S2.** Active plantarflexion mobility execution. a) starting position in neutral dorsiflexion, b) end position in maximal active plantarflexion.

### **Weight bearing ankle dorsiflexion mobility**

Weight bearing ankle DF ROM is measured in a standing lunge position with the player's foot placed upon a metric ruler 10 cm away from a wall to the player's greater toe<sup>46 49</sup>. The player is instructed to lunge forward, directing the knee in line with her second toe, until contact with the wall is achieved; without allowing the heel to lift off the ground, which is continuously monitored through the availability to maintain a piece of paper against the foundation. Throughout the test, the player is allowed to provide balance by light contact with her fingers against the wall.

Firstly, three consecutive warm-up trials are performed from the 10 cm mark to familiarize the player with the test. Thereafter, the test leader measures the following three trials. In each trial, the player begins from the reference mark (10 cm) and progresses 1 cm away from the wall at a time, until unable to perform a successful repetition. If the player is unable to perform an approved repetition at the 10 cm reference mark, she is asked to progress 1 cm forward until able to complete a successful repetition. Once the player achieves knee-wall contact, the DF ROM is measured with a digital inclinometer (Clinometer, Plaincode, Stephanskirchen, Germany) and the distance from the wall to the greater toe is measured in cm in the repetition furthest away from the wall in each trial.



**Figure S3.** Weight bearing ankle dorsiflexion mobility.


### **Trunk mobility**

Mobility in trunk rotation are measured in a cross-legged seated position, and in a lunge position with the player on a gym mat, graded with 5 degrees increments, from zero to one hundred and eighty degrees<sup>50-52</sup>.


In the seated test (modified seated rotation test), the player is positioned at the center of the gym mat, in a cross-legged position with a wooden stick resting on the shoulders whilst keeping her arms crossed. If the player is unable to achieve the cross-legged sitting position, she is allowed to sit comfortable in an ordinary sitting position, which is noted by the test leader. Once in the starting position, the player is instructed to keep an upright posture and maximally rotate alternating between right and left for three times, whilst the test leader measures the rotational degrees in the end range.

The same procedure is thereafter repeated in a lunge position with the wooden stick resting on the player's shoulders. The player is positioned in a lunge position with her posterior knee at the center of the gym mat, and with her feet aligned on the zero-degree mark. Three consecutive maximal rotations are carried out alternating between right and left and is conducted in a lunge position for both the dominant, and non-dominant limb.






**Figure S4.** Modified seated rotation test. a) starting position, b) end position (right).



**Figure S5.** lunge rotation test. a) starting position left leg, b) end position (right).



**Figure S6.** lunge rotation test. a) starting position right leg, b) end position (left).

### ***Trunk strength***

Isometric trunk rotational strength is measured in a modified standing wood chopper test utilizing a force gauge to evaluate force output (RS Pro Digital Force Gauge, RS Components Ltd., Corby, UK) <sup>53-55</sup>.

In a standing position with extended arms, the player holds a handle in shoulder height, which is attached to the force gauge. The test leader positions the player in a 30-degree trunk rotation in the horizontal plane towards the anchor point (see figure S7).

The player is thereafter instructed to maximally generate force through her trunk and isometrically rotate in the opposite direction for five seconds whilst maintaining straight arms. Three consecutive repetitions are conducted for both right and left, and the maximal force output generated is used in the analyses. The order of execution is randomized prior to performing the test.



**Figure S7.** Modified standing wood chopper test (isometric rotation to the right).

### Deep neck flexor endurance

Deep neck flexor muscle endurance is assessed through a modified version of the Cranio-cervical flexion test (CCFT) with a pressure sensor (Stabilizer Pressure Bio-Feedback, Chattanooga Group inc, Hixson, TN)<sup>51 56 57</sup>.

Prior to executing the test, the player is instructed in how to perform a correct cranio-cervical flexion motion in standing and supine position through a gentle ‘head nodding’ cue. The player is positioned in supine position on a treatment table with her hands placed upon her abdomen or at the side of the body and with her feet on the table, with flexed hips and knees. With the player’s head and neck in a neutral position, the pressure stabilizer is positioned sub-occipitally, and inflated to a baseline pressure of 20 mmHg. Firstly, a pre-test is conducted and later an endurance test.

During the pre-test, the player is instructed to perform a gentle cranio-cervical flexion to increase the pressure starting from a baseline of 20 mmHg with 2 mmHg increments to a maximum of 30 mmHg. 3x3 second contractions are carried out at each target pressure (TP) with a three second rest in between each contraction whilst the test leader monitors for potential compensational strategies: excessive use of global neck musculature, chin jerking, cervical spine retraction, jaw clenching, breath holding and a pressure loss of  $\geq 2$  mmHg. A stopwatch time the contractions and visual feedback of pressure level is provided by the test leader who holds the manometer dial so that both the player and the test leader can read it throughout the procedure.

The endurance test is conducted if the player completes each of the five TP (22, 24, 26, 28 and 30 mmHg) without exhibiting any of the compensational strategies and/or experiencing pain during the pre-test. During the endurance test, the same setup and procedure as in the



pre-test is carried out. The player is now instructed to hold each contraction at the TP for 3x10 seconds with a ten second rest in between contractions. The highest completed TP with a full set of 3x10 seconds contractions is registered by the test leader and later used for analysis.



**Figure S8.** Modified cranio-cervical flexion test.


### **Hip- and knee strength**

Isometric hip flexion, extension, adduction, and abduction strength as well as eccentric hip abduction and adduction and isometric knee extension strength are measured with a hand-held dynamometer (HHD) (MicroFet2, Hoggan Health Industries inc. West Jordan, UT, USA)<sup>58 59</sup>.

Prior to executing the strength tests, two submaximal isometric contractions in each direction are performed to familiarize the player with the procedures. Three isometric contractions with gradually increasing force output for five seconds, and three maximal eccentric contractions for three seconds are performed in the isometric and eccentric tests, respectively, with a 10 second rest in between each contraction. The maximal force output for each position is registered by the test leader and later used for analysis. The order of execution and starting side is randomized prior to performing the tests at the particular test station (see table S1).

#### *Isometric hip flexion strength*


The player is positioned in a seated position at the edge of an treatment table, with 90-degrees of hip- and knee flexion. The HHD is positioned two centimeters proximal to the patella, and are externally fixated with a belt, which is secured under the leg of the treatment table, limiting hip flexion movement. The player is instructed to perform three isometric contractions on each leg.



**Figure S9.** Isometric hip flexion strength (right hip).

*Isometric knee extension strength*

Seated in the same position as during the isometric hip flexion strength test, with a slightly extended knee joint, the HHD is positioned two centimeters proximal to the malleoli on the anterior aspects of the player's tibia and are externally fixated with a belt. The player is instructed to perform three isometric contractions on each leg respectively.



**Figure S10.** Isometric knee extension strength (right leg).

*Isometric hip extension strength*

With the player positioned in a prone position on a treatment table and with her feet off the edge of the table, the test leader externally fixates the HHD two centimeters proximal to the malleoli with a belt. Furthermore, the player is instructed to perform three maximal isometric contractions on each leg respectively



**Figure S11.** Isometric hip extension strength (left hip).

*Isometric hip abduction strength*

The player is positioned in a supine position on an treatment table, with the tested leg extended, and the non-tested leg flexed. The test leader positions and fixates the HHD two centimeters proximal to the lateral malleolus with a belt, which limits hip abduction movement. Thereafter, the player is instructed to perform three maximal isometric hip abductions, whilst the test leader measures the force output for both the left, and right side.





**Figure S12.** Isometric hip abduction strength (left hip).

*Isometric hip adduction*

Lying in the same position as during the isometric hip abduction the test leader places and fixates the HHD two centimeters proximal to the medial malleolus with a belt. Consequently, the player executes three maximal isometric hip adductions on the left and right side, whilst the test leader registers the force output.



**Figure S13.** Isometric hip adduction strength (right hip).

### *Eccentric hip abduction strength*


The player is in a side-lying position on an treatment table with the test leg extended, and the opposite leg flexed to 90 degrees in the knee- and hip joint, whilst a neutral hip position is maintained. The player is subsequently instructed to place the test leg in approximately 40 degrees of hip abduction, and the test leader places a HHD one centimeter proximally to the lateral malleolus. The test leader initiates the test by saying “push”, and when the player has built up a maximal isometric contraction, the test leader begins to apply a downward directed force with the HHD whilst the player resists eccentrically for five seconds. Three repetitions are carried out on both the right and leg left, and the maximal force output is later used for analysis.



**Figure S14.** Eccentric hip abduction strength. a) starting position (right hip), b) end position (right hip).

### *Eccentric hip adduction strength*

The player is positioned in the same manner as in the eccentric hip abduction strength test, with the tested leg extended, and the non-tested leg flexed in the hip- and knee joint. Thereafter, the player is instructed to place the test leg in a maximal adduction position, whereupon the test leader positions a HHD one centimeter proximally to the medial malleolus. The test is initiated when the test leaders says “push”, whereupon a downward directed force is applied with the HHD whilst the player resists eccentrically for five seconds. Three consecutive trials are conducted on both sides, and the test leader registers the force output.



**Figure S15.** Eccentric hip abduction strength. a) starting position (left hip), b) end position (left hip).

### ***Hip mobility***

Measures of passive hip ROM in flexion, extension, abduction, internal- and external rotation are obtained using a universal clear plastic goniometer<sup>60 61</sup>. Three consecutive measurements for each position are performed for both the dominant and the non-dominant leg and the mean value for each position is later used for analysis. If the same value is obtained during the first and second measurement for a particular movement, a third one is not performed. The order of execution (side and movement) is randomized prior to performing the measures.

### ***Passive hip flexion ROM***

The player is positioned in supine position on a treatment table. With the player's leg held in a 90-degree knee flexion, test leader 1 moves the player's leg into a passive hip flexion until a firm end feel is achieved, and a posterior pelvic tilt occurs. Once the end feel is achieved, test leader 2 places the center of the goniometer at the greater trochanter and aligns one of the goniometer's arms with the player's femur, and the other one horizontally with the treatment table to read the goniometer. Three consecutive measures are conducted on each hip.






**Figure S16.** Passive hip flexion ROM (left hip).

*Passive hip abduction ROM*

The player is in a supine position on an treatment table with extended legs. While palpating the player's ipsilateral anterior superior iliac spine, test leader 1 holds the player's leg by the ankle and moves the leg into passive hip abduction until a firm end feel is achieved, and motion is felt at the pelvis. Thereafter, test leader 2 positions the goniometer at the player's hip, aligning the lever arms with the player's anterior superior iliac spine and femur, and reads the degrees of abduction. The test is repeated three times on each hip.



**Figure S17.** Passive hip abduction ROM (left hip).

### *Passive hip extension ROM*

In prone position with extended legs, test leader 1 fixates the player's pelvis by placing a hand at the ipsilateral posterior superior iliac spine. Thereafter, while holding the player's leg at the knee, test leader 1 moves the player's leg into passive hip extension, until an end feel is achieved, indicated by an anterior tilt of the pelvis. Test leader 2 measures the degrees of passive hip flexion with the goniometer's center positioned at the greater trochanter, and the lever arms in line with the player's femur and the treatment table horizontally. Three measures are performed on each leg.



**Figure S18.** Passive hip extension ROM (right hip).

### *Passive hip internal- and external rotation ROM*

In prone position, the player's leg is flexed to 90 degrees in the knee joint. Consequently, test leader 1 fixates the pelvis by placing his/her hand on the player's posterior superior iliac spine and performs a passive internal and external hip rotation, respectively, until an end feel is felt, indicated by an anterior pelvic tilt. Test leader 2 measures the degree of rotational mobility with a goniometer positioned at the knee, with the levers aligned with the player's tibia and with the treatment table horizontally. Three consecutive measures are conducted on each leg.




**Figure S19.** Passive hip rotational ROM (right hip). a) internal rotation, b) external rotation.

### ***Jump performance tests***

To assess the player's unilateral jump performance, the One-leg Long Box Jump Test (OLLBJ) and square hop test are performed<sup>62 63</sup>. A 40x40 cm square is marked on the foundation and later utilized as a reference mark in both tests. During the jump tests, players wear indoor sporting shoes.

#### ***One-leg long box jump test (OLLBJ)***


Firstly, the starting position, i.e. the distance player's jump from to the 40x40 cm square is calculated by dividing the player's height in cm with 1.6 ( $\text{height}/1.6 = \text{distance to the square}$ ). The player is instructed to stand on one leg at the starting position, and to perform a one-legged jump aiming inside the boundaries of the 40x40 cm square, and to maintain balance after landing. A trial is considered approved on the basis that the player land inside the 40x40 cm square, and adequately maintains balance after landing. The player performs three warm up trials on each leg, to familiarize with the procedure, and later five consecutive test trials. The test leader registers the total number of approved trials on each leg (0 to 5).



**Figure S20.** One-leg long box jump test (right leg). a) starting position, b) landing, c) balance maintained.

#### ***Square hop test***

During the square hop test, the player is instructed to hop on one leg in and out of the 40x40 cm square as many times as possible for 15 seconds in a clockwise direction, timed with a stopwatch, whilst the test leader registers the number of approved hops. A hop is classified as approved on the basis whether the player begins a hop in the starting position (outside the square) and then executes the short hop task inside the square and then in the correct direction outside the square. Prior to the test, the player performs two warm up trials on each foot.



**Figure S21.** A series of square hop tests illustrated on the player's right leg.

### ***Ankle- & knee stability***

#### ***Modified anterior drawer test (ankle)***

To assess talocrural stability or pain, a modified anterior drawer test is performed<sup>64 65</sup>. With the player in supine position with the test limb in knee flexion and the on the treatment table, the test leader applies an anteriorly directed force to the player's talus and a concurrent posteriorly directed force to the calcaneus. The test is conducted once on both the dominant and non-dominant foot and are considered positive if the player experiences any pain or discomfort during the procedure.





**Figure S22.** Modified anterior drawer test (right ankle).

*Modified Fairbank's apprehension test (patellofemoral)*

A modified version of Fairbank's apprehension test is conducted to evaluate stability or pain in the patellofemoral joint<sup>66</sup>. In supine position with extended legs, the test leader applies a laterally and subsequently medially directed force to the patella. The test is considered positive if the player experiences any pain or discomfort during the test, and/or an involuntary contraction of the quadriceps musculature. The test is carried out once on the player's dominant and non-dominant limb.



**Figure S23.** Modified Fairbank's apprehension test (left patellofemoral joint).  
a) lateral translation, b) medial translation.

### ***Isometric back extensor endurance***

Isometric back extensor endurance is assessed by a modified Sorensen test<sup>67-69</sup>. In prone position, the player's anterior superior iliac spine is positioned at the edge of the treatment table. The player's lower body is supported to the treatment table with three straps positioned over the player's ankles, knees, and pelvis. Whilst the test leader fastens the player's lower body to the treatment table with the three straps, the player uses a box/stool for support.

The player is thereafter instructed to keep her arms folded across the chest and isometrically maintain the upper body in a horizontal position until failure whilst the test leader register the time elapsed. A digital inclinometer (Clinometer, Plaincode, Stephanskirchen, Germany) is placed upon a metric ruler at the level of the 5<sup>th</sup> vertebra of the thoracal spine to monitor sagittal plane movement. If the player's upper body deviate greater than 10 degrees in the sagittal plane on more than two occasions and/or experience pain during the procedure, the test is stopped. Prior to the test, the player completes a shorter warmup trial of 5 seconds to orient the desired sagittal plane target angle.



**Figure S24.** Modified Sorensen test.

**Table S1.** Test stations, number of test leaders and randomization of the physical test protocol.

Test	Test station	Number of test leaders	Randomized order of execution
Calf heel raises	1	1	Yes
Active plantarflexion mobility	2	1	No
Weight bearing ankle dorsiflexion mobility	2	1	No
Ankle- & Knee stability	2	1	No
Hip mobility	3	2	Yes
Isometric Knee extension, hip flexion & extension strength	4	2	Yes
Trunk mobility	5	1	No
Trunk strength	5	1	Yes
Isometric and eccentric hip abduction and adduction	6	2	Yes
Deep neck flexor endurance	7	1	n/a
Functional performance tests	8	1	Yes
Isometric back extensor endurance	9	1	n/a

n/a-not applicable