

BMJ Open Interventions to reduce post-traumatic stress disorder symptoms in health care professionals from 2011 to 2021: a scoping review

Jialu Qian,¹ Weihong Wang,² Shiwen Sun,³ Lu Liu,¹ Yaping Sun,¹ Xiaoyan Yu ³

To cite: Qian J, Wang W, Sun S, *et al.* Interventions to reduce post-traumatic stress disorder symptoms in health care professionals from 2011 to 2021: a scoping review. *BMJ Open* 2022;**12**:e058214. doi:10.1136/bmjopen-2021-058214

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-058214>).

Received 20 October 2021
Accepted 05 January 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹School of Nursing, Zhejiang University School of Medicine, Hangzhou, China

²Department of Obstetrics, Ninghai Maternal and Child Health Hospital, Ningbo, China

³Department of Obstetrics, Women's Hospital School of Medicine Zhejiang University, Hangzhou, Zhejiang, China

Correspondence to
Professor Xiaoyan Yu;
yuxy@zju.edu.cn

ABSTRACT

Objectives This scoping review aimed to describe available interventions for decreasing (post-traumatic stress disorder) PTSD symptoms among healthcare professionals in hospital care.

Method A scoping review was conducted following Arksey and O'Malley's framework. PubMed, EMBASE, Cochrane Library, CINAHL, PsycINFO, Web of Science, Scopus and ProQuest were searched for original research published in English from 2011 to 2021, on 8 July 2021. We included studies that described interventions that focused on reducing the PTSD symptoms of healthcare professionals. A narrative synthesis was adopted to synthesise the data.

Results A total of eight studies out of 2558 articles were identified. Six used a quantitative study design and two adopted qualitative methods. cognitive behavioural therapy and mindfulness-based interventions were the most commonly adopted. Most studies used a combination of different intervention strategies. Trauma-related knowledge, emotion regulation and relaxation skill training, and psychological support from peers and psychologists were three core intervention components. The duration ranged from 2 weeks to 6 months. Healthcare professionals who participated in training programmes reported both positive experiences and suggestions for the improvement of PTSD-reducing interventions in their qualitative feedback.

Conclusions The scoping review provides a practical summary of the intervention characteristics for reducing the PTSD symptoms of healthcare professionals. Hospitals and managers could use the overview of interventions to assist healthcare professionals with PTSD symptoms. More research investigating the effects of PTSD symptom-reducing interventions for healthcare professionals with appropriate follow-up assessments is needed in the future.

INTRODUCTION

Post-traumatic stress disorder (PTSD) is a mental disorder that may occur following exposure to exceptionally horrifying events. It can develop after a single traumatic event or from long-term exposure to trauma.¹ Four main symptoms characterise PTSD: re-experiencing the event, avoidance of reminders, negative emotions related to the event and

Strengths and limitations of this study

- To the best of the researchers' knowledge, this is the first scoping review on interventions for reducing the post-traumatic stress disorder symptoms of healthcare professionals.
- This review was rigorous and based on the Preferred Reporting Items for Systematic Reviews and Meta-Analysis extension for Scoping Reviews checklist.
- We only searched English electronic databases, which means that some articles in this area published in languages other than English may have been missed.

chronic hyperarousal symptoms.² PTSD has a close relationship with substantial psychosocial morbidity,^{3 4} increased risk of suicide⁵ and poor physical health outcomes.^{3 6}

It has been reported that 3.0% of adults screen positive for PTSD at any time.⁷ PTSD occurs in approximately 8% of men and 20% of women after exposure to traumatic events.⁸ The estimated lifetime prevalence of PTSD is 8.3%.⁹ Due to occupational specialty, healthcare professionals have more opportunities for indirect or direct exposure to trauma by providing care to patients, and it has the potential to induce the symptomatic responses of PTSD.^{10 11} PTSD symptoms have been reported by a variety of healthcare professional groups, including physicians,^{12 13} nurses^{14 15} and midwives.^{16 17} The respective prevalence of PTSD in ambulance personnel, emergency physicians and midwives were 11%, 22% and 36%, respectively.^{18–20}

Healthcare professionals are facing the dilemma of PTSD symptom distress and insufficient organisational support. It has been reported that a stressful work environment and lack of sufficient organisational support could be antecedents of PTSD.^{11 15} Healthcare professionals described their experiences of trauma and PTSD as feeling alone

and unsupported and lacking emotional support.^{21 22} This may lead to adverse consequences among healthcare professionals, such as burnout, job dissatisfaction and the intention to leave their job.¹⁵ It is also likely to cause other mental disorders, including depression²³ and secondary stress disorder.²⁴ The poor mental states of healthcare professionals may increase the risk of medication errors and lower the quality of care provided for patients.^{25–27}

Only one previous meta-analysis examined the efficacy of psychological interventions for healthcare professionals with PTSD in a life-threatening pandemic.²⁸ This review paid more attention to the effectiveness of psychological interventions and included only one study in each meta-analysis. Studies of interventions conducted in healthcare professionals focusing on reducing PTSD symptoms are accumulating, especially amid the COVID-19 pandemic. Mindfulness approaches,²⁹ resilience training,³⁰ online three good things intervention³¹ and so on were conducted in this field. Considering that the intervention types and study designs varied, it is more appropriate to perform a scoping review, as it could give a general map of the interventions in this field. To the best of our knowledge, this is the first scoping review on interventions for reducing the PTSD symptoms of healthcare professionals. Previous studies demonstrated the importance of managing healthcare professionals' PTSD^{32–34}; therefore, it is meaningful to provide a comprehensive review in this area, thereby enhancing their psychological well-being and the quality of health services. This scoping review focused on intervention details (eg, intervention type, content, format, duration, frequency) to present a much-needed and broad overview of current interventions proposed to reduce the PTSD symptoms of healthcare professionals and to show gaps in published research that can be further explored.

METHODS

A scoping review was chosen, as this methodology is ideal for mapping the field of study's extent based on the review question's broad nature.³⁵ It facilitates information collection from different sources and study designs concerned with various research questions.³⁶ This scoping review was conducted based on a five-step methodological framework proposed by Arksey and O'Malley.³⁶ The Preferred Reporting Items for Systematic Reviews and Meta-Analysis extension for Scoping Reviews (PRISMA-ScR) checklist was followed.³⁷

Stage 1: identify the research question

This review sought to answer the following research question: What interventions are proposed to reduce the level of PTSD symptoms of healthcare professionals in hospital care?

Stage 2: identify the relevant studies

A systematic search was undertaken in PubMed, EMBASE, Cochrane Library (CENTRAL trials database), CINAHL,

PsycINFO, Web of Science, Scopus and ProQuest for articles published from 2011 until July 2021 to obtain the latest evidence in this area. We identified the main concepts, including 'healthcare professionals', 'intervention' and 'PTSD'. A combination of Medical Subject Headings/Emtree terms and free terms was used following the population, intervention, outcome scheme. The different search strings adapted for each database are displayed in online supplemental file 1. Two independent reviewers (JQ and WW) performed the search process following the same strategies. Reference lists of included articles were reviewed thoroughly to search for additional studies.

Stage 3: study selection

Studies were selected based on the inclusion and exclusion criteria. The inclusion criteria were as follows: (a) research studies conducted among healthcare professionals who have experiences of exposure to trauma in the context of the hospital; (b) original studies focusing on the evaluation of the effectiveness or experiences of any intervention with the primary or secondary aim of reducing healthcare professionals' PTSD symptoms; (c) PTSD symptoms were evaluated using valid measurements, such as the Impact of Event Scale—Revised (IES-R) and PTSD Symptom Checklist (PCL-C); (d) qualitative, quantitative or mixed methods studies and (e) published in English between 2011 and 2021. Restrictions on the date were made to concentrate on recent clinical practice and to update previous literature. This review focused on primary research studies; therefore, conference abstracts, reviews, news, study protocol commentaries, case reports, letters, guidelines and books were excluded. Two reviewers (JQ and WW) independently determined the eligibility of studies using a two-step screening process: (1) titles and abstract screening and (2) full-text screening. Any discrepancy was resolved by discussion with a third reviewer (SS).

Stage 4: charting the data

The research team designed a standardised table, and pilot testing was performed on two articles. A consensus was reached to extract data that answered the review question.³⁸ The extracted data included the first author, year of publication, country, study design, study aim, participants, sample size, intervention details (type, duration, frequency and so on), PTSD measurement and results. The first author carefully read and assessed each identified article, which another reviewer (WHW) double checked to ensure accuracy. We did not undertake a systematic quality appraisal of the included studies because it was a scoping review rather than a systematic review.³⁹

Stage 5: collating, summarising and reporting the results

A narrative synthesis was adopted to collate the extracted data.⁴⁰ The Synthesis Without Meta-analysis guideline was followed.⁴¹ Groupings of intervention types, strategies, content, format, duration, session, frequency, assessment




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analysis flow diagram illustrating the search process. PTSD, post-traumatic stress disorder.

time and effectiveness of the interventions were used in the synthesis. We used descriptive tables to synthesise and compare the broad findings. Similarities and differences between the identified studies were compared with generate common themes. Research team members participated in the discussion and revision of the results and assisted in validating the theme identification and description throughout the process.

Patient and public involvement

There was no direct patient or public involvement in this review.

RESULTS

Search outcomes and study characteristics

Electronic searches and other sources identified 2558 citations, resulting in 2154 unique citations to be screened by title and abstract following the removal of 404 duplicates. Overall, 2126 publications were excluded based on title and abstract screening, which left 28 citations for full-text review. Ultimately, eight studies were identified as focusing on reducing the PTSD symptoms of healthcare professionals.^{29–31 42–46} A PRISMA flow diagram illustrating the search process is shown in [figure 1](#).

Details on the study characteristics are displayed in [table 1](#). These studies were conducted in a variety of countries, including the USA (n=3)^{30 31 43} and the UK (n=2),^{45 46} with one study each from Australia (n=1),⁴² Spain (n=1)²⁹ and Mexico (n=1).⁴⁴ The publication year ranged from

2012 to 2021. Of the eight empirical publications, four used a randomised controlled trial design,^{29 30 42 44} two reported qualitative methods,^{31 46} one was a pre-post pilot test study⁴³ and one was a longitudinal design.⁴⁵ The included studies were mainly conducted among healthcare professionals working in a relatively stressful environment, such as the intensive care unit, emergency room, palliative care unit or maternity unit. A total of 859 participants were included in this review. The sample size in each study ranged from 21 to 482. In quantitative studies, the Clinician-Administered PTSD Scale, Davidson Trauma Scale, Post-traumatic Diagnostic Scale (PDS), PCL-C, and IES-R were used to measure PTSD symptoms.

Intervention types and strategies

See online supplemental file 2 for details on interventions for reducing PTSD in healthcare professionals and online supplemental file 3 for a summary of the intervention characteristics. Five studies used a combination of different intervention strategies^{29 30 43 45 46} and three studies^{31 42 44} used a single intervention. Cognitive behavioural therapy (CBT)^{29 30 42 45 46} was the most commonly adopted intervention. Four studies conducted mindfulness-based interventions.^{29 30 43 44} One study used the three good things intervention.³¹

Among the eight included studies, various strategies were used. CBT interventions were chosen for psychoeducation, emotion regulation training, exposure therapy and event-triggered counselling. Mindfulness-based interventions usually used relaxation exercises through breathing or imagination, stretching and deep breathing exercises and mindfulness-based stress reduction exercises (eg, body scan and sitting meditation). Four studies^{30 43 45 46} employed a group workshop. Writing therapy, including the three good things intervention and written exposure,^{30 31} was applied. Two studies provided available referral and access to psychological assessment from a clinical psychologist.^{45 46}

Intervention content and format

Interventions on reducing the PTSD symptoms of healthcare professionals mainly contained the following three core components: (1) trauma-related knowledge: giving theoretical guidance to help healthcare professionals understand trauma reactions and management; (2) emotion regulation and relaxation skill training and (3) psychological support from peers and psychologist: strengthening the feelings of support of healthcare professionals to express trauma experiences and offering a professional psychological assessment.

Six of eight studies were conducted offline,^{30 42–46} and only two studies used a mobile-based intervention and an online intervention.^{29 31}

Duration, session and frequency

The duration of intervention ranged from 2 weeks to 6 months. Generally, the intervention session varied from 5 to 16. The length of each session ranged from 15 to

Table 1 Characteristics of included studies (n=8)

Author & year (Country)	Study design	Study aims	Participants	Sample size	PTSD measurement
Bryant, 2019 (Australia) ⁴²	RCT	To assess the efficacy of cognitive behavioural therapy (CBT) for PTSD in emergency service personnel and examined if brief exposure to trauma memories is no less efficacious than prolonged exposure.	Emergency service personnel with PTSD	100	CAPS
Fiol-DeRoque, 2021 (Spain) ²⁹	RCT	To evaluate the effectiveness of a mobile phone-based intervention to reduce mental health problems in healthcare workers during the COVID-19 pandemic.	Healthcare workers providing face-to-face healthcare to patients with COVID-19	482	DTS
Gerhart, 2016 (USA) ⁴³	A pre–post pilot test	To examine the efficacy of mindfulness-based communication training on depression, burnout and PTSD symptoms.	Palliative care providers	21	PCL-C
Kim, 2012 (Mexico) ⁴⁴	RCT	To determine whether mindfulness-based stretching and deep breathing exercise reduces PTSD symptom severity among nurses with subclinical features of PTSD.	Nurses	22	PCL-C
Mealer, 2014 (USA) ³⁰	RCT	To determine if a multimodal resilience training programme for ICU nurses was feasible to perform and acceptable to the study participants.	ICU nurses	27	PDS
Rippstein, 2017 (USA) ³¹	A qualitative study using thematic analysis	To explore the content of the good things reported by healthcare workers participating in the Three Good Things intervention, which were used to strengthen healthcare workers' skills to deal with stressful events	Neonatal ICU healthcare professionals	32	/
Slade, 2018 (UK) ⁴⁵	Longitudinal design	To identify the programme for the prevention of PTSD in midwifery on their psychological well-being and job satisfaction.	Midwives	153	IES-R
Slade, 2020 (UK) ⁴⁶	A qualitative interview study	To examine the perceived acceptability, utility and relevance of the POPPY resources.	Midwives and midwifery managers	22	/

CAPS, Clinician-Administered PTSD Scale; CBT, cognitive behavioural therapy; DTS, Davidson Trauma Scale; ICU, intensive care unit; IES-R, Impact of Event Scale – Revised; PCL-C, PTSD Symptom Checklist; PDS, Post-traumatic Diagnostic Scale; POPPY, programme for the prevention of PTSD in midwifery; RCT, randomised controlled trial.

120 min. The frequencies were weekly,^{30 42 43} semiweekly⁴⁴ or daily.^{29 31}

Assessment time and effectiveness of the interventions

In six quantitative studies,^{29 30 42–44 46} all the studies conducted preintervention and postintervention assessments. One study added a mid-training assessment.⁴³ Additionally, only one study reported follow-up assessment scores at 9 months post intervention.⁴² Significantly improved symptoms of PTSD were observed in quantitative studies at postintervention assessment.^{29 30 42–44 46} The effectiveness of CBT in reducing PTSD was still maintained at the 9-month follow-up.⁴²

Findings from qualitative studies

There were two studies describing participants' experiences after the PTSD symptom-reducing interventions.

Benefits of the PTSD symptom-reducing interventions

Two studies both obtained positive feedback from participants. In the study that conducted the three good things intervention,³¹ participants reported promoted well-being and resilience. They proposed that good teamwork, supportive coworkers, communication and relaxing sleep were important factors contributing to their well-being. Another qualitative study interviewed midwives and midwifery managers.⁴⁶ Midwives reported that this training raised awareness of the impact of trauma and made them feel more informed about managing their trauma responses after the intervention. The availability of trusted peer supporters and access to psychological assessments was helpful. From the point of midwifery managers, they reported that the programme enabled acknowledgement and management of trauma in midwifery. Midwifery managers thought the programme filled a clinical gap for midwives seeking support pathways after workplace trauma.

Suggestions and challenges of PTSD symptom-reducing interventions

Midwives reported uncertainty, fear of judgement and stigma as barriers to accessing peer support. Midwives suggested peer supporters should have sufficient clinical experience to fully understand the situation being discussed. For access to psychological support, easy access and confidentiality were essential. Midwifery managers thought the programme should be integrated into the organisational context and require financial investment from the organisation. It could be mandatory training and applied among the wider maternity workforce.

DISCUSSION

The objective of this scoping review was to describe the range of interventions designed to reduce PTSD symptoms among healthcare professionals. This review synthesises intervention settings and the available quantitative and qualitative results regarding the effectiveness of interventions, thereby providing a basis for applying interventions

for healthcare professionals' PTSD symptom reduction. Eight studies were identified in our scoping review. Considering the high prevalence rate of PTSD among healthcare professionals⁴⁶ and the numerous adverse consequences caused by it,^{24 26 27 47} the importance of this scoping review becomes more apparent.

Included studies all showed positive effects in reducing PTSD symptoms in healthcare professionals. In terms of intervention types and strategies, CBT and mindfulness-based interventions were commonly used. This might be explained by previous research which found CBT and mindfulness-based interventions to be effective for PTSD treatment in previous studies.^{48 49} One study focused on resilience training.³⁰ Resilience has been implicated a positive and dynamic outcome following stressors such as traumatic events.⁵⁰ It protects individuals against stress-related mental problems.⁵¹ Interventions focused on enhancing resilience can help prevent and treat individuals with PTSD.^{52 53} The 'three good things' writing therapy showed benefits in relieving PTSD,³¹ which is consistent with previous studies.^{54–56} With the development of positive psychology, many studies have tried to improve PTSD symptoms by increasing positive psychological changes, which may help individuals re-establish a new understanding of life.^{57 58} Group format workshops tend to be used because they can help share useful information on trauma management and help them gain sufficient peer support and organisational support.^{30 43 46}

Trauma-related knowledge, emotion regulation and relaxation skill training, and psychological support from peers and psychologists were summarised as three core intervention components. On the one hand, the knowledge and skills of PTSD management should be improved by providing training to healthcare professionals. On the other hand, there is a need to establish a communication platform to provide adequate psychological and professional support for providers.²² Comprehensive PTSD symptom-reducing interventions based on these three core components could be referred in clinical implementation.

In regard to intervention format, two studies used a mobile-based intervention and an online intervention.^{29 31} In the era of rapid electronic technology development, requirements for healthcare services have promoted instant communication and accelerated efficiency in the transmission of information.⁵⁹ Compared with offline interventions, it might be more convenient to offer long-term support for healthcare professionals via internet technology⁶⁰ or smartphones.⁶¹

Regarding findings from qualitative studies, healthcare professionals gave their positive feedback, which illustrated that these interventions might be effective in PTSD symptom reduction. Meanwhile, suggestions and challenges of PTSD symptom-reducing interventions were proposed. Good teamwork and supportive coworkers were considered important to their mental health. However, the reality is that healthcare professionals feel that support from colleagues and managers is insufficient

after experiencing traumatic events.^{15 21 62 63} Our findings indicated that fear of judgement and stigma were barriers for healthcare professionals to access peer support. This finding is similar to previous findings that some healthcare professionals suffered stigma from their colleagues and managers.⁶⁴ Organisational support from the hospital is insufficient⁴⁶; therefore, hospitals should first attach full importance to healthcare professionals' PTSD management and make clear policies. PTSD symptom screening should be provided for healthcare professionals so that early intervention can be given, thereby decreasing negative psychological outcomes.⁶⁵ The availability of peer support and psychological assessments were viewed as useful resources for healthcare professionals.⁴⁶ Hospitals and managers should try to establish a comprehensive support system for PTSD management, thereby providing strong backing for healthcare professionals.

Only a small number of studies describing PTSD symptom-reducing interventions for healthcare professionals were included in this scoping review, which suggested that little attention has been given to the psychological well-being of healthcare professionals. More research should investigate the effects of PTSD symptom-reducing interventions for healthcare professionals in the future, given inadequate research in this area. Due to the combination of diverse intervention techniques, future studies would be needed to clearly determine the effectiveness of respective strategies. Internet technology or smartphones could be integrated into interventions for convenient and long-term support. Only one study reported follow-up assessment scores.⁴² Research should have appropriate follow-up times to comprehensively understand the maximal efficacy of PTSD severity in the future.

Strengths and limitations

A strength of this scoping review is that it provides a broad overview of interventions aimed at reducing the PTSD symptoms of healthcare professionals. It summarised detailed intervention characteristics and components, which could provide guidance for applying PTSD symptom-reducing interventions for healthcare professionals. This review adheres to the PRISMA-ScR checklist, which ensures the quality and rigour of the study.

There are several limitations of this review. This review has not been preregistered. And also, although we searched eight databases and thoroughly reviewed reference lists of identified studies, we only included eight studies in this review. Therefore, to some extent, this review contributed limited information. Another limitation is that this study is limited to papers published in the English language, which means that some articles in this area may have been missed. Finally, a quality assessment of the included research was not performed considering the scoping review methodology. It is not always necessary for a scoping review. Nevertheless, a scoping review aims to give a general map of the research on a general topic to

inform the reader and to clarify the directions for further research,⁶⁶ which has been achieved in our study.

Conclusions

The scoping review provides a useful summary of the intervention characteristics for reducing the PTSD symptoms of healthcare professionals. This review highlights that hospitals and managers could use the overview of interventions and strengthen support for healthcare professionals' PTSD symptoms. Due to inadequate research in this area, more research should investigate the effect of PTSD symptom-reducing interventions for healthcare professionals with appropriate follow-up assessments in the future.

Contributors JQ, WW, SS, LL, YS and XY all contributed to the conception and design of the review. JQ and WW independently searched electronic databases and screened titles and abstracts. JQ and WW were responsible for reviewing the included studies. Disagreements were discussed with a third researcher (SS) to reach a consistent conclusion. This paper was drafted by JQ and reviewed by SS, LL, YS and XY critically prior to submission; XY served as scientific advisor. XY is responsible for the overall content as guarantor.

Funding This work was supported by Medical and Health Research Project of Zhejiang Province (grant number 2020KY173).

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study does not involve human participants.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Xiaoyan Yu <http://orcid.org/0000-0003-3861-7340>

REFERENCES

- 1 Bisson JI, Cosgrove S, Lewis C, *et al*. Post-traumatic stress disorder. *BMJ* 2015;351:h6161.
- 2 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. Washington, DC: American Psychiatric Association, 2013.
- 3 Sommer JL, El-Gabalawy R, Mota N. Understanding the association between posttraumatic stress disorder characteristics and physical health conditions: a population-based study. *J Psychosom Res* 2019;126:109776.
- 4 Kessler RC, Sonnega A, Bromet E, *et al*. Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 1995;52:1048–60.
- 5 Gradus JL. Posttraumatic stress disorder and death from suicide. *Curr Psychiatry Rep* 2018;20:98.
- 6 Gupta MA. Review of somatic symptoms in post-traumatic stress disorder. *Int Rev Psychiatry* 2013;25:86–99.

- 7 McManus S, Meltzer H, Brugha T. *Adult psychiatric morbidity in England 2007: results of a household survey*, 2009.
- 8 Warner CH, Warner CM, Appenzeller GN, et al. Identifying and managing posttraumatic stress disorder. *Am Fam Physician* 2013;88:827–34.
- 9 Kilpatrick DG, Resnick HS, Milanak ME, et al. National estimates of exposure to traumatic events and PTSD prevalence using DSM-IV and DSM-5 criteria. *J Trauma Stress* 2013;26:537–47.
- 10 Elwood LS, Mott J, Lohr JM, et al. Secondary trauma symptoms in clinicians: a critical review of the construct, specificity, and implications for trauma-focused treatment. *Clin Psychol Rev* 2011;31:25–36.
- 11 Schuster M, Dwyer PA. Post-Traumatic stress disorder in nurses: an integrative review. *J Clin Nurs* 2020;29:2769–87.
- 12 Lazarus A. Traumatized by practice: PTSD in physicians. *J Med Pract Manage* 2014;30:131–4.
- 13 Walker AL, Gamble J, Creedy DK, et al. Impact of traumatic birth on Australian obstetricians: a pilot feasibility study. *Aust N Z J Obstet Gynaecol* 2020;60:555–60.
- 14 Park E, Meyer RML, Gold JL. The role of medical specialization on posttraumatic symptoms in pediatric nurses. *J Pediatr Nurs* 2020;53:22–8.
- 15 Levi P, Patrician PA, Vance DE, et al. Post-Traumatic stress disorder in intensive care unit nurses: a concept analysis. *Workplace Health Saf* 2021;69:224–34.
- 16 Ravaldi C, Carelli E, Frontini A, et al. The BLOSSoM study: Burnout after perinatal loss in midwifery. Results of a nation-wide investigation in Italy. *Women and Birth* 2021;1. doi:10.1016/j.wombi.2021.01.003
- 17 McDaniel LR, Morris C. The second victim phenomenon: how are midwives affected? *J Midwifery Womens Health* 2020;65:503–11.
- 18 Petrie K, Milligan-Saville J, Gayed A, et al. Prevalence of PTSD and common mental disorders amongst ambulance personnel: a systematic review and meta-analysis. *Soc Psychiatry Psychiatr Epidemiol* 2018;53:897–909.
- 19 DeLucia JA, Bitter C, Fitzgerald J, et al. Prevalence of post-traumatic stress disorder in emergency physicians in the United States. *West J Emerg Med* 2019;20:740–6.
- 20 Beck CT, LoGiudice J, Gable RK. A mixed-methods study of secondary traumatic stress in certified nurse-midwives: Shaken belief in the birth process. *J Midwifery Womens Health* 2015;60:16–23.
- 21 Ryan T, McLeod N. Alone and unsupported: a mental health nurse's experience of trauma, PTSD, institutional response and resilience. *Int J Ment Health Nu* 2018;27:42.
- 22 Qian J-L, Pan P-E, Wu M-W, et al. The experiences of nurses and midwives who provide surgical abortion care: a qualitative systematic review. *J Adv Nurs* 2021;77:3644–56.
- 23 Hamama-Raz Y, Walker R, Palgi Y, et al. Comorbidity of post-traumatic stress symptoms and depressive symptoms among obstetric nurses with perinatal death exposure. *Isr J Psychiatry Relat Sci* 2016;53:58–62.
- 24 Roden-Foreman JW, Bennett MM, Rainey EE, et al. Secondary traumatic stress in emergency medicine clinicians. *Cogn Behav Ther* 2017;46:522–32.
- 25 Karanikola M, Giannakopoulou M, Mpouzika M, et al. Dysfunctional psychological responses among intensive care unit nurses: a systematic review of the literature. *Rev Esc Enferm USP* 2015;49:847–57.
- 26 Almutairi AF, Adlan AA, Balkhy HH. "It feels like I'm the dirtiest person in the world.": exploring the experiences of healthcare providers who survived MERS-CoV in Saudi Arabia. *J Infect Public Health* 2018; 11:187–91.
- 27 Harapan H, Alleta A, Anwar S, et al. Attitudes towards Zika virus infection among medical doctors in Aceh Province, Indonesia. *J Infect Public Health* 2018;11:99–104.
- 28 Sun Z, Yu C, Zhou Y, et al. Psychological interventions for healthcare providers with PTSD in life-threatening pandemic: systematic review and meta-analysis. *Front Psychiatry* 2021;12:697783.
- 29 Fiol-DeRoque MA, Serrano-Ripoll MJ, Jiménez R, et al. A mobile Phone-Based intervention to reduce mental health problems in health care workers during the COVID-19 pandemic (PsyCovidApp): randomized controlled trial. *JMIR Mhealth Uhealth* 2021;9:e27039.
- 30 Mealer M, Conrad D, Evans J, et al. Feasibility and acceptability of a resilience training program for intensive care unit nurses. *Am J Crit Care* 2014;23:e97–105.
- 31 Rippstein-Leuenberger K, Mauthner O, Bryan Sexton J, et al. A qualitative analysis of the three good things intervention in healthcare workers. *BMJ Open* 2017;7:e015826.
- 32 D'Ettoire G, Pellicani V, Ceccarelli G. Post-traumatic stress disorder symptoms in healthcare workers: a ten-year systematic review. *Acta Biomed* 2020;91:e2020009.
- 33 Chew NWS, Lee GKH, Tan BYQ, et al. A multinational, multicentre study on the psychological outcomes and associated physical symptoms amongst healthcare workers during COVID-19 outbreak. *Brain Behav Immun* 2020;88:559–65.
- 34 Carmassi C, Foghi C, Dell'Oste V, et al. PTSD symptoms in healthcare workers facing the three coronavirus outbreaks: what can we expect after the COVID-19 pandemic. *Psychiatry Res* 2020;292:113312.
- 35 Peters MDJ, Godfrey CM, Khalil H, et al. Guidance for conducting systematic scoping reviews. *Int J Evid Based Healthc* 2015;13:141–6.
- 36 Arskey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Method* 2005;8:19–32.
- 37 Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018;169:467–73.
- 38 Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci* 2010;5:69.
- 39 Munn Z, Peters MDJ, Stern C, et al. Systematic review or scoping review? guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol* 2018;18:143.
- 40 Barnett-Page E, Thomas J. Methods for the synthesis of qualitative research: a critical review. *BMC Med Res Methodol* 2009;9:59.
- 41 Campbell M, McKenzie JE, Sowden A, et al. Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline. *BMJ* 2020;368:l6890.
- 42 Bryant RA, Kenny L, Rawson N, et al. Efficacy of exposure-based cognitive behaviour therapy for post-traumatic stress disorder in emergency service personnel: a randomised clinical trial. *Psychol Med* 2019;49:1565–73.
- 43 Gerhart J, O'Mahony S, Abrams I, et al. A pilot test of a mindfulness-based communication training to enhance resilience in palliative care professionals. *J Contextual Behav Sci* 2016;5:89–96.
- 44 Kim S, Burge M. P02.137. Mindfulness-based stretching and deep breathing exercise reduces symptoms of posttraumatic stress disorder. *BMC Complement Altern Med* 2012;12.
- 45 Slade P, Sheen K, Collinge S, et al. A programme for the prevention of post-traumatic stress disorder in midwifery (POPPY): indications of effectiveness from a feasibility study. *Eur J Psychotraumatol* 2018;9:1518069–11.
- 46 Slade P, Sheen K, Collinge S. Acceptability of a programme for the prevention of post-traumatic stress disorder in midwifery: a qualitative investigation with midwives and midwifery managers following feasibility testing. *MIDIRS Midwifery Digest* 2020;30:27–32.
- 47 Colville GA, Smith JG, Brierley J, et al. Coping with staff burnout and work-related posttraumatic stress in intensive care. *Pediatr Crit Care Med* 2017;18:e267–73.
- 48 Lewis C, Roberts NP, Andrew M, et al. Psychological therapies for post-traumatic stress disorder in adults: systematic review and meta-analysis. *Eur J Psychotraumatol* 2020;11:1729633.
- 49 Goldstein E, McDonnell C, Atchley R, et al. The impact of psychological interventions on posttraumatic stress disorder and pain symptoms: a systematic review and meta-analysis. *Clin J Pain* 2019;35:703–12.
- 50 Kalisch R, Cramer AOJ, Binder H, et al. Deconstructing and reconstructing resilience: a dynamic network approach. *Perspect Psychol Sci* 2019;14:765–77.
- 51 Kalisch R, Baker DG, Basten U, et al. The resilience framework as a strategy to combat stress-related disorders. *Nat Hum Behav* 2017;1:784–90.
- 52 Rakesh G, Morey RA, Zannas AS, et al. Resilience as a translational endpoint in the treatment of PTSD. *Mol Psychiatry* 2019;24:1268–83.
- 53 Horn SR, Feder A. Understanding resilience and preventing and treating PTSD. *Harv Rev Psychiatry* 2018;26:158–74.
- 54 Qian J, Zhou X, Sun X, et al. Effects of expressive writing intervention for women's PTSD, depression, anxiety and stress related to pregnancy: A meta-analysis of randomized controlled trials. *Psychiatry Res* 2020;288:112933.
- 55 Thompson-Hollands J, Marx BP, Sloan DM. Brief novel therapies for PTSD: written exposure therapy. *Curr Treat Options Psychiatry* 2019;6:99–106.
- 56 van Emmerik AAP, Reijntjes A, Kamphuis JH. Writing therapy for posttraumatic stress: a meta-analysis. *Psychother Psychosom* 2013;82:82–8.
- 57 Zebrack B, Kwak M, Salsman J, et al. The relationship between posttraumatic stress and posttraumatic growth among adolescent and young adult (AYA) cancer patients. *Psychooncology* 2015;24:162–8.
- 58 Qian J, Sun S, Zhou X, et al. Effects of an expressive writing intervention in Chinese women undergoing pregnancy termination for fetal abnormality: a randomized controlled trial. *Midwifery* 2021;103:103104.



- 59 Khan S, Dasrath F, Farghaly S, *et al.* Unmet communication and information needs for patients with IBD: implications for mobile health technology. *Br J Med Med Res* 2016;12:1–11.
- 60 Bevan Jones R, Thapar A, Rice F, *et al.* A web-based psychoeducational intervention for adolescent depression: design and development of MoodHwb. *JMIR Ment Health* 2018;5:e13.
- 61 Chan KL, Leung WC, Tiwari A, *et al.* Using smartphone-based psychoeducation to reduce postnatal depression among first-time mothers: randomized controlled trial. *JMIR Mhealth Uhealth* 2019;7:e12794.
- 62 Mayers PM, Briony P, Beryl G. Experiences of registered midwives assisting with termination of pregnancies at a tertiary level Hospital. *Health Sa Gesondheid* 2008;10.
- 63 Mamabolo L, Tjallinks JE. Experiences of registered nurses at one community health centre near Pretoria providing termination of pregnancy services. *Africa J Nurs Mid* 2010;12:73–86.
- 64 Teffo ME, Levin J, Rispel LC. Compassion satisfaction, burnout and secondary traumatic stress among termination of pregnancy providers in two South African provinces. *J Obstet Gynaecol Res* 2018;44:1202–10.
- 65 Si M-Y, Su X-Y, Jiang Y, *et al.* Psychological impact of COVID-19 on medical care workers in China. *Infect Dis Poverty* 2020;9:113.
- 66 Estupiñán Artiles C, Regan J, Donnellan C. Dysphagia screening in residential care settings: a scoping review. *Int J Nurs Stud* 2021;114:103813.