


BMJ Open Transcutaneous auricular vagus nerve stimulation in poststroke cognitive impairment: protocol for a randomised controlled trial

Zhen-Dong Li,¹ Hang-Jian Qiu,² Xiao-Qian Wang,² Cheng-Cheng Zhang,¹ Yue-Juan Zhang ¹

To cite: Li Z-D, Qiu H-J, Wang X-Q, *et al.* Transcutaneous auricular vagus nerve stimulation in poststroke cognitive impairment: protocol for a randomised controlled trial. *BMJ Open* 2022;**12**:e063803. doi:10.1136/bmjopen-2022-063803

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-063803>).

Received 12 April 2022
Accepted 23 September 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Nursing, The First Affiliated Hospital of Hunan University of Chinese Medicine, Changsha, Hunan, China

²School of Nursing, Hunan University of Chinese Medicine, Changsha, Hunan, China

Correspondence to

Dr Yue-Juan Zhang;
zhangyuejuanhuli@163.com

ABSTRACT

Background As one of the most common stroke sequelae, poststroke cognitive impairment significantly impacts 17.6%–83% of survivors, affecting their rehabilitation, daily living and quality of life. Improving cognitive abilities among patients in stroke recovery is therefore critical and urgent. Transcutaneous auricular vagus nerve stimulation (TAVNS) is a non-invasive, safe, cost-effective treatment with great potential for improving the cognitive function of poststroke patients. This clinical research will evaluate the effectiveness, and help elucidate the possible underlying mechanisms, of TAVNS for improving poststroke cognitive function.

Methods and analysis A single-centre, parallel-group, allocation concealment, assessor-blinded randomised controlled clinical trial. We will allocate 88 recruited participants to the TAVNS or sham group for an intervention that will run for 8 weeks, 5 days per week with twice daily sessions lasting 30 min each. Blood tests will be performed and questionnaires issued at baseline and 8-week and 12 week follow-ups. Primary outcomes will be changes in cognitive function scores. Secondary outcomes will be changes in activities of daily living, quality of life and serum oxidative stress indicators.

Ethics and dissemination The Ethics Committee of the First Affiliated Hospital of Hunan University of Chinese Medicine has approved the protocol (No. HN-LL-YJSLW-2022200). Findings will be published in peer-reviewed academic journals and presented at scientific conferences.

Trial registration number ChiCTR2200057808.

INTRODUCTION

Among stroke sequelae, poststroke cognitive impairment (PSCI) is common, affecting 17.6%–83% of stroke survivors, depending on assessment timing, research setting, demographics and cognitive test/cut-off.¹ PSCI is a clinical syndrome characterised by cognitive deterioration after a clinical cerebrovascular incident, which persists for up to 6 months; it includes both PSCI no dementia and the more severe form, post-stroke dementia.^{2–3} Core impaired domains

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The first single-centre, parallel-group, assessor-blinded, randomised controlled clinical trial to test the effects of transcutaneous auricular vagus nerve stimulation (TAVNS) on poststroke cognitive impairment.
- ⇒ Integrates diverse outcomes to examine the effects and underlying mechanisms of TAVNS in poststroke cognitive improvement.
- ⇒ Adherence to out-of-hospital self-administered TAVNS treatments may influence study results.
- ⇒ Because stimulation site differences are obvious, incomplete patient and assessor blinding may limit study results.

include executive function, memory, attention, language and visuospatial function, which not only lead to disability, dependence and low quality of life,⁴ but are closely associated with high risk of recurrent ischaemic stroke⁵ and reduced 5-year survival.² As the tremendous burdens of stroke continue to rise,^{6–7} PSCI has become a mounting public healthcare challenge that urgently needs to be addressed.

Currently, the most relevant PSCI-prevention measures are acute therapy and recurrence prevention.¹ However, there are no unambiguously successful therapeutic techniques, and prospective pharmaceuticals have yet to show efficacy in decreasing or preventing cognitive deterioration following stroke. Acetylcholinesterase inhibitors (AChEIs) and memantine, which are approved for use in Alzheimer's disease (AD), have shown some clinical benefits in vascular dementia.^{8–9} However, a recent study revealed little evidence that pharmaceutical therapies, including AChEIs and memantine, help reduce symptoms or slow dementia progression in patients with PSCI.¹⁰ Pharmaceutical therapies also cause side effects like

gastrointestinal issues (diarrhoea, constipation), headaches and dizziness.¹¹ Thus, it is critical to develop viable treatments that slow or stop PSCI progression.

Vagus nerve stimulation (VNS) is approved by the United States Food and Drug Administration as an adjunct to medication for treating partial epilepsy, depression and primary headache disorders¹²; it was also recently recognised to have the potential for improving cognitive impairment.^{13 14} The vagus nerve connects to the nucleus of the solitary tract (NTS), which activates noradrenergic neurons in the locus coeruleus (LC). This causes norepinephrine (NE) release in memory-forming areas like the hippocampus, basolateral amygdala and medial prefrontal cortex.^{15–18} Because noradrenergic stimulation has been demonstrated to improve memory performance, the NTS-LC-NE pathway is thought to play a key role in memory enhancement, which may be activated with VNS.^{19–21} In clinical practice, there are two VNS types: invasive (iVNS) and non-invasive (nVNS). Interest in nVNS is greater, because iVNS necessitates a surgical procedure under general anaesthesia and is associated with potential treatment-related complications like cardiac arrhythmias, peritracheal haematomas and vocal cord dysfunction.²²

Transcutaneous auricular VNS (TAVNS) is a safe, well-tolerated, non-invasive alternative²³ that activates the vagal projections and vagally mediated pathways similar to iVNS.²⁴ The auricular branch of the vagus nerve (ABVN) is the only branch of the vagus to reach the body surface, at the cymba and cavum conchae.²⁵ Its afferent fibres pass through the cervical ganglion into the vagal trunk and project to the NTS.²⁶ Functional brain imaging has shown that TAVNS and iVNS have similar activation patterns, suggesting that the former effectively activates vagal trajectories and has the potential to induce similar effects.^{27 28} Clinical studies have also shown that TAVNS can effectively improve cognitive function in those with epilepsy²⁹ and depression,³⁰ and in healthy individuals.^{31 32} However, whether TAVNS has the same effects in those with PSCI is unclear.

PSCI has a complex set aetiologies, including dysfunctions of the neurovascular unit,³³ neuroinflammation response^{34 35} and glymphatic pathway,³⁶ and in relation to infarcts with AD pathology.³⁷ Recent studies have shown that low serum superoxide dismutase (SOD) is associated with a high risk of cognitive impairment after acute ischaemic stroke (AIS), suggesting that SOD is a potentially modifiable factor in PSCI.³⁸ Consistent with this, two animal model studies have shown that iVNS increases serum SOD, attenuating myocardial ischaemic injury and renal ischaemic injury.^{39 40} However, it remains to be determined whether TAVNS has the same effects, and is effective, in PSCI.

In this single-centre randomised clinical trial, we aim to assess the efficacy of TAVNS vs sham in PSCI. We hypothesise that compared with sham, TAVNS improves cognitive function, activities of daily living (ADL), quality of life and serum oxidative stress indicators.




Figure 1 Study flow chart. AE, adverse event; TAVNS, transcutaneous auricular vagus nerve stimulation.

METHODS

Design

This study will be conducted at the First Affiliated Hospital of Hunan University of Chinese Medicine as a single-centre, parallel-group, allocation concealment, assessor-blinded randomised controlled clinical trial. The Ethics Committee of the First Affiliated Hospital of Hunan University of Chinese Medicine has approved the protocol (No. HN-LL-YJSLW-2022200).

Baseline measures will include demographic information, epidemiological data, neuropsychological scores and serum oxidative stress indicators. Data relevant to clinical symptoms and cognitive domains (eg, stroke location and volume) will be collected and included in analyses as confounding factors. Eighty-eight recruited patients with PSCI will then be randomly allocated to either TAVNS or sham group at a 1:1 ratio for eight treatment weeks, according to a randomised number table generated via SPSS. Immediately postintervention (8-week follow-up) and 12 weeks later (20-week follow-up), participants will repeat the baseline neuropsychological scales and serum measurements to test whether TAVNS improves cognitive function and serum oxidative stress levels. The study design flow chart is shown in figure 1 and a timeline summary is presented in table 1. This protocol follows the

Table 1 Summary of enrolment, intervention and assessment timelines

Study period	Screening	Allocation	Intervention	Follow-up
Timepoint	-2-0 weeks	0 weeks	8 weeks	20 weeks
Enrolment				
Eligibility screen	x			
Basic information	x			
Inclusion, exclusion criteria	x			
Informed consent		x		
Randomisation		x		
Interventions				
TAVNS				
Sham TAVNS				
Assessments				
Medical history	x			
MMSE		x	x	x
MoCA		x	x	x
ADL		x	x	x
SF-36		x	x	x
Blood samples		x	x	x
Adverse events			x	x
Adherence			x	x
Treatment and rehabilitation			x	x
Daily activities diary			x	x

ADL, activities of daily living; MMSE, Mini-Mental State Examination; MoCA, Montreal Cognitive Assessment; SF-36, 36-Item Short-Form Health Survey; TAVNS, transcutaneous auricular vagus nerve stimulation.

Recommendations for Interventional Trials guidelines.⁴¹ A Standard Protocol Items: Recommendations for Interventional Trials Checklist is provided in online supplemental file.

Patient and public involvement

There will be no patient or public involvement in conducting the study.

Sample size

Sample size calculation was based on Mini-Mental State Examination (MMSE) score improvements, which will be the primary study outcome indicator. Since no studies have evaluated the effects of TAVNS on MMSE scores in PSCI, the minimal clinically relevant effect size is unknown. The sample size was thus calculated based on our preliminary study, in which we assessed the effectiveness of TAVNS (n=5) and sham (n=5) on MMSE scores in patients with PSCI. According to these data, mean pre-post treatment MMSE score differences and SDs in the TAVNS and sham groups were 7 ± 2.73 and 5.4 ± 1.14 , respectively. The estimated sample size was calculated using the formula:

$$n = 2[(z_{\alpha} + z_{\beta})^2 \sigma^2] / \delta^2$$

Alpha (α)=0.05 and power $(1-\beta)$ =0.9 determined that a minimum sample of 38 patients per group are needed.

Adjusting for a potential 15% study drop-out rate, a total of 88 participants are required (44 per group).

Inclusion criteria

The inclusion criteria are: (1) Age 45–75 years; (2) AIS diagnosis⁴² confirmed by CT or MRI; (3) Conscious, with stable vital signs; (4) First-time stroke victim, within 2–4 weeks of onset; (5) National Institutes of Health Stroke Scale (NIHSS) score 5–15; (6) Cognitive impairment, with MMSE score ≤ 26 and (7) Agree to participate and sign informed consent.

Exclusion criteria

The exclusion criteria are: (1) Concomitant disorder associated with cognitive impairment (eg, epilepsy, AD, brain trauma) or a history of medication use for cognitive impairment prior to stroke; (2) Severe language, vision or psychiatric disorder precluding a cognitive examination; (3) Severe heart, kidney, liver or endocrine disease; (4) Alcohol or drug abuse; (5) Participation in another, conflicting clinical study and (6) Currently taking medication to improve cognitive function.

Withdrawal or drop-out criteria

Participants will be terminated from the trial according to the following criteria: (1) Wanting to discontinue

the study; (2) Not cooperating with the intervention, outcomes assessments or follow-up; (3) Experiencing a serious study-related adverse event (AE).

Recruitment and screening

Recruitment advertising will be via posters on hospital bulletin boards, flyers distributed in rehabilitation and encephalopathy departments, and a recruitment station in the First Affiliated Hospital of the Hunan University of Chinese Medicine. Potential participants will be screened according to inclusion and exclusion criteria to determine enrolment eligibility. Eligible patients will be given detailed information about the trial by one of two trained research assistants. Those who decide to participate will then be asked to sign written informed consent, after which the baseline assessment will be arranged.

Baseline assessments

Relevant demographic and epidemiological data will be collected, including gender, age, education, smoking, alcohol use, illness duration, stroke location and volume, and NIHSS score. Cognitive function will then be assessed via the MMSE and Montreal Cognitive Assessment (MoCA); the Modified Barthel Index (MBI) will be used to assess daily activities and the 36-Item Short-Form Health Survey (SF-36) will be used to assess the quality of life. Blood samples will also be collected to quantify serum oxidative stress indicators, including SOD, glutathione (GSH), reactive oxygen species (ROS) and malondialdehyde (MDA).

Randomisation and allocation concealment

Following baseline assessments, eligible participants will be randomly allocated to the TAVNS or sham group at a 1:1 ratio. Random allocation sequencing will be generated using the RV.UNIFORM procedure in SPSS V.20.0 (IBM) by an independent research assistant who will not be engaged in study recruitment, assessment or intervention. Therapists will be notified of each patient's allocation and charged with arranging their allotted treatment conditions. Allocation results will be concealed from participants, outcome assessors and statistical analysts and not revealed until the completion of the final data analysis report.

Blinding

Patients, outcome assessors and statistical analysts will remain unaware of patient study allocation. To minimise discussion of the intervention, patients in different groups will not be housed in the same hospital ward. Neither outcome assessors nor statistical analysts will be engaged in intervention administration. Assessors will avoid asking for information about the stimulation location. Group allocation will be coded (A/B) and kept in sealed, opaque envelopes by a specified research assistant. Code identifiers will be unblinded after the completion of all data analyses.



Figure 2 (A) Stimulation device (SDZIIb, Huatuo). (B) Ear clip electrodes. (C) TAVNS (cymba conchae) and sham (earlobe) stimulation locations. TAVNS, transcutaneous auricular vagus nerve stimulation.

Intervention

All participants will continue to receive routine medical or rehabilitative treatment, and regular communication with their primary care physicians will be maintained throughout the study. Routine treatment and rehabilitation will be developed and implemented by the primary care physician in accordance with the Chinese Stroke Association guidelines for ischaemic cerebrovascular diseases⁴³ and Chinese Stroke Association guidelines for stroke rehabilitation.⁴⁴

Stimulation locations

The right ABVN is closely connected to the sinoatrial node, and its stimulation is more likely to cause changes in heart rate than stimulation of the left ABVN⁴⁵; thus, stimulation will be applied to the left ear in both groups. The TAVNS group stimulation target is the left auricular cymba conchae and the sham group stimulation target is the left earlobe (figure 2). These sites were chosen because the cymba conchae has the highest ABVN projection density (100%), while the earlobe has none.²⁵ Furthermore, a recent study comparing the effects of different TAVNS stimulation sites on vagal evoked potential responses showed that the cymba and concha regions are active TAVNS targets, while the earlobe is not.⁴⁶

Intervention procedure

TAVNS will be administered via an electronic acupuncture treatment instrument manufactured by Suzhou Medical Instruments Factory (SDZIIb, Huatuo, Suzhou, China, <http://www.hwato-med.com>). First, patients will take a comfortable position, seated or laying. Next, the skin on and around their left ear will be wiped with alcohol, to eliminate surplus oils and guarantee optimal conductivity. Ear electrode clips will then be placed on the left cymba conchae or earlobe, for TAVNS and sham groups, respectively. Stimulation parameters will be adjusted

for: (1) waveform: biphasic, continuous, asymmetrically balanced; (2) wave width: 200 μ s; (3) wave density: 5 Hz; (4) intensity: adjusted to the strongest sensation the individual can tolerate without pain, typically 4–6 mA; and (5) duration: 30 min/session, twice daily, 5 days/week, for 8 weeks. Other than stimulation location, the two groups' parameters will be consistent, including intervention duration and frequency.

Intervention quality control

All interventions during hospitalisation, for both groups, will be delivered by trained researchers. Patients discharged before the end of the 8-week intervention will be asked to participate in a TAVNS training course, in which a researcher will show them how to operate the TAVNS, including stimulation location and parameter settings. Participants who are approved by the trainer as familiar with the TAVNS device will self-administer the intervention at home, and complete a daily diary about the treatment.

Follow-up

After the 8-week intervention, all participants will be followed for another 12 weeks. During the follow-up period, all participants may return to their original lifestyle but will be asked to record daily activities, rehabilitation exercises and medication use. Research assistants will conduct weekly follow-up via telephone or video call to maintain participant interest and retention. Finally, all outcome indicators will be reassessed at the end of follow-up.

Treatment adherence

To enhance intervention adherence after hospital discharge, researchers will send treatment reminders twice daily via WeChat and require patients to keep a daily treatment diary, including details of when each treatment began, when it ended, side effects and improvements. Regular assessments will also be made to check all diaries and weekly telephone or video calls will be made throughout the treatment period. Participants will receive RMB 100 if they complete the project successfully, regardless of group allocation. If a patient misses two consecutive treatment weeks, or more than two total weeks, a judicious discussion and decision about whether to exclude them will occur.

Outcome assessment

All outcomes will be measured at baseline (–2 to 0 weeks), after the intervention period (8 weeks), and at the end of the study period (20 weeks). All outcome assessments will be administered by two neurologists from the First Affiliated Hospital of Hunan University of Chinese Medicine, who will be unaware of the participants' group allocation.

Primary Outcomes

The MMSE (Chinese version) will be used to measure cognitive function.⁴⁷ The MMSE is a brief 30-point questionnaire on which scores ≤ 26 indicate that cognitive

impairment may be present. Ten of the items are focused on time and place orientation, short-term memory, attention and calculation, recall, language skills, repetition, complex commands, reading, writing and pentagram reproductions.⁴⁸ However, the MMSE may be inadequate for detecting clinical signs of mild cognitive impairment and dementia^{49 50}; thus, the MoCA will also be used.

The MoCA is a short, 30-point test on which higher scores indicate better cognitive functioning. The scale assesses global cognitive abilities in attention, naming, visuospatial/executive functions, memory, language, visual structure skills, abstraction, calculation and orientation.⁵¹ When participants have received fewer than 12 years of education, one point will be added to their overall score. A score of ≤ 26 is considered intellectually deficient. The revised Chinese version (Beijing version) is widely used and has strong validity and reliability.

Secondary outcomes

The MBI, a commonly used, standardised scale, will be used to measure ADL. The MBI consists of 10 items totalling 100 points, with higher scores indicating greater daily self-care.⁵²

Quality of life will be measured by the Medical Outcomes Study SF-36, which is currently the most common clinical quality of life evaluation used in both patient and general populations. The 36-item SF-36 (Chinese version) measures eight health concepts, with excellent reliability and validity in Chinese samples⁵³; higher scores indicate better health.⁵⁴

Venous blood samples will be collected to measure oxidative stress indicators and their change rate for both groups. At each sampling, 10 mL of venous blood will be collected from the participant's left antecubital vein and sealed in vacutainer tubes with 0.5 mm EDTA. The whole blood sample will then be centrifuged at 2000 r/min for 10 min at room temperature, after which the supernatant serum will be transferred to individual, clean tubes and stored at -80°C for analysis. SOD will be measured by pyrogallol autoxidation (Superoxide Dismutase Detection Kit, Fuyuan, Fujian, China); GSH will be measured by Glutathione Assay Kit according to the manufacturer's instructions (Sigma-Aldrich R., California, USA); ROS will be measured by ROS assay kit (Beyotime Biotechnology, Shanghai, China); MDA will be measured by condensation with thiobarbituric acid at 532 nm using a Teb Pazhouhan Razi Company kit (Tehran, Iran).

Safety evaluation

AEs including dizziness, pain and unpleasant sensations at the stimulus point have been reported by a few patients in previous studies,²³ although these were mostly transitory and mild. Here, if any AE occurs during the treatment or follow-up period, it will be responded to with the appropriate intervention and management, and documented via case report form. Serious AEs will be reported immediately to the Ethics Committee to determine whether study withdrawal is necessary. Patients may also withdraw



for any safety reason or if it is in their best interest to do so.

Data management and monitoring

Data will be collected by the outcome assessors using a printed case report form, which will be independently entered into the electronic data capture system by two research assistants who are not engaged in study allocation, treatment, or assessment. The electronic data system will be accessible to the Hospital Research Centre, which will be responsible for research data monitoring and auditing. All private participant data will be coded and protected. After study completion, all documents will be securely stored for 5 years.

Statistical analysis plan

A statistician who is uninvolved with study allocation, treatment and outcome evaluation will conduct statistical analysis using SPSS V.20.0. Missing outcome data will be extrapolated using a multiple imputation method. For continuous variables, a Shapiro-Wilks distribution normality test will be performed, followed by log-transformed adjustment for non-Gaussian distributed variables. Normally distributed continuous variables will be presented as mean, SD and non-normally as median, IQR. Categorical variables will be presented as frequency or percentage.

Two-sided tests with a 5% significance level will be used for hypothesis testing. For between-groups baseline characteristic comparisons, continuous variables will be analysed using the t-test or Mann-Whitney U test and categorical variables will be analysed using the Pearson χ^2 or Fisher exact probability test. One-way analysis of variance with repeated measures (Kruskal-Wallis test) will be used to test primary and secondary outcome data. If the omnibus variance analysis is significant, then Bonferroni-adjusted post hoc analyses will be used to compare paired study periods. The Pearson χ^2 or Fisher exact probability test will be used to analyse change rates of oxidative stress indicators. To measure clinical effect magnitude, we will calculate effect size estimates using Cohen's d.

Ethics

The study has been approved by the Ethics Committee of First Affiliated Hospital of the Hunan University of Chinese Medicine and will be conducted in strict accordance with the Declaration of Helsinki. All participants will provide written informed consent, after a minimum of 3 days' consideration. All study staff will be trained in good clinical practice prior to the intervention, to achieve better study outcomes. Raw data will be anonymised and encrypted, ensuring that only the researchers actively participating in the study will have access to them.

Dissemination

The study protocol has been registered and is available on the Chinese Clinical Trial Registry website (<http://www.chictr.org.cn>; No. ChiCTR2200057808). Study

results, positive and/or negative, will be published in peer-reviewed academic journals.

DISCUSSION

This will be the first randomised controlled trial to compare the efficacy of a TAVNS vs sham intervention for improving PSCI. It will explore whether TAVNS is effective for improving cognitive function, daily functioning and quality of life among patients with PSCI. In addition, serum oxidative stress indicators SOD, GSH, ROS and MDA will be measured to help explain the neural mechanism(s) by which TAVNS improves PSCI. The study results may advance a novel, safe, effective PSCI treatment strategy with few adverse effects.

Compared with VNS, TAVNS is more convenient and economical, has fewer side effects and has been used in patients with epilepsy,⁵⁵ depression⁵⁶ and tinnitus.⁵⁷ However, the effects of TAVNS in PSCI have yet to be tested. Our stimulation parameters are based on a published study showing that TAVNS can help enhance adults' ability to learn novel letter-sound relationships.⁵⁸ In their study, Thakkar *et al* reported that TAVNS at 5 Hz and 200 μ s can improve automaticity and decoding task performance.⁵⁸ As participants vary in the stimuli intensity they can perceive and tolerate, 4–6 mA stimulus frequency will be used herein, adjusted below participants' pain tolerance levels.

Notably, TAVNS treatment duration has varied across studies,⁵⁹ including in stroke rehabilitation for 10 days,⁶⁰ 6 weeks⁶¹ and 8 weeks.⁶² Because no study to date has examined the effects of TAVNS on PSCI prevention or treatment, our design is based on assessments of verbal memory performance in patients with epilepsy. Mertens *et al*²⁹ reported improved verbal memory performance after 6 weeks of VNS. Considering that our TAVNS intervention will begin around week 4 poststroke, and that 12 weeks is widely recommended as the diagnostic time point for PSCI, it is reasonable to extend treatment to 8 weeks so that we can examine cognitive function changes.

Cognitive function tends to decline rapidly over a short period after acute stroke, and then progressively (and at least partly) recover.¹ Yet because of the unpredictability of recovery and symptom onset latency, identification is often challenging and time-consuming. In this study, we will limit inclusion to participants who are 2–4 weeks poststroke onset, to identify early cognitive impairment. The assessment period is also therefore 3–6 months poststroke, which is widely recommended as the diagnosis timeline. Thus, our timeline will best explain the TAVNS intervention effects and allow evaluation of relatively long-term efficacy.

Limitations

There are several possible study limitations. First, it is possible that some enrolled patients will be discharged before completing the 8-week intervention. Quality and adherence to out-of-hospital self-management

interventions may influence the primary or secondary outcomes. Thus, we will provide discharged patients with special TAVNS training and require that they complete a daily treatment diary. Second, differences in patient lifestyles, treatment and rehabilitation during the follow-up period may affect results. To this end, we will collect relevant data during the follow-up period and adjust statistical analyses, as needed. Finally, patient and assessor blinding may be incomplete, as differences in stimulation sites are easily recognisable.

Trial status

This trial has been registered since 17 March 2022, and was approved by the Ethics Committee of the First Affiliated Hospital of Hunan University of Chinese Medicine in April 2022. The trial began on 1 May 2022 and its completion is expected by the end of June 2023.

Contributors Z-DL conceived, designed and contributed to drafting the study protocol. H-JQ and X-QW wrote several protocol sections. C-CZ participated in study coordination and implementation. Y-JZ designed and revised the study protocol. All authors contributed to writing the protocol description and have read and approved the final version for submission.

Funding This work is supported by grants from the Hunan Provincial Health Commission Scientific Research Project (No. 202114011882) and the Hunan Provincial Traditional Chinese Medicine Scientific Research Project (No.2020004).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patient(s)

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Yue-Juan Zhang <http://orcid.org/0000-0001-7309-4173>

REFERENCES

- Huang Y-Y, Chen S-D, Leng X-Y, *et al.* Post-Stroke cognitive impairment: epidemiology, risk factors, and management. *J Alzheimers Dis* 2022;86:1–17.
- Wang K, Dong Q, YU JT. Expert consensus on the management of post-stroke cognitive impairment 2021. *Chinese Journal of Stroke* 2021;16:376–89.
- Verdelho A, Wardlaw J, Pavlovic A, *et al.* Cognitive impairment in patients with cerebrovascular disease: a white paper from the links between stroke ESO dementia Committee. *Eur Stroke J* 2021;6:5–17.
- Rimmele DL, Thomalla G. [Long-term consequences of stroke]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2022;65:498–502.
- Kwon HS, Lee D, Lee MH, *et al.* Post-Stroke cognitive impairment as an independent predictor of ischemic stroke recurrence: PICASSO sub-study. *J Neurol* 2020;267:688–93.
- Feigin VL, Brainin M, Norrving B, *et al.* World stroke organization (WSO): global stroke fact sheet 2022. *Int J Stroke* 2022;17:18–29.
- Rochmah TN, Rahmawati IT, Dahlui M, *et al.* Economic burden of stroke disease: a systematic review. *Int J Environ Res Public Health* 2021;18. doi:10.3390/ijerph18147552. [Epub ahead of print: 15 07 2021].
- McShane R, Westby MJ, Roberts E, *et al.* Memantine for dementia. *Cochrane Database Syst Rev* 2019;3:CD003154.
- Battle CE, Abdul-Rahim AH, Shenkin SD, *et al.* Cholinesterase inhibitors for vascular dementia and other vascular cognitive impairments: a network meta-analysis. *Cochrane Database Syst Rev* 2021;2:D13306.
- Quinn TJ, Richard E, Teuschl Y, *et al.* European stroke organisation and European Academy of Neurology joint guidelines on post-stroke cognitive impairment. *Eur Stroke J* 2021;6:1-XXXVIII..
- Farooq MU, Min J, Goshgarian C, *et al.* Pharmacotherapy for vascular cognitive impairment. *CNS Drugs* 2017;31:759–76.
- Ma J, Qiao P, Li Q, *et al.* Vagus nerve stimulation as a promising adjunctive treatment for ischemic stroke. *Neurochem Int* 2019;131:104539.
- Vonck K, Raedt R, Naulaerts J, *et al.* Vagus nerve stimulation...25 years later! what do we know about the effects on cognition? *Neurosci Biobehav Rev* 2014;45:63–71.
- Aniwattanapong D, List JJ, Ramakrishnan N, *et al.* Effect of vagus nerve stimulation on attention and working memory in neuropsychiatric disorders: a systematic review. *Neuromodulation* 2022;25:343–355.
- Hassert DL, Miyashita T, Williams CL. The effects of peripheral vagal nerve stimulation at a memory-modulating intensity on norepinephrine output in the basolateral amygdala. *Behav Neurosci* 2004;118:79–88.
- Roosevelt RW, Smith DC, Clough RW, *et al.* Increased extracellular concentrations of norepinephrine in cortex and hippocampus following vagus nerve stimulation in the rat. *Brain Res* 2006;1119:124–32.
- Raedt R, Clinckers R, Mollet L, *et al.* Increased hippocampal noradrenaline is a biomarker for efficacy of vagus nerve stimulation in a limbic seizure model. *J Neurochem* 2011;117:461–9.
- Follesa P, Biggio F, Gorini G, *et al.* Vagus nerve stimulation increases norepinephrine concentration and the gene expression of BDNF and bFGF in the rat brain. *Brain Res* 2007;1179:28–34.
- McGaugh JL. Consolidating memories. *Annu Rev Psychol* 2015;66:1–24.
- Harley CW. Norepinephrine and the dentate gyrus. *Prog Brain Res* 2007;163:299–318.
- Mueller D, Porter JT, Quirk GJ. Noradrenergic signaling in infralimbic cortex increases cell excitability and strengthens memory for fear extinction. *J Neurosci* 2008;28:369–75.
- Ben-Menachem E, Revesz D, Simon BJ, *et al.* Surgically implanted and non-invasive vagus nerve stimulation: a review of efficacy, safety and tolerability. *Eur J Neurol* 2015;22:1260–8.
- Redgrave J, Day D, Leung H, *et al.* Safety and tolerability of transcutaneous vagus nerve stimulation in humans; a systematic review. *Brain Stimul* 2018;11:1225–38.
- Zhang Y, Huang Y, Li H, *et al.* Transcutaneous auricular vagus nerve stimulation (taVNS) for migraine: an fMRI study. *Reg Anesth Pain Med* 2021;46:145–50.
- Peuker ET, Filler TJ. The nerve supply of the human auricle. *Clin Anat* 2002;15:35–7.
- Wang Y, Li S-Y, Wang D, *et al.* Transcutaneous auricular vagus nerve stimulation: from concept to application. *Neurosci Bull* 2021;37:853–62.
- Yakunina N, Kim SS, Nam E-C. Optimization of transcutaneous vagus nerve stimulation using functional MRI. *Neuromodulation* 2017;20:290–300.
- Kraus T, Kiess O, Hösl K, *et al.* CNS BOLD fMRI effects of sham-controlled transcutaneous electrical nerve stimulation in the left outer auditory canal - a pilot study. *Brain Stimul* 2013;6:798–804.
- Mertens A, Gadeyne S, Lescauwae E, *et al.* The potential of invasive and non-invasive vagus nerve stimulation to improve verbal memory performance in epilepsy patients. *Sci Rep* 2022;12:1984.
- Austelle CW, O'Leary GH, Thompson S, *et al.* A comprehensive review of vagus nerve stimulation for depression. *Neuromodulation* 2022;25:309–15.
- Giraudier M, Ventura-Bort C, Weymar M. Transcutaneous vagus nerve stimulation (tvNS) improves High-Confidence recognition memory but not emotional word processing. *Front Psychol* 2020;11:1276.



- 32 De Smet S, Baeken C, Seminck N, *et al.* Non-Invasive vagal nerve stimulation enhances cognitive emotion regulation. *Behav Res Ther* 2021;145:103933.
- 33 Doyle KP, Quach LN, Solé M, *et al.* B-lymphocyte-mediated delayed cognitive impairment following stroke. *J Neurosci* 2015;35:2133–45.
- 34 Chen A, Akinyemi RO, Hase Y, *et al.* Frontal white matter hyperintensities, clasmotodendrosis and gliovascular abnormalities in ageing and post-stroke dementia. *Brain* 2016;139:242–58.
- 35 Alawieh A, Tomlinson S. Injury site-specific targeting of complement inhibitors for treating stroke. *Immunol Rev* 2016;274:270–80.
- 36 Back DB, Kwon KJ, Choi D-H, *et al.* Chronic cerebral hypoperfusion induces post-stroke dementia following acute ischemic stroke in rats. *J Neuroinflammation* 2017;14:216.
- 37 Sweeney MD, Montagne A, Sagare AP, *et al.* Vascular dysfunction-The disregarded partner of Alzheimer's disease. *Alzheimers Dement* 2019;15:158–67.
- 38 Zhang M-S, Liang J-H, Yang M-J, *et al.* Low serum superoxide dismutase is associated with a high risk of cognitive impairment after mild acute ischemic stroke. *Front Aging Neurosci* 2022;14:834114.
- 39 Wang M, Deng J, Lai H, *et al.* Vagus Nerve Stimulation Ameliorates Renal Ischemia-Reperfusion Injury through Inhibiting NF- κ B Activation and iNOS Protein Expression. *Oxid Med Cell Longev* 2020;2020:7106525.
- 40 Yi C, Zhang C, Hu X, *et al.* Vagus nerve stimulation attenuates myocardial ischemia/reperfusion injury by inhibiting the expression of interleukin-17A. *Exp Ther Med* 2016;11:171–6.
- 41 Chan A-W, Tetzlaff JM, Gotsche PC, *et al.* SPIRIT 2013 explanation and elaboration: guidance for protocols of clinical trials. *BMJ* 2013;346:e7586.
- 42 Peng B, Wu B. Chinese guidelines for the diagnosis and treatment of acute ischemic stroke 2018. *Chinese Journal of Neurology* 2018;51:666–82.
- 43 Liu L, Chen W, Zhou H, *et al.* Chinese stroke association guidelines for clinical management of cerebrovascular disorders: Executive summary and 2019 update of clinical management of ischaemic cerebrovascular diseases. *Stroke Vasc Neurol* 2020;5:159–76.
- 44 Zhang T, Zhao J, Li X, *et al.* Chinese stroke association guidelines for clinical management of cerebrovascular disorders: Executive summary and 2019 update of clinical management of stroke rehabilitation. *Stroke Vasc Neurol* 2020;5:250–9.
- 45 Muppidi S, Gupta PK, Vernino S. Reversible right vagal neuropathy. *Neurology* 2011;77:1577–9.
- 46 de Gurtubay IG, Bermejo P, Lopez M, *et al.* Evaluation of different vagus nerve stimulation anatomical targets in the ear by vagus evoked potential responses. *Brain Behav* 2021;11:e2343.
- 47 Brodaty H, Connors MH, Loy C, *et al.* Screening for dementia in primary care: a comparison of the GPCOG and the MMSE. *Dement Geriatr Cogn Disord* 2016;42:323–30.
- 48 Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189–98.
- 49 Carnero-Pardo C. Should the Mini-Mental state examination be retired? *Neurologia* 2014;29:473–81.
- 50 Carnero-Pardo C. Reasons for retiring the Mini-Mental state examination. *Neurologia* 2015;30:588–9.
- 51 Nasreddine ZS, Phillips NA, Bédirian V, *et al.* The Montreal cognitive assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* 2005;53:695–9.
- 52 MacIsaac RL, Ali M, Taylor-Rowan M, *et al.* Use of a 3-Item short-form version of the Barthel index for use in stroke: systematic review and external validation. *Stroke* 2017;48:618–23.
- 53 Lin Y, Yu Y, Zeng J, *et al.* Comparing the reliability and validity of the SF-36 and SF-12 in measuring quality of life among adolescents in China: a large sample cross-sectional study. *Health Qual Life Outcomes* 2020;18:360.
- 54 Ware JE, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. conceptual framework and item selection. *Med Care* 1992;30:473–83.
- 55 von Wrede R, Rings T, Schach S, *et al.* Transcutaneous auricular vagus nerve stimulation induces stabilizing modifications in large-scale functional brain networks: towards understanding the effects of taVNS in subjects with epilepsy. *Sci Rep* 2021;11:7906.
- 56 Li S, Rong P, Wang Y, *et al.* Comparative effectiveness of transcutaneous auricular vagus nerve stimulation vs citalopram for major depressive disorder: a randomized trial. *Neuromodulation* 2022;25:450–60.
- 57 Zeng X, Zheng K, Cen J, *et al.* [Efficacy of percutaneous vagal stimulation in the treatment of chronic tinnitus]. *Lin Chung Er Bi Yan Hou Tou Jing Wai Ke Za Zhi* 2021;35:105–8.
- 58 Thakkar VJ, Engelhart AS, Khodaparast N, *et al.* Transcutaneous auricular vagus nerve stimulation enhances learning of novel letter-sound relationships in adults. *Brain Stimul* 2020;13:1813–20.
- 59 Baig SS, Kamarova M, Ali A, *et al.* Transcutaneous vagus nerve stimulation (tVNS) in stroke: the evidence, challenges and future directions. *Auton Neurosci* 2022;237:102909.
- 60 Capone F, Miccinilli S, Pellegrino G, *et al.* Transcutaneous vagus nerve stimulation combined with robotic rehabilitation improves upper limb function after stroke. *Neural Plast* 2017;2017:7876507.
- 61 Baig SS, Falidas K, Laud PJ, *et al.* Transcutaneous auricular vagus nerve stimulation with upper limb repetitive task practice may improve sensory recovery in chronic stroke. *J Stroke Cerebrovasc Dis* 2019;28:104348.
- 62 Subrahmanyam S, Suresh J. Effectiveness of transcutaneous electrical stimulation of vagus nerve among post stroke urinary incontinence. *Eur J Mol Clin Med* 2020;7:3895–913.