







# BMJ Open Experiences and well-being of healthcare professionals working in the field of ultrasound in obstetrics and gynaecology as the SARS-CoV-2 pandemic were evolving: a cross-sectional survey study

Tom Bourne <sup>1,2</sup>, Christopher Kyriacou <sup>1</sup>, Harsha Shah <sup>1</sup>, Jolien Ceusters,<sup>3</sup> Jessica Preisler,<sup>4,5</sup> Ulrike Metzger,<sup>6</sup> Chiara Landolfo <sup>7</sup>, Christoph Lees <sup>8</sup>, Dirk Timmerman <sup>2,9</sup>

**To cite:** Bourne T, Kyriacou C, Shah H, *et al.* Experiences and well-being of healthcare professionals working in the field of ultrasound in obstetrics and gynaecology as the SARS-CoV-2 pandemic were evolving: a cross-sectional survey study. *BMJ Open* 2022;**12**:e051700. doi:10.1136/bmjopen-2021-051700

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-051700>).

Received 12 April 2021

Accepted 05 January 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Professor Tom Bourne;  
t.bourne@imperial.ac.uk

## ABSTRACT

**Objective** Assess experience of healthcare professionals (HCPs) working with ultrasound in obstetrics and gynaecology during the evolving SARS-CoV-2 pandemic, given the new and unprecedented challenges involving viral exposure, personal protective equipment (PPE) and well-being.

**Design** Prospective cross-sectional survey study.

**Setting** Online international survey. Single-best, open box and Hospital Anxiety and Depression Scale (HADS) questions.

**Participants** The survey was sent to 35 509 HCPs in 124 countries and was open from 7 to 21 May 2020. 2237/3237 (69.1%) HCPs from 115 countries who consented to participate completed the survey. 1058 (47.3%) completed the HADS.

**Primary outcome measures** Overall prevalence of SARS-CoV-2, depression and anxiety among HCPs in relation to country and PPE availability.

**Analyses** Univariate analyses were used to investigate associations without generating erroneous causal conclusions.

**Results** Confirmed/suspected SARS-CoV-2 prevalence was 13.0%. PPE provision concerns were raised by 74.1% of participants; highest among trainees/resident physicians (83.9%) and among HCPs in Spain (89.7%). Most participants worked in self-perceived high-risk areas with SARS-CoV-2 (67.5%–87.0%), with proportionately more trainees interacting with suspected/confirmed infected patients (57.1% vs 24.2%–40.6%) and sonographers seeing more patients who did not wear a mask (33.3% vs 13.9%–7.9%). The most frequent PPE combination used was gloves and a surgical mask (22.3%). UK and US respondents reported spending less time self-isolating (8.8 days) and lower satisfaction with their national pandemic response (37.0%–43.0%). 19.8% and 8.8% of respondents met the criteria for moderate to severe anxiety and depression, respectively.

**Conclusions** Reported prevalence of SARS-CoV-2 in HCPs is consistent with literature findings. Most respondents

## Strengths and limitations of this study

- Strengths of our study include the sample size, the assessment of healthcare professional (HCP) responses internationally and the ability to compare responses between different HCPs.
- Four languages enabled greater participant inclusion, and responses between International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) and non-ISUOG members were compared prior to grouping for analysis.
- Key weaknesses include possible selection bias due to the method of sample selection and the low population response rate.
- It is important not to draw conclusions about causation as this is a descriptive study performed as the pandemic was evolving.
- At the time of survey participation, the majority of HCPs did not have the same access to a swab or antibody test that is now available.

used gloves and a surgical mask, with a greater SARS-CoV-2 prevalence compared with those using 'full' PPE. HCPs with the least agency (trainees and sonographers) were not only more likely to see high-risk patients but also less likely to be protected. A fifth of respondents reported moderate to severe anxiety.

## INTRODUCTION

### Background

The SARS-CoV-2 pandemic has caused unprecedented challenges to all healthcare systems.<sup>1</sup> These include patient protection, managing the burden of disease on service capacity and mitigating adverse effects on healthcare professionals (HCPs: physicians, sonographers, nurses and allied HCPs).<sup>2</sup>

Guidance has internationally focused on clinician redeployment to front-line settings, prioritisation of services, provision and rationalisation of personal protective equipment (PPE).<sup>3,4</sup> During the pandemic, access to PPE for HCPs has remained a key concern. Despite HCPs being prioritised in most countries, PPE shortages have been described universally<sup>1</sup> and have been a particular concern in the early months of the pandemic. Lack of sufficient and adequate PPE is important in obstetrics and gynaecology, where HCPs work in a variety of settings, including ultrasound scanning, may not ordinarily be considered high-risk but involves close patient proximity for extended periods.<sup>5</sup> These areas are easily overlooked but may pose a high risk of SARS-CoV-2 infection.<sup>6</sup>

A further concern has been the availability of SARS-CoV-2 testing among HCPs. Initially, provision for this was variable, with, for example, relatively easy access in Hong Kong compared with virtually none in the UK. This is important as, in the absence of testing, HCP risk becoming COVID-19 vectors within hospitals and the wider community. Lack of testing may also lead to unnecessary isolation from work. There are limited data regarding HCP testing availability and time HCPs have taken off work due to suspected or confirmed SARS-CoV-2.<sup>7</sup>

Less attention has been focused on the implications of infectious diseases on HCP psychological health, which previous pandemics have shown to be significant.<sup>8,9</sup> HCPs are confronted with ongoing resource, shift pattern, PPE and testing uncertainty,<sup>1,10</sup> as well as COVID-19 exposure risk and the implications of this on their families.<sup>11</sup> Protecting the psychological health of the medical workforce is critical, particularly as anxiety, depression and burnout are recognised complications for HCPs working in high-stress environments.<sup>12</sup>

## Objectives

We aimed to assess the scale, experience and psychological well-being of clinicians who have experienced the evolving SARS-CoV-2 pandemic working with ultrasound in obstetrics and gynaecology internationally between 7 and 21 May 2020.

## METHODS

### Study design

We used a cross-sectional survey design with two main sections assessing physical aspects, including underlying medical conditions, comorbidities and personal SARS-CoV-2 experience, and psychological well-being (online supplement 1: The International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) SARS-CoV-2). Participants were initially asked to provide informed consent prior to continuing to the online survey. If they did not give consent by answering 'yes' to the first question, they were not directed to the survey. Single-best answer and open box questions were proposed by a senior gynaecologist. These were then piloted, vetted and modified by the authors before being translated from English to French,

Spanish and Italian in order to increase survey accessibility. We estimated that the time taken to complete the questionnaire was 10 min.

### Setting

The online platform SurveyMonkey (2020) was selected to upload and disseminate the questionnaire, which was encrypted, and multiple responses were disabled. The survey design ensured the data collected was completely anonymous for all respondents.

### Participants

All HCPs on the International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) mailing list were invited to participate in this study between 7 and 21 May 2020. There were 1237953 confirmed cases of SARS-CoV-2 infection and 63313 deaths in the time the survey was open.<sup>13</sup> Various countries had instigated local and national lockdowns at this time, including the UK. Members were sent an email containing information describing the study and individual links to the questionnaire in English, Spanish, Italian and French. We made it clear to the participants that their participation was voluntary and that responses would be both anonymous and untraceable.

During the survey period, two reminders were sent out. As this was a survey study of a relatively new condition at the time, a power calculation was not performed. Exclusion criteria included retired doctors as this was deemed to not be reflective of the working environment during the SARS-CoV-2 pandemic. A copy of the questionnaire in English can be viewed in online supplement 1: The International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) SARS-CoV-2.

### Patient and public involvement statement

Given the HCP participant population and the acute nature of the SARS-CoV-2 pandemic, patient and public involvement was not performed.

### Variables and data sources

#### Physical aspects

This section was divided into subsections: Introduction; Demographics; Underlying medical conditions; Medications; Smoking and BMI; Role, shift pattern, PPE; SARS-CoV-2 diagnosis; Personal SARS-CoV-2 pathway; Household; Colleagues; and Support (online supplement 1: The International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) SARS-CoV-2: questions 2–40).

Participants were asked details about their job and organisational attributes, as well as on a variety of personal comorbidities relevant to SARS-CoV-2 infection.<sup>14</sup> They were then questioned on their personal experience of SARS-CoV-2 at home and in the workplace.

#### Psychological well-being - Hospital Anxiety and Depression Scale (HADS)

Participants were given the option to complete the validated HADS, which is a 14-item (seven questions related

to anxiety and depression each) questionnaire<sup>15</sup> (online supplement 1: The International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) SARS-CoV-2: questions 41–56). Each subscale measures symptom severity (the score ranges between 0 and 21, a score greater or equal to 11 indicates moderate-to-severe symptoms). If respondents chose not to complete the HADS, the survey ended.

### Outcome measures

The primary outcome was the overall prevalence of SARS-CoV-2, depression and anxiety among HCPs in relation to country and PPE availability.

### Statistical analyses

Univariate analyses were used to investigate the association between country of origin, type of HCP and PPE without generating erroneous causal conclusions. Results were presented as percentages (%), mean, 95% CIs, medians and ranges, depending on the variable. Missing data, that is, questions that were skipped by HCPs as they completed their questionnaire, were not included in the analyses.

HCPs and areas of work were divided into groups to allow reasonable comparable analyses to be performed. For HCPs, there were three main categories: consultants/attending physicians (gynaecologists, obstetricians, obstetricians and gynaecologists or radiologists/sonologists aged 35 years and older); trainees/residents (gynaecologists, obstetricians, obstetricians and gynaecologists or radiologists/sonologists aged 34 years and younger, doctors in research and doctors in training); and sonographers (sonographers and allied HCPs (midwives and nurses)) (online supplement 3: Breakdown of healthcare professional (HCP) roles for analysis).

Areas of work were defined as high-risk (gynaecology (early pregnancy unit, acute gynaecology and ultrasound), obstetrics (birth centre/birthing centre, labour ward, maternity triage/hospital obstetric triage and ultrasound) and radiology (sonology–ultrasound)); moderate risk (obstetrics (antenatal clinic, antenatal ward, community clinics and postnatal ward) and radiology (non-ultrasound)); or low-risk (working from home, gynaecology (benign and oncology) and obstetrics (maternity helplines)) (online supplement 4: Breakdown of risk by area of work for analysis). This categorisation approximated risk based on HCP proximity and time spent with patients, allowing meaningful data analysis.

Anxiety and depression scores were divided by case-ness in order to differentiate the proportion of the study population that had no anxiety or depression (HADS score 0–7), those experiencing mild anxiety or depression (HADS score 8–10), and those suffering from moderate-to-severe anxiety or depression (HADS 11–21).

The statistical analyses were performed using R V.3.5.1.

## RESULTS

### Summary

#### Response

The survey was sent to 35 509 HCPs in 124 countries. A total of 3287 (9.3%) HCPs clicked the survey link and 3237 (98.5%) of them consented to complete it. A total of 1960 (60.5%) responded in English, 881 (27.2%) in Spanish, 267 (8.2%) in Italian and 129 (4.0%) in French. The response rate for ISUOG members was 23.1% (2441/10589), while that for non-ISUOG members was 3.2% (796/24920) (online supplement 5: Flowchart summarizing survey participants, differentiating International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) members and non-ISUOG members (n = 3287)). Of 3237 HCPs from 115 countries who started the survey, 2237 (69.1%) completed it in full. Mean age of respondents was 47.2 years, with a range from 18 to 82 years. The majority (83%) were aged 31–60 years old. A total of 1474 (66%) of respondents were female and 755 (34%) were male, with 4 intersex or undisclosed. Of 2237 participants, 1058 (47.3%) completed the HADS, and the remainder (1179/2237, 52.7%) did not. Survey participation and demographics are summarised in table 1.

Direct comparison of data between the characteristics of ISUOG members and non-ISUOG members confirmed findings were similar (online supplement 6: Demographics of survey participation, differentiated based on International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) membership (n = 3237)). It was therefore deemed appropriate to combine these populations for the analysis.

### COVID-19 prevalence

Confirmed or suspected SARS-CoV-2 prevalence was 13.0% (290/2237). Among this group, 11.4% had a history of heart/lung disease; 13.4% were taking antihypertensives; and 4.5% were smokers (online supplement 7: Comparison of comorbidity in those with (n = 290) and without suspected or confirmed SARS-CoV-2 (n = 290) (Total n = 2237)).

### PPE use and provision

Of all HCPs, 12.5% reported not using PPE, while 87.5% reported variable use. Of the total HCPs, 74.1% had concerns about provision or shortages of one or more items of PPE, in particular N95/FFP3 masks (57.2%). PPE was reported to be recycled or reused by 61.3%.

### Anxiety and depression

Of the total participants, 21.6% met the criteria for mild anxiety, and 19.8% met the criteria for moderate-to-severe anxiety; 19.4% met the criteria for mild depression and 8.8% for moderate-to-severe depression. Of those with suspected/confirmed SARS-CoV-2, 27.2% and 12.3% reported moderate-to-severe anxiety and depression, compared with 18.4% and 8.1% of those without (online supplement 8: Comparison of Hospital Anxiety and Depression scale (HADS) scores in those with (n =



**Table 1** Demographics of total survey participation (N=3237)

	Total (N=3237)
Total ISUOG mailing list (n)	35509
Response of total ISUOG mailing list, N (%)	3237 (9)
Countries (n)	124
Respondents in English (n)	1960
Respondents in Spanish (n)	881
Respondents in Italian (n)	267
Respondents in French (n)	129
ISUOG members, n (%)	2441 (23)
Completed survey, n (%)	2237 (69)
ISUOG members, n (%)	1795 (80)
Countries (n)	115
Age (years), mean (range)	47.2 (18–82)
≤20, n (%)	1 (0)
21–30, n (%)	117 (5)
31–40, n (%)	558 (25)
41–50, n (%)	686 (31)
51–60, n (%)	596 (27)
61–70, n (%)	251 (11)
>70, n (%)	24 (1)
Gender, n (%)	
Female	1474 (66)
Male	755 (34)
Intersex	1 (0)
Prefer not to say	3 (0)
Prevalence suspected/confirmed SARS-CoV-2 based on symptoms±PCR testing, n (%)	290 (13)
Is PPE used by the respondents? n (%)	
Not yet	279 (12)
Yes—for aerosol-generating procedures only	133 (6)
Yes—for all patients	1392 (62)
Yes—for suspected/positive patients only	429 (19)
What PPE is used by the respondents? n (%)	
None	76 (3)
Gloves only	49 (2)
Gloves+surgical mask	497 (22)
Gloves+surgical mask+visor	152 (7)
Gloves+FFP3 mask	95 (4)
Gloves+FFP3 mask+visor	51 (2)
Gloves+FFP3 mask+visor+gown+surgical hat	90 (4)
Patient wearing mask	1841 (82)
Concerns regarding a lack of PPE, n (%)	
No	579 (26)
Gloves	278 (12)
Gown	629 (28)
N95/FFP3 mask	1279 (57)
Surgical hat	233 (10)
Surgical mask	820 (37)
Visor	582 (26)

Continued

**Table 1** Continued

	Total (N=3237)
Surgical mask (patient)	609 (27)
Type of PPE that has been reused or recycled, n (%)	
No	865 (39)
Gloves	34 (2)
Gown	287 (13)
N95/FFP3 mask	886 (40)
Surgical hat	92 (4)
Surgical mask	565 (25)
Visor	621 (28)
Anxiety, mean (95% CI)*	6.9 (6.7 to 7.2)
0–7 (none), n (%)	620 (59)
8–10 (mild), n (%)	229 (22)
11–21 (moderate to severe), n (%)	209 (20)
Depression, mean (95% CI)*	5.3 (5.1 to 5.5)
0–7 (none), n (%)	760 (72)
8–10 (mild), n (%)	205 (19)
11–21 (moderate to severe), n (%)	93 (9)

\*Numbers of Hospital and Anxiety Depression Scale (HADS) respondents lower than the main survey (n=1058).  
ISUOG, International Society of Ultrasound in Obstetrics and Gynecology; PCR, Polymerase Chain Reaction; PPE, personal protective equipment.

162) and without suspected or confirmed SARS-CoV-2 (n = 896) (Total n = 1058)). A greater proportion of female participants scored for moderate-to-severe anxiety (156/754, 20.7%) and depression (79/754, 10.5%) compared with male participants (41/303, 13.5% and 14/303, 4.6%) (online supplement 9: Hospital Anxiety and Depression scale (HADS) breakdown by gender (n = 1058)).

Four out of 2237 participants were excluded from the univariate analyses, having completed the survey following retirement.

### Univariate analysis by country (N=2233)

A breakdown of findings by countries that had most respondents is presented in [table 2](#).

### COVID-19 prevalence

The highest prevalence of SARS-CoV-2 infection based on symptoms and/or a positive PCR swab was reported in the UK (31.1%), followed by Italy (14.9%), Spain (14.7%) and the USA (13.7%). Respondents from India and the Philippines reported a prevalence of 2.6% and 8.8%, respectively. Only 0.9% and 1.0% of Indian and UK participants, respectively, had an antibody test, compared with 52.9% and 60.3% of participants in Italy and Spain, respectively. Of participants from the USA and Spain who had a SARS-CoV-2 antibody test, 25.0% and 13.4%, respectively, had a positive result.

Participants in the UK reported spending an average of 8.8 days self-isolating away from the workplace, the same as those in the USA. Mean self-isolation time was only

**Table 2** Breakdown of findings by country (N=2233)

	Italy	Spain	India	Philippines	UK	USA	Other*
Respondents, n (%)	174 (8)	136 (6)	115 (5)	114 (5%)	103 (5)	102 (5)	1489 (67)
Prevalence of suspected/confirmed SARS-CoV-2 based on symptoms±PCR testing, n (%)	26 (15)	20 (15)	3 (3)	10 (9)	32 (31)	14 (14)	185 (12)
Had SARS-CoV-2 antibody test, n (%)	92 (53)	82 (60)	1 (1)	21 (18)	1 (1)	16 (16)	155 (10)
Had positive SARS-CoV-2 antibody, n (%)	6 (7)	11 (13)	0 (0)	0 (0)	0 (0)	4 (25)	16 (10)
Is PPE used by the respondents? n (%)							
Not yet	17 (10)	28 (21)	31 (27)	6 (5)	7 (7)	5 (5)	185 (12)
Yes—for aerosol-generating procedures only	8 (5)	6 (4)	9 (8)	7 (6)	6 (6)	3 (3)	94 (6)
Yes—for all patients	107 (61)	33 (24)	65 (57)	96 (84)	76 (74)	83 (81)	932 (63)
Yes—for suspected/positive patients only	42 (24)	69 (51)	10 (9)	5 (4)	14 (14)	11 (11)	278 (19)
What PPE is used by the respondents? n (%)							
None	0 (0)	1 (1)	1 (1)	1 (1)	1 (1)	0 (0)	72 (5)
Gloves only	0 (0)	0 (0)	0 (0)	0 (0)	2 (2)	0 (0)	47 (3)
Gloves+surgical mask	79 (45)	42 (31)	7 (6)	0 (0)	41 (40)	45 (44)	283 (19)
Gloves+surgical mask+visor	6 (3)	11 (8)	6 (5)	1 (1)	17 (17)	12 (12)	99 (7)
Gloves+FFP3 mask	12 (7)	10 (7)	8 (7)	0 (0)	0 (0)	6 (6)	59 (4)
Gloves+FFP3 mask+visor	1 (1)	4 (3)	6 (5)	2 (2)	0 (0)	3 (3)	35 (2)
Gloves+FFP3 mask+visor+gown+surgical hat	3 (2)	5 (4)	13 (11)	23 (20)	0 (0)	1 (1)	45 (3)
Patient wearing mask	173 (99)	133 (98)	109 (95)	111 (97)	33 (32)	91 (89)	1191 (80)
Concerns regarding a lack of PPE, n (%)							
No	35 (20)	14 (10)	52 (45)	20 (18)	46 (45)	24 (24)	388 (26)
Gloves	25 (14)	19 (14)	5 (4)	12 (11)	1 (1)	9 (9)	207 (14)
Gown	53 (30)	45 (33)	23 (20)	54 (47)	25 (24)	23 (23)	406 (27)
N95/FFP3 mask	118 (68)	110 (81)	51 (44)	84 (74)	33 (32)	61 (60)	822 (55)
Surgical hat	15 (9)	10 (7)	10 (9)	15 (13)	4 (4)	8 (8)	171 (11)
Surgical mask	84 (48)	62 (46)	10 (9)	21 (18)	24 (23)	45 (44)	574 (39)
Visor	51 (29)	51 (38)	11 (10)	19 (17)	19 (18)	24 (24)	407 (27)
Surgical mask (patient)	44 (25)	47 (35)	10 (9)	32 (28)	13 (13)	31 (30)	432 (29)
Type of PPE that has been reused or recycled, n (%)							
No	55 (32)	31 (23)	40 (35)	25 (22)	62 (60)	12 (12)	640 (43)
Gloves	2 (1)	3 (2)	5 (4)	2 (2)	0 (0)	1 (1)	21 (1)
Gown	15 (9)	17 (13)	23 (20)	70 (61)	3 (3)	4 (4)	155 (10)
N95/FFP3 mask	82 (47)	82 (60)	59 (51)	66 (58)	10 (10)	68 (67)	519 (35)
Surgical hat	2 (1)	7 (5)	1 (1)	16 (14)	1 (1)	4 (4)	61 (4)
Surgical mask	83 (48)	60 (44)	7 (6)	13 (11)	14 (14)	65 (64)	323 (22)
Visor	23 (13)	48 (35)	31 (27)	56 (49)	28 (27)	28 (27)	407 (27)
Mean days of self-isolation (range)	11.3 (0–30)	11.6 (0–30)	17.3 (0–30)	16.9 (0–30)	8.8 (0–30)	8.8 (0–30)	8.4 (0–30)
Mean days of household self-isolating (range)	21.1 (14–30)	10.5 (2–15)	18 (10–30)	15.3 (7–30)	11.1 (2–14)	15 (2–30)	12.2 (1–30)
Respondents satisfied with local unit SARS-CoV-2 response, n (%)	89 (51)	97 (71)	89 (77)	89 (78)	75 (73)	74 (73)	1018 (68)
Mean % level of respondent satisfaction with government SARS-CoV-2 response (95% CI)	38 (34 to 41)	23 (19 to 26)	61 (56 to 65)	48 (44 to 52)	43 (38 to 48)	37 (31 to 43)	53 (52 to 55)
Anxiety, mean (95% CI)†	6.3 (5.6 to 7.1)	6.9 (6.1 to 7.7)	7.4 (6.1 to 8.8)	7.1 (5.6 to 8.6)	7.2 (6.2 to 8.2)	7.3 (6.1 to 8.4)	6.9 (6.6 to 7.2)
0–7 (none), n (%)	74 (71)	51 (61)	22 (51)	18 (50)	36 (58)	35 (59)	384 (57)
8–10 (mild), n (%)	13 (12)	21 (25)	13 (30)	10 (28)	12 (19)	8 (14)	152 (23)
11–21 (moderate to severe), n (%)	17 (16)	12 (14)	8 (19)	8 (22)	14 (23)	16 (27)	134 (20)
Depression, mean (95% CI)†	5.8 (5.2 to 6.5)	5 (4.2 to 5.7)	5.1 (4.1 to 6.1)	5.3 (4.2 to 6.5)	5.3 (4.3 to 6.2)	4.8 (3.8 to 5.7)	5.3 (5.0 to 5.6)

Continued

Table 2 Continued

	Italy	Spain	India	Philippines	UK	USA	Other*
0–7 (none), n (%)	74 (71)	67 (80)	31 (72)	27 (75)	44 (71)	46 (78)	471 (70)
8–10 (mild), n (%)	20 (19)	11 (13)	10 (23)	8 (22)	12 (19)	9 (15)	135 (20)
11–21 (moderate to severe), n (%)	10 (10)	6 (7)	2 (5)	1 (3)	6 (10)	4 (7)	64 (10)

\*The 1489 within the 'other' cohort make up data from participants of the remaining 109 countries. We provide a breakdown of the six countries that had the most respondents.

†Numbers of Hospital and Anxiety Depression Scale (HADS) respondents lower than the main survey (n, Italy=104; Spain=84; India=43; Philippines=36; UK=62; USA=59; Other=670).

PCR, Polymerase Chain Reaction; PPE, personal protective equipment.

greater than 2 weeks in India and in the Philippines (17.3 and 16.9 days, respectively), where SARS-CoV-2 prevalence among HCPs in the study was lowest at the time of the study.

### PPE use and provision

The use of PPE when performing ultrasound was reported by 73.0%–95.1% of participants. The most common combination of PPE used while performing ultrasonography was gloves and a surgical mask, with the highest rates of use in Italy (45.4%), the UK (39.8%) and the USA (44.1%). No participants from the Philippines and only 6.1% from India reported using only gloves and a surgical mask. Visor use, in addition to gloves and a surgical mask, was highest in the UK (16.5%) and the USA (11.8%). Full PPE (gloves, FFP3/N95 mask, visor, gown and surgical hat) was used by 4% (90/2233) of the respondents. Full PPE use was highest in the Philippines (20.2%) and lowest in the UK (0%). Apart from the UK (32.0%), high rates of patients wearing a mask were noted in every other country analysed (80.0%–99.4%).

PPE provision or supply concerns were highest in Spain (89.7%), Italy (79.9%) and the Philippines (82.5%), and lowest in the UK (55.3%) and India (54.8%). Less PPE was reported as being recycled or reused in the UK (39.8%) compared with any other country analysed. In the USA, 88.2% of HCPs reported PPE recycling or reuse.

### Anxiety, depression and support

The HADS revealed that in the UK, 22.6%, and in the USA, 27.1% of participants met the criteria for moderate to severe anxiety, higher than any other country analysed, with Spanish participants having the lowest rate (14.3%). Italian, UK and 'other countries' participants had the highest rates of moderate to severe depression (9.6%–9.7%), with Philippines HCPs having the lowest rates (2.8%).

Levels of satisfaction with the support offered by local units during the pandemic ranged from 68% to 78% in all countries except Italy, where the level was markedly lower (51.1%). Indians had the highest satisfaction rate with their government response to the pandemic with a mean of 61%, compared with 43% in the UK, 37% in the USA, 38% in Italy and 23% in Spain.

### Univariate analysis by HCP (N=2233)

Table 3 focuses on HCPs—66.3% consultants/attending physicians, 22.3% sonographers and 11.4% trainee/resident physicians (who are the youngest cohort overall). Of physicians and sonographers, 62.4%–63.3% and 54.3%–55.6%, respectively, reported changes to their duties and patient contact throughout the pandemic.

### COVID-19 prevalence

Suspected or confirmed SARS-CoV-2 prevalence was similar across the groups (12.6% to 14.2%). 87.0% of trainees worked in high-risk areas, compared with 77.3% of consultants and 67.5% of sonographers. Trainees self-isolated for 1 day less on average (8.5 days) compared with consultants and sonographers (9.6 and 9.7 days, respectively).

### PPE use and provision

The PPE reported in each HCP group as most frequently used when performing ultrasonography was gloves and a surgical mask (21.0%–30.3%), with 11.6%–15.0% reporting no PPE use. Sonographers saw more patients who did not wear a mask (33.3%) compared with trainees (7.9%) and consultants (13.9%).

A larger proportion of trainees (57.1%) interacted with patients with suspected or confirmed SARS-CoV-2 compared with consultants (40.6%) and sonographers (24.2%) but were less likely to use PPE for all patients (48.8%), compared with consultants (63.1%) and sonographers (66.9%).

Of the total trainees, 83.9% reported PPE concerns (compared with 73.1% of consultants and 71.9% of sonographers), with 74.0% reusing or recycling PPE (compared with 59.9% consultants and 58.9% sonographers).

### Anxiety, depression and support

Rates of anxiety (39.4%–47.7%) and depression (26.9%–31.0%) were similar across the HCP groups, with 18.3%–25.2% and 7.5%–9.2% of participants meeting the criteria for moderate to severe anxiety and depression, respectively. Those working in high-risk areas appear more inclined to complete the HADS (online supplement 10: Hospital Anxiety and Depression scale (HADS) breakdown by risk of working area (n = 1058)). However, rates of mild to severe anxiety are overall similar, with more depression reported by those working in lower-risk areas.

**Table 3** Breakdown of findings by HCP (N=2233)

	Consultants/attending	Sonographer	Trainees/residents
Respondents, n (%)	1480 (66)	499 (22)	254 (11)
Changes in shift pattern as a result of the pandemic, n (%)			
Unable to work	34 (2)	21 (4)	11 (4)
Change in duties due to SARS-CoV-2 pandemic, n (%)			
Increased work	185 (13%)	64 (13%)	37 (15%)
Similar work	570 (39%)	228 (46%)	82 (32%)
Decreased work	691 (47%)	186 (37%)	124 (49%)
Change in patient contact due to the SARS-CoV-2 pandemic, n (%)			
Increased patient contact	114 (8)	35 (7)	22 (9)
Similar patient contact	530 (36)	212 (42)	90 (35)
Decreased patient contact	743 (50)	214 (43)	119 (47)
No patient contact	59 (4)	17 (3)	12 (5)
Level of COVID-19 risk of working environment reported by HCPs, n (%)			
High risk of COVID-19 infection	1144 (77%)	337 (68%)	221 (87%)
Moderate risk of COVID-19 infection	128 (9%)	32 (6%)	14 (6%)
Low risk of COVID-19 infection	208 (14%)	130 (26%)	19 (7%)
Interaction with patients with suspected/confirmed SARS-CoV-2, n (%)	601 (41%)	121 (24%)	145 (57%)
Is PPE used by the respondents? n (%)			
Not yet	171 (12)	75 (15)	33 (13)
Yes—for aerosol-generating procedures only	89 (6)	28 (6)	16 (6)
Yes—for all patients	934 (63)	334 (67)	124 (49)
Yes—for suspected/positive patients only	286 (19)	62 (12)	81 (32)
What PPE is used by the respondents? n (%)			
None	47 (3)	21 (4)	8 (3)
Gloves only	20 (1)	25 (5)	4 (2)
Gloves+surgical mask	315 (21)	105 (21)	77 (30)
Gloves+surgical mask+visor	111 (8)	35 (7)	6 (2)
Gloves+FFP3 mask	70 (5)	13 (3)	12 (5)
Gloves+FFP3 mask+visor	41 (3)	8 (2)	2 (1)
Gloves+FFP3 mask+visor+gown+surgical hat	59 (4)	24 (5)	7 (3)
Patient wearing mask	1274 (86)	333 (67)	234 (92)
Concerns regarding a lack of PPE, n (%)			
No	398 (27)	140 (28)	41 (16)
Gloves	176 (12)	63 (13)	39 (15)
Gown	403 (27)	133 (27)	93 (37)
N95/FFP3 mask	860 (58)	239 (48)	180 (71)
Surgical hat	142 (10)	61 (12)	30 (12)
Surgical mask	516 (35)	189 (38)	115 (45)
Visor	379 (26)	122 (24)	81 (32)
Surgical mask (patient)	395 (27)	133 (27)	81 (32)
Type of PPE that has been reused or recycled, n (%)			
No	594 (40)	205 (41)	66 (26)
Gloves	22 (1)	9 (2)	3 (1)
Gown	166 (11)	92 (18)	29 (11)
N95/FFP3 mask	596 (40)	152 (30)	138 (54)
Surgical hat	59 (4)	22 (4)	11 (4)

Continued

Table 3 Continued

	Consultants/attending	Sonographer	Trainees/residents
Surgical mask	357 (24)	116 (23)	92 (36)
Visor	401 (27)	138 (28)	82 (32)
Prevalence of suspected/confirmed SARS-CoV-2-based on symptoms±PCR testing, n (%)	187 (13)	71 (14)	32 (13)
Had SARS-CoV-2 antibody test, n (%)	259 (18)	55 (11)	54 (21)
Had positive SARS-CoV-2 antibody, n (%)	28 (11)	5 (9)	4 (7)
Mean days of self-isolation (range)	9.6 (0–30)	9.7 (0–30)	8.5 (0–30)
Mean days of household self-isolating (range)	13 (1–30)	14.3 (2–30)	8.8 (2–14)
Respondents satisfied with local unit SARS-CoV-2 response, n (%)	1043 (70)	340 (68)	148 (58)
Mean % level of respondent satisfaction with government SARS-CoV-2 response (95% CI)	47 (46 to 49)	57 (54 to 59)	44 (40 to 47)
Anxiety, mean (95% CI)*	6.7 (6.4 to 7.0)	7.7 (7.1 to 8.3)	6.9 (6.3 to 7.6)
0–7 (none), n (%)	433 (61)	112 (52)	75 (58)
8–10 (mild), n (%)	151 (21)	48 (22)	30 (23)
11–21 (moderate to severe), n (%)	131 (18)	54 (25)	24 (19)
Depression, mean (95% CI)*	5.3 (5.0 to 5.5)	5.5 (5.0 to 6.1)	5.3 (4.7 to 5.9)
0–7 (none), n (%)	523 (73)	148 (69)	89 (69)
8–10 (mild), n (%)	126 (18)	50 (23)	29 (22)
11–21 (moderate to severe), n (%)	66 (9)	16 (7)	11 (9)

\*Numbers of Hospital and Anxiety Depression Scale (HADS) respondents lower than the main survey (n, Consultants/attending=715; Sonographer=214; Trainees/residents=129).

HCP, healthcare professional; PCR, Polymerase Chain Reaction; PPE, personal protective equipment.

Trainees reported lower satisfaction/support from their local unit (58.3%, compared with 70.5% and 68.1% in consultants and sonographers, respectively) and government (44%, compared with 47% and 57% in consultants and sonographers, respectively).

### Univariate analysis by PPE (N=2233)

Table 4 focuses on common combinations of PPE used when performing an ultrasound scan.

### COVID-19 prevalence

Of the total respondents, 2.2% reported using gloves in isolation. Twenty-nine per cent of this group reported suspected or confirmed COVID-19.

The most used combination of PPE when performing an ultrasound scan was gloves and a surgical mask (used by 22.3% of respondents). Of participants using this combination of PPE, 45.4% (226/497) reported interacting with patients with confirmed SARS-CoV-2, with 16.9% of this group reporting suspected or confirmed COVID-19.

The use of gloves with an FFP3/N95 mask instead of a surgical mask when performing ultrasound was reported by 4.3% (without visor) and 2.3% (with visor) of HCPs. Of those using gloves and FFP3/N95 mask, 10.5% reported suspected or confirmed COVID-19, compared with 3.9% of those who also used a visor.

Four per cent of all HCPs reported using 'full' PPE, 81.1% of whom worked in a COVID-19 high-risk setting. Full PPE advised for use by clinicians when interacting with a suspected or confirmed SARS-CoV-2 patient

includes gloves, gown, N95/FFP3 mask, a surgical hat and visor. Of those using full PPE, 37.8% interacted at least once with patients with suspected or confirmed SARS-CoV-2. Of this group, 6.7% reported suspected or confirmed SARS-CoV-2.

Of the HCPs, 82.4% reported patients wearing a mask during clinical interactions. This was within a high-risk area of work in 76.5% cases.

### Anxiety, depression and support

HCPs who reported PPE shortages had higher rates of anxiety (45.4%) and depression (31.5%) compared with those without shortages (29.6% and 18.5%, respectively) (online supplement 11: Comparison of Hospital Anxiety and Depression scale (HADS) scores in those with (n = 788) and without personal protective equipment (PPE) shortages (n = 270) (Total n = 1058)).

Lower satisfaction/support from their local unit ranged between 66% and 75% with a mean level of government satisfaction ranging between 40% and 68% between the PPE subgroups.

## DISCUSSION

### Summary

We found that trainees and sonographers are generally more exposed to SARS-CoV-2, with sonographers seeing more patients who did not wear a mask and trainees working the most in higher risk areas. Trainees are thus



**Table 4** Breakdown of findings by combinations of PPE (N=2233)\*

	None	Gloves only	Gloves+surgical mask	Gloves +surgical mask+visor	Gloves +FFP3 mask	Gloves +FFP3 mask+visor	Gloves +FFP3 mask +visor+gown +surgical hat
Respondents (n, %)	76 (3)	49 (2)	497 (22)	152 (7)	95 (4)	51 (2)	90 (4)
Level of COVID-19 risk of working environment reported, n (%)							
High risk of COVID-19 infection	50 (66)	34 (69)	394 (79)	104 (68)	60 (63)	37 (73)	73 (81)
Moderate risk of COVID-19 infection	9 (12)	3 (6)	38 (8)	16 (11)	7 (7)	6 (12)	6 (7)
Low risk of COVID-19 infection	17 (22)	12 (24)	65 (13)	32 (21)	28 (29)	8 (16)	11 (12)
Interaction with patients with suspected/confirmed SARS-CoV-2, n (%)	22 (29)	16 (33)	226 (45)	47 (31)	31 (33)	15 (29)	34 (38)
Prevalence of suspected/confirmed SARS-CoV-2 based on symptoms±PCR testing, n (%)	10 (13)	14 (29)	84 (17)	17 (11)	10 (11)	2 (4)	6 (7)
Respondents satisfied with local unit SARS-CoV-2 response, n (%)	57 (75)	33 (67)	326 (66)	112 (74)	65 (68)	38 (75)	60 (67)
Mean % level of respondent satisfaction with government SARS-CoV-2 response (95% CI)	63 (56 to 69)	68 (60 to 76)	50 (47 to 52)	48 (43 to 53)	42 (36 to 48)	40 (32 to 47)	47 (41 to 53)

\*All possible combinations are not included in this table; therefore, the denominator of the table is not 2233 (n=1010). Main relevant PPE combinations are described. PCR, Polymerase Chain Reaction; PPE, personal protective equipment.

more likely to interact with patients with suspected or confirmed SARS-CoV-2 and have greater PPE concerns with poorer satisfaction and support. The most common PPE combination in use by HCPs were gloves and a surgical mask, with 18% of patients reporting not wearing a mask. Prevalence of SARS-CoV-2 was lower among HCPs when visors, filtering face masks or full PPE was used. Up to one in five respondents met the criteria for either moderate to severe anxiety or depression. Our findings suggest that suspected or confirmed SARS-CoV-2 prevalence among HCPs working with ultrasound in obstetrics and gynaecology was at least 13% at the time of the survey, with a higher prevalence in the UK.

### Strengths

The main strengths of our study are the relatively large sample size, the assessment of HCP responses internationally and the ability to compare responses between different HCPs. We translated the survey into four languages to include as many potential participants as possible. Furthermore, the response rate of ISUOG members was greater than non-ISUOG members, as expected, given the target population. We compared responses between ISUOG and non-ISUOG members to ensure the data could be reliably grouped for the analysis.

### Weaknesses

The key weakness of the study was the relatively low response rate and the method of sample selection that introduces possible selection bias when interpreting the results. As this is a descriptive study discussing early experience and well-being of HCPs working with ultrasound in obstetrics and gynaecology during the evolving pandemic, it is important not to draw conclusions about causation, one reason why univariate analyses were performed. For example, we cannot link use of PPE to levels of anxiety and depression in this study. We combined suspected and confirmed SARS-CoV-2, at the time of writing, as the majority of HCP did not have the same access to a swab or antibody test that is now available.

### COVID-19 prevalence

Although international prevalence varies, at the time of writing, confirmed global SARS-CoV-2 cases had surpassed 90 million. The suspected or confirmed SARS-CoV-2 prevalence of 13% in this international study is consistent with other published reports on SARS-CoV-2 and HCP, where prevalence ranges from 1.6% to 22%.<sup>16–22</sup> These studies have now been conducted across multiple specialities in many countries, with prevalence determined by PCR as well as antibody testing. One UK study within the field of obstetrics and gynaecology identified an HCP SARS-CoV-2 prevalence of 22%.<sup>18</sup> Given that the advice for use of PPE is consistent within obstetrics and gynaecology,<sup>6</sup> the higher prevalence may relate to respondents not using PPE or not using it correctly. The prevalence suggested by our dataset may be associated with PPE supply concerns, the need to recycle or reuse PPE, a lack

of guidance regarding when to use various levels of PPE, or due to many respondents experiencing high-risk interactions with patients, for example, scanning in proximity for prolonged periods of time.

### PPE use and provision

Countries with greatest concern for PPE from our survey included Spain, Italy, the Philippines and the USA, with similarly high rates of PPE recycling. In the UK and India, over half of respondents expressed concern. The SARS-CoV-2 pandemic led to imposed export restrictions as domestic needs increased, demand shock, rush of PPE acquisition and supply chain failures with variation of national healthcare advice.<sup>23–26</sup> As an example, China was responsible for mass production of clinical gowns, manufacturing 50% of the world's demand prepandemic.<sup>23</sup> This all contributed to the soaring global costs of PPE and the financial consequences of obtaining the necessary materials to safely support each healthcare system. In the USA, gowns were 7.5 times more expensive in 2020 compared with 2019.<sup>24</sup> Face masks were nine times more costly; N95 respirators were priced eight times the prepandemic rate; and gloves cost 2.5 times more.<sup>24</sup> PPE concern and recycling may have been a consequence of the logistical and financial complications seen as part of the SARS-CoV-2 response. However, as each healthcare system had varying prepandemic funding concerns and constraints, as well as varying HCP advice, these may have also compounded PPE supply and distribution further at a time of extreme need.

More trainee doctors reported working in higher-risk areas, as well as being more likely to interact with patients with suspected or confirmed SARS-CoV-2 than consultants or sonographers. They also had higher levels of concern regarding PPE, reusing and recycling more PPE, with a poorer perception of unit and governmental satisfaction and support. It is well documented that HCP burnout relates to anxiety and depression and that burnout is particularly prevalent among trainees.<sup>27</sup> Fears of infection and death during the pandemic may exacerbate this problem<sup>28</sup> and may also relate to inadequate support or protection.

Sonographers also reported seeing more patients who did not wear a mask. The survey shows a worrying pattern where HCPs with the least agency (trainees and sonographers) were not only more likely to see more high-risk patients but were less likely to be protected. It is perhaps not surprising they expressed greater concerns than other colleagues. PPE issues may relate to systematic financial healthcare consequences previously described but, in this instance, may also have local aetiology. Those with least agency also tend to have the least power to instigate change in their department. Concerns may never reach managers and administrators who also may not be clinically trained or orientated. Communication failure may have led to the PPE concerns expressed by HCPs.

Consensus statements and national and international guidance have been published on advised levels of PPE in

relation to SARS-CoV-2, with increasing requirements based on clinical risk. At the top end of the scale, disposable gloves, disposable fluid-repellent coverall/gown, filtering face piece respirator and eye/face protection should be used when performing aerosol-generating procedures on possible or confirmed cases of SARS-CoV-2.<sup>6 29 30</sup> Of the study participants, 22.3% reported using gloves and a surgical mask, the combination recommended by the ISUOG consensus statement for HCPs performing ultrasound scans.<sup>6</sup> Others reported using various combinations of PPE, some of which is recommended for high-risk, aerosol-generating patient–HCP interactions. However, a proportion also reported not using PPE. This variation in PPE may be due to local policy, dependent on wards or clinical areas where HCP duties extend beyond ultrasound, or a result of the PPE supply issues described previously. The use of full PPE or the addition of eye/face protection and/or filtering face piece respirators to gloves is associated with lower reports of suspected or confirmed SARS-CoV-2 when compared with gloves and surgical mask alone. This corroborates literature findings.<sup>19 22</sup>

### Anxiety and depression

In our study, just under half of the survey respondents completed the HADS questionnaire. This may have been due to a language barrier. However, there is also stigma related to HCPs seeking help and support, even though it is known HCPs experience high levels of psychological morbidity.<sup>31–34</sup> This may have discouraged HCPs to provide information relating to anxiety and depression.

From those who completed the HADS, almost half the respondents reported some level of anxiety and depression, with 19.8% and 8.8% experiencing moderate-to-severe anxiety and depression, respectively. Greater proportions were among female participants, with more depression reported by those working in lower-risk areas. Again, this may relate to the stigma for HCPs reporting psychological morbidity and seeking support, but also may be explained by a completion bias, that is, those with psychological morbidity may be more inclined to complete the HADS to increase awareness of these issues.<sup>31–34</sup>

Given the design of this study, we must be careful not to relate this to COVID-19 and be aware that there may be selection bias. There are aspects of the obstetrics and gynaecology environment which may encourage higher pre-existing levels of anxiety and depression among HCPs prior to the SARS-CoV-2 pandemic. The investigating traumatic work-related events in obstetrics and gynaecology (INDIGO) study reports two-thirds of obstetrics and gynaecology trainees and consultants in the UK have been personally traumatised by work-related events unrelated to SARS-CoV-2, with 31% of them affected by post-traumatic stress disorder symptoms.<sup>35</sup> Of the total paediatricians, 14.1% and 7.3% reported mild to severe anxiety and depression using HADS in a survey published prior to the pandemic,<sup>36</sup> a specialty that reflects similar intensity.

Our findings strongly support the need for reliable infrastructure that provide HCP counselling and psychological

support without stigma in every medical and surgical specialty. In the UK, services such as NHS Practitioner Health and the British Medical Association offer confidential support and protect HCPs at the national level, while locally, seniors provide regular debriefing sessions. However, provision of local and global support is not consistent, and thus movement beyond a culture of HCP stigma is essential.<sup>31</sup>

### CONCLUSION

This study provides insight into the experience and well-being of clinicians working in the field of ultrasound in obstetrics and gynaecology during the early phase of the SARS-CoV-2 pandemic. Although further work is required to unpick the associations of the pandemic, PPE availability, SARS-CoV-2 prevalence, and anxiety and depression, we hope our report highlights the importance of provision of PPE and the need for national and international consistencies in advice regarding PPE requirements, regulations and use. HCPs all take risk daily, particularly trainees and sonographers who are generally more exposed to SARS-CoV-2 and less unable to communicate their concerns and needs. As many patients may harbour infection with minimal or without symptoms, there is a real need to ensure HCPs are consistently and adequately supported.

### Author affiliations

<sup>1</sup>Early Pregnancy and Acute Gynaecology Unit, Department of Obstetrics and Gynaecology, Queen Charlotte's and Chelsea Hospital, Imperial College London, London, UK

<sup>2</sup>Department of Development and Regeneration, KU Leuven, Leuven, Belgium

<sup>3</sup>Department of Oncology, Laboratory of Tumour Immunology and Immunotherapy, KU Leuven, Leuven, Belgium

<sup>4</sup>Facultad de Medicina, Clínica Alemana, Universidad del Desarrollo, Santiago de Chile, Chile

<sup>5</sup>Facultad de Medicina, Clínica Alemana, Hospital Clínico de la Universidad de Chile José Joaquín Aguirre, Santiago, Chile

<sup>6</sup>Département d'échographie en Gynécologie et Obstétrique, Centre d'Échographie de l'Odéon, Paris, France

<sup>7</sup>Dipartimento Scienze della Salute della Donna, del Bambino e di Sanità Pubblica, Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Roma, Italy

<sup>8</sup>Centre for Fetal Care, Department of Obstetrics and Gynaecology, Queen Charlotte's and Chelsea Hospital, Imperial College London, London, UK

<sup>9</sup>Department of Obstetrics and Gynaecology, KU Leuven University Hospitals Leuven, Leuven, Belgium

**Twitter** Tom Bourne @proftombourne, Christopher Kyriacou @ChristopherKyr4, Harsha Shah @Harsha\_Shah86 and Christoph Lees @Christoph\_Lees

**Contributors** TB, CK, HS and DT participated in the conception and design of the study. TB, CK and HS created the English version of the survey. JP, UM and CL translated survey to Spanish, French and Italian. JC performed the statistical analysis. TB, CK, HS, JC and DT interpreted the results. TB, CK and HS wrote the initial version of the manuscript. All authors critically revised the manuscript and approved the final version. TB is the author acting as guarantor.

**Funding** TB is supported by the National Institute for Health Research Imperial Biomedical Research Centre (grant number IS-BRC-1215–20013). CK is supported by Imperial Health Charity (grant number RFPD1920/116). DT is fundamental clinical researcher of Research Foundation – Flanders (FWO). JC and DT are supported by FWO (grant number G0B4716N) and Internal Funds KU Leuven (grant number C24/15/037).

**Competing interests** None declared.

**Patient consent for publication** Not applicable.



**Ethics approval** This study involves human participants. Ethical approval for the distribution of this online survey was obtained from the ethics committee of KU Leuven, Belgium (S-64016). Participants gave informed consent to participate in the study before taking part. A copy of the protocol can be viewed in online supplemental material 2.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All data relevant to the study are included in the article or uploaded as supplemental information. No additional data available.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

#### ORCID iDs

Tom Bourne <http://orcid.org/0000-0003-1421-6059>

Christopher Kyriacou <http://orcid.org/0000-0001-9001-5545>

Harsha Shah <http://orcid.org/0000-0003-3866-7946>

Chiara Landolfo <http://orcid.org/0000-0001-9808-7957>

Christoph Lees <http://orcid.org/0000-0002-2104-5561>

Dirk Timmerman <http://orcid.org/0000-0002-3707-6645>

#### REFERENCES

- 1 The Lancet. COVID-19: protecting health-care workers. *The Lancet* 2020;395:922.
- 2 Adams JG, Walls RM. Supporting the health care workforce during the COVID-19 global epidemic. *JAMA* 2020;323:1439.
- 3 Abu-Rustum RS, Akolekar R, Sotiriadis A, et al. ISUOG consensus statement on organization of routine and specialist obstetric ultrasound services in context of COVID-19. *Ultrasound Obstet Gynecol* 2020;55:863–70.
- 4 Bourne T, Kyriacou C, Coomarasamy A, et al. ISUOG consensus statement on rationalization of early-pregnancy care and provision of ultrasonography in context of SARS-CoV-2. *Ultrasound Obstet Gynecol* 2020;55:871–8.
- 5 Wilson AN, Ravalid C, Scoullar MJL, et al. Caring for the carers: ensuring the provision of quality maternity care during a global pandemic. *Women Birth* 2021;34:206–9.
- 6 Abramowicz JS, Basseal JM, Brezinka C, et al. ISUOG safety Committee position statement on use of personal protective equipment and hazard mitigation in relation to SARS-CoV-2 for practitioners undertaking obstetric and gynecological ultrasound. *Ultrasound Obstet Gynecol* 2020;55:886–91.
- 7 Lai J, Ma S, Wang Y, et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Netw Open* 2020;3:e203976.
- 8 Lee AM, Wong JGWS, McAlonan GM, et al. Stress and psychological distress among SARS survivors 1 year after the outbreak. *Can J Psychiatry* 2007;52:233–40.
- 9 Bansal P, Bingemann TA, Greenhawt M, et al. Clinician wellness during the COVID-19 pandemic: extraordinary times and unusual challenges for the Allergist/Immunologist. *J Allergy Clin Immunol Pract* 2020;8:1781–90.
- 10 Albott CS, Wozniak JR, McGlinch BP, et al. Battle Buddies: rapid deployment of a psychological resilience intervention for health care workers during the COVID-19 pandemic. *Anesth Analg* 2020;131:43–54.
- 11 Shanafelt T, Ripp J, Trockel M. Understanding and addressing sources of anxiety among health care professionals during the COVID-19 pandemic. *JAMA* 2020;323:2133.
- 12 Dewey C, Hingle S, Goelz E, et al. Supporting clinicians during the COVID-19 pandemic. *Ann Intern Med* 2020;172:752–3.
- 13 WHO. WHO coronavirus disease (COVID-19). Dashboard: Word Health Organization, 2020. <https://covid19.who.int/>
- 14 Yang J, Zheng Y, Gou X, et al. Prevalence of comorbidities and its effects in patients infected with SARS-CoV-2: a systematic review and meta-analysis. *Int J Infect Dis* 2020;94:91–5.
- 15 Zigmund AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361–70.
- 16 Kluytmans-van den Bergh MFQ, Buiting AGM, Pas SD, et al. Prevalence and clinical presentation of health care workers with symptoms of coronavirus disease 2019 in 2 Dutch hospitals during an early phase of the pandemic. *JAMA Netw Open* 2020;3:e209673.
- 17 Sikkema RS, Pas SD, Nieuwenhuijsen DF, et al. COVID-19 in health-care workers in three hospitals in the South of the Netherlands: a cross-sectional study. *Lancet Infect Dis* 2020;20:1273–80.
- 18 Khalil A, Hill R, Wright A, et al. SARS-CoV-2-Specific antibody detection in healthcare workers in a UK maternity Hospital: correlation with SARS-CoV-2 RT-PCR results. *Clin Infect Dis* 2021;72:1680–1.
- 19 Korth J, Wilde B, Doff S, et al. SARS-CoV-2-specific antibody detection in healthcare workers in Germany with direct contact to COVID-19 patients. *J Clin Virol* 2020;128:104437.
- 20 Rivett L, Sridhar S, Sparkes D, et al. Screening of healthcare workers for SARS-CoV-2 highlights the role of asymptomatic carriage in COVID-19 transmission. *eLife* 2020;9.
- 21 Chen Y, Tong X, Wang J, et al. High SARS-CoV-2 antibody prevalence among healthcare workers exposed to COVID-19 patients. *J Infect* 2020;81:420–6.
- 22 Chatterjee P, Anand T, Singh KJ, et al. Healthcare workers & SARS-CoV-2 infection in India: A case-control investigation in the time of COVID-19. *Indian J Med Res* 2020;151:459–67.
- 23 Burki T. Global shortage of personal protective equipment. *Lancet Infect Dis* 2020;20:785–6.
- 24 Mena Lora A, Ali M, Spencer S, et al. Cost of personal protective equipment during the first wave of COVID-19. *ASHE* 2021;1:s49.
- 25 Walters W, Collins T, Evans A. Additional cost in personal protective equipment to NHS services during COVID-19: a review of the expense incurred by aneurin BEVAN Healthboards maxillofacial unit over a one-year period to meet current clinical guidelines. *Br J Oral Maxillofac Surg* 2020;58:e320–2.
- 26 Rowan NJ, Laffey JG. Challenges and solutions for addressing critical shortage of supply chain for personal and protective equipment (PPE) arising from coronavirus disease (COVID19) pandemic – case study from the Republic of Ireland. *Sci Total Environ* 2020;725:138532.
- 27 Bourne T, Shah H, Falconieri N, et al. Burnout, well-being and defensive medical practice among obstetricians and gynaecologists in the UK: cross-sectional survey study. *BMJ Open* 2019;9:e030968.
- 28 Hu D, Kong Y, Li W, et al. Frontline nurses' burnout, anxiety, depression, and fear statuses and their associated factors during the COVID-19 outbreak in Wuhan, China: a large-scale cross-sectional study. *EClinicalMedicine* 2020;24:100424.
- 29 PHE. COVID-19 personal protective equipment (PPE), 2020. Available: <https://www.gov.uk/government/publications/wuhan-novel-coronavirus-infection-prevention-and-control/covid-19-personal-protective-equipment-ppe>
- 30 WHO. Rational use of personal protective equipment for coronavirus disease (COVID-19) and considerations during severe shortages, 2020. Available: [https://www.who.int/publications/i/item/rational-use-of-personal-protective-equipment-for-coronavirus-disease-\(covid-19\)-and-considerations-during-severe-shortages](https://www.who.int/publications/i/item/rational-use-of-personal-protective-equipment-for-coronavirus-disease-(covid-19)-and-considerations-during-severe-shortages)
- 31 Knaak S, Mantler E, Szeto A. Mental illness-related stigma in healthcare: barriers to access and care and evidence-based solutions. *Health Manage Forum* 2017;30:111–6.
- 32 Adams EFM, Lee AJ, Pritchard CW, et al. What stops us from healing the healers: a survey of help-seeking behaviour, stigmatisation and depression within the medical profession. *Int J Soc Psychiatry* 2010;56:359–70.
- 33 Wallace JE. Mental health and stigma in the medical profession. *Health* 2012;16:3–18.
- 34 Wijeratne C, Johnco C, Draper B, et al. Doctors' reporting of mental health stigma and barriers to help-seeking. *Occup Med* 2021;71:366–74.
- 35 Slade P, Balling K, Sheen K, et al. Work-related post-traumatic stress symptoms in obstetricians and gynaecologists: findings from INDIGO, a mixed-methods study with a cross-sectional survey and in-depth interviews. *BJOG* 2020;127:600–8.
- 36 van Steijn ME, Scheepstra KWF, Yasar G, et al. Occupational well-being in pediatricians-a survey about work-related posttraumatic stress, depression, and anxiety. *Eur J Pediatr* 2019;178:681–93.



## SUPPLEMENTARY MATERIAL

### The experiences and wellbeing of healthcare professionals working in the field of ultrasound in Obstetrics and Gynaecology as the SARS-CoV-2 pandemic was evolving: a cross-sectional survey study

#### Authors

Tom Bourne<sup>1,2</sup>, Christopher Kyriacou<sup>1</sup>, Harsha Shah<sup>1</sup>, Jolien Ceusters<sup>3</sup>, Jessica Preisler<sup>4,5</sup>, Ulrike Metzger<sup>6</sup>, Chiara Landolfo<sup>7</sup>, Christoph Lees<sup>8</sup>, Dirk Timmerman<sup>2,9</sup>

#### Affiliations

1. Early pregnancy and Acute Gynaecology Unit, Department of Obstetrics and Gynaecology, Queen Charlotte's and Chelsea Hospital, Imperial College, London, UK.
2. Department of Development and Regeneration, KU Leuven, Leuven, Belgium.
3. Department of Oncology, Laboratory of Tumour Immunology and Immunotherapy, KU Leuven, Leuven, Belgium.
4. Facultad de Medicina, Clínica Alemana – Universidad del Desarrollo, Santiago, Chile
5. Hospital Clínico de la Universidad de Chile, Santiago, Chile.
6. Centre d'Échographie de l'Odéon, Paris, France.
7. Dipartimento Scienze della Salute della Donna, del Bambino e di Sanità Pubblica, Fondazione Policlinico Universitario Agostino Gemelli, IRCCS, Rome, Italy.
8. Centre for Fetal Care, Department of Obstetrics and Gynaecology, Queen Charlotte's and Chelsea Hospital, Imperial College, London, UK.
9. Department of Obstetrics and Gynaecology, University Hospitals Leuven, Leuven, Belgium.

#### Correspondence to:

Professor Tom Bourne  
Queen Charlotte's and Chelsea Hospital, Imperial College London  
Du Cane Road, London, W12 0HS, UK  
[t.bourne@imperial.ac.uk](mailto:t.bourne@imperial.ac.uk)

**Supplement 1: The International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) SARS-CoV-2 Questionnaire for healthcare professionals working with ultrasound in Obstetrics and Gynecology.**

**Informed consent**

This questionnaire is fully anonymous - answers cannot be traced back to participants. The anonymised data will be collected for analysis in order to provide a report and publication at a later date.

**1. Please click 'yes' to participate in this survey. Otherwise, click 'no': (please choose one option)**

Yes

No

**Introduction**

**2. What title best describes your job: (please choose one option)**

Sonographer – Obstetrics

Sonographer – Gynecology

Sonographer – Obstetrics and Gynecology

Obstetrician

Gynecologist

Obstetrician and Gynecologist

Radiologist

Doctor in training – Obstetrics

Doctor in training – Gynecology

Doctor in training – Obstetrics and Gynecology

Doctor in training – Radiologist

Doctor in research – Obstetrics

Doctor in research – Gynecology

Doctor in research – Obstetrics and Gynecology

Doctor in research – Radiology

Nurse – Obstetrics

Nurse – Gynecology

Nurse – Obstetrics and Gynecology

Nurse – Early pregnancy only

Midwife – Obstetrics

Midwife – Early pregnancy only

Retired

**3. In what country do you practice? (please state below)**

Afghanistan

Albania

Algeria

Andorra

Angola

- 1 Antigua and Barbuda
- 2 Argentina
- 3 Armenia
- 4 Australia
- 5 Austria
- 6 Azerbaijan
- 7 The Bahamas
- 8 Bahrain
- 9 Bangladesh
- 10 Barbados
- 11 Belarus
- 12 Belgium
- 13 Belize
- 14 Benin
- 15 Bhutan
- 16 Bolivia
- 17 Bosnia and Herzegovina
- 18 Botswana
- 19 Brazil
- 20 Brunei
- 21 Bulgaria
- 22 Burkina Faso
- 23 Burundi
- 24 Cabo Verde
- 25 Cambodia
- 26 Cameroon
- 27 Canada
- 28 Central African Republic
- 29 Chad
- 30 Chile
- 31 China
- 32 Colombia
- 33 Comoros
- 34 Congo, Democratic Republic of the
- 35 Congo, Republic of the
- 36 Costa Rica
- 37 Côte d'Ivoire
- 38 Croatia
- 39 Cuba
- 40 Cyprus
- 41 Czech Republic
- 42 Denmark
- 43 Djibouti
- 44 Dominica
- 45 Dominican Republic
- 46 East Timor (Timor-Leste)
- 47 Ecuador

- 1 Egypt
- 2 El Salvador
- 3 Equatorial Guinea
- 4 Eritrea
- 5 Estonia
- 6 Eswatini
- 7 Ethiopia
- 8 Fiji
- 9 Finland
- 10 France
- 11 Gabon
- 12 The Gambia
- 13 Georgia
- 14 Germany
- 15 Ghana
- 16 Greece
- 17 Grenada
- 18 Guatemala
- 19 Guinea
- 20 Guinea-Bissau
- 21 Guyana
- 22 Haiti
- 23 Honduras
- 24 Hungary
- 25 Iceland
- 26 India
- 27 Indonesia
- 28 Iran
- 29 Iraq
- 30 Ireland
- 31 Israel
- 32 Italy
- 33 Jamaica
- 34 Japan
- 35 Jordan
- 36 Kazakhstan
- 37 Kenya
- 38 Kiribati
- 39 Korea, North
- 40 Korea, South
- 41 Kosovo
- 42 Kuwait
- 43 Kyrgyzstan
- 44 Laos
- 45 Latvia
- 46 Lebanon
- 47 Lesotho



- 1 Liberia
- 2 Libya
- 3 Liechtenstein
- 4 Lithuania
- 5 Luxembourg
- 6 Madagascar
- 7 Malawi
- 8 Malaysia
- 9 Maldives
- 10 Mali
- 11 Malta
- 12 Marshall Islands
- 13 Mauritania
- 14 Mauritius
- 15 Mexico
- 16 Micronesia, Federated States of
- 17 Moldova
- 18 Monaco
- 19 Mongolia
- 20 Montenegro
- 21 Morocco
- 22 Mozambique
- 23 Myanmar (Burma)
- 24 Namibia
- 25 Nauru
- 26 Nepal
- 27 Netherlands
- 28 New Zealand
- 29 Nicaragua
- 30 Niger
- 31 Nigeria
- 32 North Macedonia
- 33 Norway
- 34 Oman
- 35 Pakistan
- 36 Palau
- 37 Panama
- 38 Papua New Guinea
- 39 Paraguay
- 40 Peru
- 41 Philippines
- 42 Poland
- 43 Portugal
- 44 Qatar
- 45 Romania
- 46 Russia
- 47 Rwanda

- 1 Saint Kitts and Nevis
- 2 Saint Lucia
- 3 Saint Vincent and the Grenadines
- 4 Samoa
- 5 San Marino
- 6 Sao Tome and Principe
- 7 Saudi Arabia
- 8 Senegal
- 9 Serbia
- 10 Seychelles
- 11 Sierra Leone
- 12 Singapore
- 13 Slovakia
- 14 Slovenia
- 15 Solomon Islands
- 16 Somalia
- 17 South Africa
- 18 Spain
- 19 Sri Lanka
- 20 Sudan
- 21 Sudan, South
- 22 Suriname
- 23 Sweden
- 24 Switzerland
- 25 Syria
- 26 Taiwan
- 27 Tajikistan
- 28 Tanzania
- 29 Thailand
- 30 Togo
- 31 Tonga
- 32 Trinidad and Tobago
- 33 Tunisia
- 34 Turkey
- 35 Turkmenistan
- 36 Tuvalu
- 37 Uganda
- 38 Ukraine
- 39 United Arab Emirates
- 40 United Kingdom
- 41 United States
- 42 Uruguay
- 43 Uzbekistan
- 44 Vanuatu
- 45 Vatican City
- 46 Venezuela
- 47 Vietnam

Yemen  
Zambia  
Zimbabwe

**4. Are you an ISUOG member? (please choose one option)**

Yes  
No

**5. If you answered no, which society or professional body did you receive this survey from?**

Open box answer

**Demographics**

**6. How old are you? (please state number of years)**

Numbers offered

**7. What sex were you assigned at birth? (please choose one option)**

Male  
Female  
Intersex  
Prefer not to say

**8. What gender do you most identify with? (please choose one option)**

Male  
Female  
Prefer not to say  
Other (please specify)

**Underlying medical conditions**

**9. Do you have any underlying medical conditions that place you at increased risk during the SARS-CoV-2 pandemic? (please tick all that apply)**

None  
Heart disease  
Hypertension  
Diabetes  
Asthma  
Other lung disease  
Kidney disease  
Cancer

**Medication****10. Are you taking any of the following medications? (please tick all that apply)**

None

Immunosuppressant medication - steroids

Immunosuppressant medication - methotrexate

Immunosuppressant medication - biological agents

Aspirin

Indomethacin

Any other non-steroidal anti-inflammatory drugs (NSAIDs) e.g. ibuprofen, diclofenac, naproxen

Angiotensin converting enzyme inhibitors (ACEi) blood pressure medication - any ending in - pril (e.g. enalapril)

Any other blood pressure medication e.g. angiotensin receptor blockers (ARB), alpha blockers, alpha-2 receptor agonists, beta blockers, combined alpha and beta blockers, calcium channel blockers (CCB), diuretics, other

**Smoking and BMI****11. Do you smoke? (please choose one option)**

Yes

Previously – now stopped

No

**12. What was your last measured weight? (please choose one option)***In kilograms (kg)* – numbers offered**13. What is your height? (please choose one option)***In centimetres (cm)* – numbers offered**Role, shift pattern, PPE****14. How has your role and shift pattern been affected by the SARS-CoV-2 pandemic? (please choose one option)**

Unable to work for the duration of the pandemic

Reduced duties with no patient contact

Reduced duties with reduced patient contact

Reduced duties with similar patient contact

Reduced duties with increased patient contact

Similar duties with no patient contact

Similar duties with reduced patient contact

Similar duties with similar patient contact

Similar duties with increased patient contact

Increased duties with no patient contact

Increased duties with reduced patient contact



1 Increased duties with similar patient contact  
2 Increased duties with increased patient contact  
3

4 **15. Where are you working during the SARS-CoV-2 pandemic? (please tick all that  
5 apply)**

6 Obstetrics – Labor ward  
7 Obstetrics – Birth center  
8 Obstetrics – Antenatal ward  
9 Obstetrics – Postnatal ward  
10 Obstetrics – Maternity triage  
11 Obstetrics – Antenatal clinic  
12 Obstetrics – Maternity helplines  
13 Obstetrics – Community clinics  
14 Obstetrics – Ultrasound  
15 Gynecology – Acute gynecology  
16 Gynecology – Benign gynecology  
17 Gynecology – Oncology  
18 Gynecology – Ultrasound  
19 Early pregnancy unit – Ultrasound and assessment  
20 Radiology – Ultrasound  
21 Radiology – non-ultrasound  
22 Obstetrics/Gynecology/Radiology – From home  
23

24 **16. Do you interact with patients confirmed or suspected to have SARS-CoV-2? (please  
25 choose one option)**

26 Yes – every day  
27 Yes – every week  
28 Yes – every so often  
29 I don't know  
30 Not yet  
31 No  
32

33 **17. Have you been using personal protective equipment (PPE) since the SARS-CoV-2  
34 outbreak? (please choose one option)**

35 No  
36 Not yet  
37 Yes – for aerosol generating procedures only  
38 Yes – for suspected/positive patients only  
39 Yes – for all patients  
40

41 **18. When you perform an ultrasound scan please state what PPE you use: (please tick  
42 all that apply)**

43 None  
44 Gloves  
45 Surgical mask  
46 N95/FFP3 mask  
47 Gown

1 Visor

2 Surgical hat

3

4 **19. Have there been any concerns regarding PPE provision or shortages since the**  
5 **SARS-CoV-2 outbreak? (note – Full PPE includes a N95/FFP3 mask, visor, surgical**  
6 **hat, gown and gloves) (please tick all that apply)**

7 No issues with provision

8 Yes – supply issues with gloves

9 Yes – supply issues with surgical masks

10 Yes – supply issues with N95/FFP3 masks

11 Yes – supply issues with gowns

12 Yes – supply issues with visors

13 Yes – supply issues with surgical hats

14 Yes – supply issues with providing patients with a surgical mask

15

16 **20. Have you been asked to re-use or recycle any PPE? (please tick all that apply**  
17 **option)**

18 No

19 Yes - Gloves

20 Yes - Surgical mask

21 Yes - N95/FFP3 mask

22 Yes - Gown

23 Yes - Visor

24 Yes - Surgical hat

25

26 **21. Do you ask patients to wear a surgical mask? (please choose one option)**

27 Yes

28 No

29

30

31 **SARS-CoV-2 diagnosis**

32

33 **22. Have you had or do you have SARS-CoV-2 based on clinical symptoms and/or a**  
34 **SARS-CoV-2 swab (PCR) diagnosis? (please choose one option)**

35 No – no test or symptoms suggestive of SARS-CoV-2

36 Yes – suspected based on symptoms but have not had PCR swab testing

37 Yes – suspected based on symptoms, had PCR testing, which was negative

38 Yes – confirmed on PCR swab testing but have had no symptoms

39 Yes – suspected based on symptoms and confirmed on PCR swab testing

40

41 **23. With regards to having a SARS-CoV-2 swab test, please choose one option:**

42 Not been offered

43 Offered as experiencing symptoms

44 Offered as in contact with confirmed or probable SARS-CoV-2 patients

45 Offered as living with someone who has probable or confirmed SARS-CoV-2

46 Not applicable

47

**24. What symptoms have you experienced or are you experiencing? (please tick all that apply)**

- None
- High fever
- Persistent dry cough
- Productive cough
- Shortness of breath/Difficulty breathing
- Fatigue
- Headache
- Loss of sense of smell and/or taste
- Sore throat
- Chest pain
- Diarrhoea
- Myalgia
- Loss of appetite
- Hoarse voice
- Abdominal pain

**Personal SARS-CoV-2 pathway**

**25. At what point were you offered a SARS-CoV-2 swab? (please state number of days into your illness with day 1 being the day you started to have symptoms. If you have not had symptoms – please choose 'not had symptoms'. If you were not offered a test, please choose 'not offered a test')**

- Not had symptoms
- Not offered a test
- Not applicable
- Numbers offered

**26. How severe was/is your infection? (please choose one option)**

- Mild COVID-19 (T< 37.8 oC, tiredness, muscle aches, headache, no breathlessness, normal appetite)
- Moderate COVID-19 (T> 37.8 oC, troublesome cough, breathless on exercise, tired but no chest or abdominal pain)
- Severe COVID-19 (T> 37.8 oC , noticeably breathless even when sitting still, unable to complete sentences when talking, chest or abdominal pain, reduced appetite)
- Critical COVID-19 (development of severe acute respiratory syndrome in need of ventilator or other respiratory support)
- Not applicable

**27. How long were you or have you been symptomatic for? (please state number of days. If you have not had symptoms – please choose 'not had symptoms')**

- Not had symptoms
- Numbers offered

1       **28. How long did you stay at home before going back to work, or if current infection,**  
2       **how long have you been at home for? (please state number of days. If you have**  
3       **not had symptoms – please choose 'not had symptoms')**

4       Not had symptoms

5       Numbers offered

7       **29. Did you or do you feel under pressure from your employers to return to work**  
8       **before making a full recovery? (please choose one option)**

9       Yes

10      No

11      Not applicable

13      **30. If you have been unwell with presumed or confirmed SARS-CoV-2, how long were**  
14      **you advised to stay at home by your employers before going back to work? (please**  
15      **state number of days. If you have not had symptoms – please choose 'not had**  
16      **symptoms')**

17      Not had symptoms

18      Numbers offered

20      **31. Have you had a SARS-CoV-2 antibody test? (please choose one option)**

21      No – no test or symptoms suggestive of SARS-CoV-2

22      No – not offered

23      No – offered and declined

24      Yes – negative

25      Yes – positive

28      **Household**

30      **32. Have you needed to, or are you currently self-isolating due to a member of your**  
31      **household having suspected or confirmed SARS-CoV-2? (please choose one option)**

32      Yes

33      No

35      **33. How long did you need or have you needed to self-isolate due to a member of your**  
36      **household having suspected or confirmed SARS-CoV-2? (please state number of**  
37      **days. If your household has been well - please choose 'household has been well')**

38      Household has been well

39      Numbers offered



**Colleagues**

**34. How many colleagues do you work with in your unit? (please state number of each group. Select 'unsure' if you do not know)**

*Sonographers*

*Consultants*

*Doctors in training*

*Doctors in research*

*Nurses*

*Midwives*

Unsure

Numbers offered

**35. To your knowledge, how many colleagues are you aware of in your unit that have had a presumed diagnosis of SARS-CoV-2 based on symptoms or a positive swab? (please state number of each group. Select 'unsure' if you have no information)**

*Sonographers*

*Consultants*

*Doctors in training*

*Doctors in research*

*Nurses*

*Midwives*

Unsure

Numbers offered

**36. To your knowledge, how many colleagues have needed to self-isolate or are self-isolating due to a member of their household having suspected or confirmed SARS-CoV-2? (please state number of each group. Select 'unsure' if you have no information)**

*Sonographers*

*Consultants*

*Doctors in training*

*Doctors in research*

*Nurses*

*Midwives*

Unsure

Numbers offered

**37. What is the longest period of time any of your colleagues have taken off in isolation because they or a household member have suspected or confirmed SARS-CoV-2? (please state number of days. Select 'unsure' if you have no information)**

Unsure

Numbers offered

**Support**

**38. Have you felt or do you feel adequately supported by your unit during the SARS-CoV-2 pandemic? (please choose one option)**

Yes

No

**39. How do you rate the response and support that was provided by your country's Ministry of Health or Public Health officials? (please answer using the scale below)**

0 - Extremely poor

50 - Adequate

100 - Excellent

**40. Please tell us, if you wish, your views regarding how well clinician safety has been protected by your employers (please state in the text box below. If you choose not to answer, please write 'not applicable')**

Open box answer

**Hospital Anxiety and Depression Scale (HADS)**

**41. Thank you for filling out this questionnaire.**

**Optional: Hospital Anxiety and Depression Scale (HADS). If you are willing to proceed, please click yes. Otherwise, click no to end the survey.**

Yes

No

**Hospital Anxiety and Depression Scale (HADS)**

**42. Choose the reply that is closest to how you have been feeling in the past week. Don't take too long over your replies: your immediate is best.**

**I feel tense or 'wound up':**

Most of the time

A lot of the time

From time to time, occasionally

Not at all

**43. Choose the reply that is closest to how you have been feeling in the past week. Don't take too long over your replies: your immediate is best.**

**I still enjoy the things I used to enjoy:**

Definitely as much

Not quite as much

- 1 Only a little  
2 Hardly at all  
3
- 4 **44. Choose the reply that is closest to how you have been feeling in the past week.**  
5 **Don't take too long over your replies: your immediate is best.**  
6
- 7 **I get a sort of frightened feeling as if something awful is about to happen:**  
8 Very definitely and quite badly  
9 Yes, but not too badly  
10 A little, but it doesn't worry me  
11 Not at all  
12
- 13 **45. Choose the reply that is closest to how you have been feeling in the past week.**  
14 **Don't take too long over your replies: your immediate is best.**  
15
- 16 **I can laugh and see the funny side of things:**  
17 As much as I always could  
18 Not quite so much now  
19 Definitely not so much now  
20 Not at all  
21
- 22 **46. Choose the reply that is closest to how you have been feeling in the past week.**  
23 **Don't take too long over your replies: your immediate is best.**  
24
- 25 **Worrying thoughts go through my mind:**  
26 A great deal of the time  
27 A lot of the time  
28 From time to time, but not too often  
29 Only occasionally  
30
- 31 **47. Choose the reply that is closest to how you have been feeling in the past week.**  
32 **Don't take too long over your replies: your immediate is best.**  
33
- 34 **I feel cheerful:**  
35 Not at all  
36 Not often  
37 Sometimes  
38 Most of the time  
39
- 40 **48. Choose the reply that is closest to how you have been feeling in the past week.**  
41 **Don't take too long over your replies: your immediate is best.**  
42
- 43 **I can sit at ease and feel relaxed:**  
44 Definitely  
45 Usually  
46 Not often  
47 Not at all

- 1  
2       **49. Choose the reply that is closest to how you have been feeling in the past week.**  
3       **Don't take too long over your replies: your immediate is best.**  
4  
5       **I feel as if I am slowed down:**  
6       Nearly all the time  
7       Very often  
8       Sometimes  
9       Not at all  
10  
11       **50. Choose the reply that is closest to how you have been feeling in the past week.**  
12       **Don't take too long over your replies: your immediate is best.**  
13  
14       **I get a sort of frightened feeling like 'butterflies' in the stomach:**  
15       Not at all  
16       Occasionally  
17       Quite often  
18       Very often  
19  
20       **51. Choose the reply that is closest to how you have been feeling in the past week.**  
21       **Don't take too long over your replies: your immediate is best.**  
22  
23       **I have lost interest in my appearance:**  
24       Definitely  
25       I don't take as much care as I should  
26       I may not take quite as much care  
27       I take just as much care as ever  
28  
29       **52. Choose the reply that is closest to how you have been feeling in the past week.**  
30       **Don't take too long over your replies: your immediate is best.**  
31  
32       **I feel restless as I have to be on the move:**  
33       Very much indeed  
34       Quite a lot  
35       Not very much  
36       Not at all  
37  
38       **53. Choose the reply that is closest to how you have been feeling in the past week.**  
39       **Don't take too long over your replies: your immediate is best.**  
40  
41       **I look forward with enjoyment to things:**  
42       As much as I ever did  
43       Rather less than I used to  
44       Definitely less than I used to  
45       Hardly at all  
46  
47

- 1       **54. Choose the reply that is closest to how you have been feeling in the past week.**  
2       **Don't take too long over your replies: your immediate is best.**  
3  
4       **I get sudden feelings of panic:**  
5       Very often indeed  
6       Quite often  
7       Not very often  
8       Not at all  
9  
10       **55. Choose the reply that is closest to how you have been feeling in the past week.**  
11       **Don't take too long over your replies: your immediate is best.**  
12  
13       **I can enjoy a good book or radio or TV program:**  
14       Often  
15       Sometimes  
16       Not often  
17       Very seldom  
18  
19  
20       **56. Thank you for filling out this questionnaire. Please complete last question to end.**  
21       **Please tell us, if you wish, how you have felt emotionally during the SARS-CoV-2**  
22       **pandemic? (please state in the text box below. If you choose not to answer, please**  
23       **write 'not applicable')**  
24       Open box answer  
25  
26  
27



**Supplement 2: Original study protocol**

**Title:** An international survey of the impact of COVID19 on healthcare professionals carrying out ultrasonography, service provision and mental state in Obstetrics and Gynecology

**Date:** 19<sup>th</sup> April 2020

**Main Sponsor:** UZ Leuven Belgium

**Carried out in collaboration with:** International Society of Ultrasound in Obstetrics and Gynecology (ISUOG)

**Version number:** 1

Principal Investigators: Professor Tom Bourne and Professor Dirk Timmerman

Co-Investigators: Dr Christopher Kyriacou, Dr Harsha Shah

**Protocol to be assessed by:** EC Onderzoek UZ Leuven, Belgium.

**Design:** Prospective cohort observational survey study.

**Aim:** To assess the services and psychological impact of the COVID-19 pandemic to healthcare professionals carrying out ultrasonography within the Obstetrics and Gynecology specialty.

**Hypothesis:**

1. Service provision to support healthcare professionals during the COVID-19 pandemic varies considerably in every unit globally;
2. Healthcare professionals will have experienced COVID-19 first-hand;
3. The COVID-19 pandemic has caused significant psychological distress to healthcare professionals which can be assessed using a validated psychological tool.

**Outcome measures:**

1. A questionnaire to healthcare professionals detailing medical history, role, area of work, personal protective equipment (PPE), COVID-19 patient contact and personal COVID-19 experience will allow data to be collected in relation to the physical impact of the COVID-19 pandemic;
2. Use of the validated Hospital Anxiety and Depression Scale (HADS) will allow psychological morbidity of healthcare professionals to be assessed.

**Eligibility:** Members of ISUOG that fill in the questions via an online survey platform. The questionnaire will in English, Spanish, Greek, Hindu, Mandarin and Cantonese.

**Duration:** The survey will be conducted over a period of two weeks in order to allow time for the questions to be answered, two reminder emails will be sent – one at a week before closure and one 24 hours before closure

**Consent:** Consent will be implied when the questionnaire is filled in and there will be a specific question asking for consent to take part in the survey, however all information is anonymized and cannot be traced to an individual. No data will be requested that is in breach of General Data Protection Regulation (GDPR).

**Background:** The current coronavirus (SARS-CoV-2) pandemic carries a number of new challenges to local, national and international healthcare agencies in order to protect patients and healthcare professionals (physicians, sonographers, allied healthcare professionals). Guidance varies from nation to nation with issues arising from redeployment of clinicians, prioritization of particular services and provision of PPE. This is important as those carrying out ultrasound scans are in close proximity to patients and are at high risk of SARS-CoV-2 infection<sup>1</sup>.

Anxiety and depression are recognized complications for healthcare professionals working in a high-stress environment, particularly one that now encompasses a novel disease for which we have yet to fully understand<sup>2</sup>.

**Methods:** This is a prospective cohort study and will recruit via the dissemination of a survey online through the ISUOG mailing list. Consent will be presumed when the form is filled in and submitted. This will allow healthcare professionals globally to answer. The questionnaire has been devised to assess the service provision in units around the world as well as question personal COVID-19 experience. Following on from this will be an optional HADS. The HADS is a 14-part questionnaire with 7 questions exploring symptoms of anxiety and depression, respectively. Each question is answered with a scale of 0-3 to represent severity of symptoms, with an independent maximum score of 21 for depression and anxiety. A score  $\geq 11$  is deemed moderate, whilst  $\geq 16$  represents severe symptoms<sup>3</sup>.

**Statistics and data analysis:** Sample size will be dependent on the number of online responders. We hope that there will be significant uptake from a number of countries globally. The recruitment will be online so no breach to social distancing rules will apply. To compare differences in binary variables we will use the likelihood ratio, chi-squared or Fisher's exact test and will consider a p value  $< 0.05$  as statistically significant. Ninety-five per cent confidence intervals for rates of psychological morbidity will be created using the Wilson method.

**Dissemination:** Following the collection and analysis of the data obtained from the questionnaire, we aim to present the findings to the ISUOG membership and publish in peer-reviewed journals – preferably Ultrasound in Obstetrics and Gynecology. The need to understand the experience of the pandemic by frontline workers is extremely important, not only to support them now, but to prepare for any future international healthcare emergency.

## References

1. Abramowicz JS, Basseal JM, Brezinka C, Dall'Asta A, Deng J, Harrison G, Marsal K, Lee JCS, Lim A, Miloro P, Poon LC, Salvesen KJ, Sande R, ter Haar G, Westerway SC, Xie MX, Lees C. ISUOG Safety Committee Position Statement on use of personal protective equipment and hazard mitigation in relation to SARS-CoV-2 for practitioners undertaking obstetric and gynecological ultrasound. *ISUOG*. 2020 March 28. Available from: <https://www.isuog.org/uploads/assets/b8dde768-08a2-424c-a4615551637515e9/ISUOG-Safety-Committee-statement-on-COVID19-and-PPE.pdf>
2. Neto MLR, Almeida HG, Esmeraldo JD, Nobre CB, Pinheiro WR, de Oliveira CRT, Sousa IDC, Lima OMML, Lima NNR, Moreira MM, Lima CKT, Junior JG, da Silva CGL.

- 1        When health professionals look death in the eye: the mental health of professionals
- 2        who deal daily with the 2019 coronavirus outbreak. *Psychiatry Res.* 2020 April
- 3        13;288:112972. doi:10.1016/j.psychres.2020.112972.
- 4        3. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta psychiatrica*
- 5        *scandinavica.* 1983 Jun;67(6):361-70.
- 6
- 7

1     **Supplement 3: Breakdown of healthcare professional (HCP) roles for analysis.**

Job title	
Doctor in research – Gynecology	Split up based on age: trainees/residents and consultants/attending
Doctor in research – Obstetrics	Split up based on age: trainees/residents and consultants/attending
Doctor in research – Obstetrics and Gynecology	Split up based on age: trainees/residents and consultants/attending
Doctor in research – Radiology/Sonologist	Split up based on age: trainees/residents and consultants/attending
Doctor in training – Gynecology	Split up based on age: trainees/residents and consultants/attending
Doctor in training – Obstetrics	Split up based on age: trainees/residents and consultants/attending
Doctor in training – Obstetrics and Gynecology	Split up based on age: trainees/residents and consultants/attending
Doctor in training – Radiologist/Sonologist	Split up based on age: trainees/residents and consultants/attending
Gynecologist	Split up based on age: trainees/residents and consultants/attending
Midwife – Early pregnancy only	Sonographer
Midwife – Obstetrics	Sonographer
Nurse – Early pregnancy only	Sonographer
Nurse – Gynecology	Sonographer
Nurse – Obstetrics	Sonographer
Nurse – Obstetrics and Gynecology	Sonographer
Obstetrician	Split up based on age: trainees/residents and consultants/attending
Obstetrician and Gynecologist	Split up based on age: trainees/residents and consultants/attending
Radiologist/Sonologist	Split up based on age: trainees/residents and consultants/attending
Retired	Exclude from analyses
Sonographer – Gynecology	Sonographer
Sonographer – Obstetrics	Sonographer
Sonographer – Obstetrics and Gynecology	Sonographer

2  
3


1 **Supplement 4: Breakdown of risk by area of work for analysis.**

Areas of work	Risk of COVID-19 infection
Early pregnancy unit - Ultrasound and assessment	High risk
Gynecology - Acute Gynecology	High risk
Gynecology - Benign Gynecology	Low risk
Gynecology – Oncology	Low risk
Gynecology - Ultrasound	High risk
Obstetrics/Gynecology/Radiology - From home	Low risk
Obstetrics - Antenatal clinic	Moderate risk
Obstetrics - Antenatal ward	Moderate risk
Obstetrics - Birth center/Birthing center	High risk
Obstetrics - Community clinics	Moderate risk
Obstetrics - Labor ward	High risk
Obstetrics - Maternity helplines	Low risk
Obstetrics - Maternity triage/Hospital obstetric triage	High risk
Obstetrics - Postnatal ward	Moderate risk
Obstetrics – Ultrasound	High risk
Radiology - non-Ultrasound	Moderate risk
Radiology - Ultrasound	High risk

2

3

1   **Supplement 5: Flowchart summarizing survey participants, differentiating International**  
2   **Society of Ultrasound in Obstetrics and Gynecology (ISUOG) members and non-ISUOG**  
3   **members (n = 3287).**



4  
5  
6   Response rate:  
7   ISUOG members - 23.1% (2441/10589)  
8   Non-ISUOG members - 3.2% (796/24920).  
9   Abbreviations: HADS (Hospital Anxiety and Depression scale)



1 **Supplement 6: Demographics of survey participation, differentiated based on**  
 2 **International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) membership (n**  
 3 **= 3237).**

	ISUOG members (n = 2441)	Others (n = 796)
<b>Total ISUOG mailing list (n)</b>	10589	24920
<b>Response of total ISUOG mailing list (n, %)</b>	2441 (23%)	796 (3%)
Countries (n)	120	78
Number of respondents in English (n)	1603	357
Number of respondents in Spanish (n)	564	317
Number of respondents in Italian (n)	195	72
Number of respondents in French (n)	79	50
<b>Completed survey</b>	<b>1795</b>	<b>442</b>
<b>Countries (n)</b>	110	65
<b>Age (mean, range)</b>	47.2 (18-82)	47.7 (25-78)
≤20 years	1 (0%)	0 (0%)
21-30 years	95 (5%)	22 (5%)
31-40 years	454 (25%)	104 (24%)
41-50 years	549 (31%)	137 (31%)
51-60 years	475 (26%)	121 (28%)
61-70 years	201 (11%)	50 (11%)
>70 years	18 (1%)	6 (1%)
<b>Gender (n, %)</b>		
Female	1176 (66%)	298 (68%)
Male	613 (34%)	142 (32%)
Intersex	1 (0%)	0 (0%)
Prefer not to say	3 (0%)	0 (0%)
<b>Prevalence suspected or confirmed SARS-CoV-2 based on symptoms +/- PCR testing (%)</b>	230 (13%)	60 (14%)
<b>Is PPE used by respondents (n, %)</b>		
Not yet	217 (12%)	62 (14%)
Yes - for aerosol generating procedures only	110 (6%)	23 (5%)
Yes - for all patients	1142 (64%)	250 (57%)
Yes - for suspected/positive patients only	324 (18%)	105 (24%)
<b>What PPE is used by respondents (n, %)</b>		
None	59 (3%)	17 (4%)
Gloves only	38 (2%)	11 (3%)
Gloves + surgical mask	393 (22%)	104 (24%)
Gloves + surgical mask + visor	116 (6%)	36 (8%)
Gloves + FFP3 mask	83 (5%)	12 (3%)
Gloves + FFP3 mask + visor	39 (2%)	12 (3%)
Gloves + FFP3 mask + visor + gown + surgical hat	70 (4%)	20 (5%)

<i>Patient wearing mask</i>	1482 (83%)	359 (82%)
<b>Concerns regarding a lack of PPE (n, %)</b>		
<i>No</i>	475 (26%)	104 (24%)
<i>Gloves</i>	223 (12%)	55 (13%)
<i>Gown</i>	499 (28%)	130 (30%)
<i>N95/FFP3 mask</i>	1014 (57%)	265 (60%)
<i>Surgical hat</i>	187 (10%)	46 (10%)
<i>Surgical mask</i>	657 (37%)	163 (37%)
<i>Visor</i>	458 (26%)	124 (28%)
<i>Surgical mask (patient)</i>	479 (27%)	130 (30%)
<b>Type of PPE that has been re-used or recycled (n, %)</b>		
<i>No</i>	690 (38%)	175 (40%)
<i>Gloves</i>	27 (2%)	7 (2%)
<i>Gown</i>	229 (13%)	58 (13%)
<i>N95/FFP3 mask</i>	722 (40%)	164 (37%)
<i>Surgical hat</i>	74 (4%)	18 (4%)
<i>Surgical mask</i>	435 (24%)	130 (30%)
<i>Visor</i>	493 (27%)	128 (29%)
<b>Anxiety (mean, 95% CI)*</b>	7 (6.7-7.3)	6.6 (6.1-7.1)
<i>0-7 (none)</i>	480 (57%)	140 (63%)
<i>8-10 (mild)</i>	180 (22%)	49 (22%)
<i>11-21 (moderate to severe)</i>	175 (21%)	34 (15%)
<b>Depression (mean, 95% CI)*</b>	5.4 (5.2-5.7)	4.9 (4.4-5.4)
<i>0-7 (none)</i>	587 (70%)	173 (78%)
<i>8-10 (mild)</i>	171 (20%)	34 (15%)
<i>11-21 (moderate to severe)</i>	77 (9%)	16 (7%)

\*Numbers of HADS respondents lower than for main survey (n = 835 ISUOG members, n = 223 Others)

Abbreviations: PCR (Polymerase Chain Reaction); PPE (Personal Protective Equipment)

1  
2  
3

1     **Supplement 7: Comparison of comorbidity in those with (n = 290) and without suspected**  
2     **or confirmed SARS-CoV-2 (n = 1947) (Total n = 2237).**

	Suspected or confirmed SARS-CoV-2	Without suspected or confirmed SARS-CoV-2
Heart disease (n, %)	3 (1%)	43 (2%)
Lung disease (n %)	30 (10%)	160 (8%)
Use of antihypertensives (n, %)	39 (13%)	283 (15%)
Smokers (n, %)	13 (4%)	125 (6%)

3  
4

1     **Supplement 8: Comparison of Hospital Anxiety and Depression scale (HADS) scores in**  
2     **those with (n = 162) and without suspected or confirmed SARS-CoV-2 (n = 896) (Total n =**  
3     **1058).**

	Suspected or confirmed SARS-CoV-2	Without suspected or confirmed SARS-CoV-2
Anxiety (mean, 95% CI)	7.4 (6.7-8.0)	6.8 (6.6-7.1)
0-7 (none)	86 (53%)	534 (60%)
8-10 (mild)	32 (20%)	197 (22%)
11-21 (moderate to severe)	44 (27%)	165 (18%)
Depression (mean, 95% CI)	5.8 (5.2-6.4)	5.2 (5.0-5.5)
0-7 (none)	105 (65%)	655 (73%)
8-10 (mild)	37 (23%)	168 (19%)
11-21 (moderate to severe)	20 (12%)	73 (8%)

1     **Supplement 9: Hospital Anxiety and Depression scale (HADS) breakdown by gender (n =**  
2     **1058).**

	Female	Male	Prefer not to say
<b>Number of respondents (n, %)</b>	754 (71%)	303 (29%)	1 (0%)
<b>Anxiety (mean, 95% CI)*</b>	6.9 (6.6-7.2)	5.7 (5.3-6.2)	-
0-7 (None) (n, %)	450 (60%)	202 (67%)	1 (100%)
8-10 (Mild) (n, %)	148 (20%)	60 (20%)	0 (0%)
11-21 (Moderate to severe) (n, %)	156 (21%)	41 (14%)	0 (0%)
<b>Depression (mean, 95% CI)*</b>	5.4 (5.2-5.7)	5.0 (4.6-5.4)	-
0-7 (none) (n, %)	529 (70%)	230 (76%)	1 (100%)
8-10 (mild) (n, %)	146 (19%)	59 (19%)	0 (0%)
11-21 (moderate to severe) (n, %)	79 (10%)	14 (5%)	0 (0%)

1     **Supplement 10: Hospital Anxiety and Depression scale (HADS) breakdown by risk of**  
2     **working area (n = 1058).**

	Low risk	Moderate risk	High risk
<b>Number of respondents (n, %)</b>	23 (2%)	28 (3%)	1007 (95%)
<b>Anxiety (mean, 95% CI)*</b>	8.3 (6.3-10.3)	5.1 (3.6-6.6)	6.6 (6.3-6.8)
0-7 (None) (n, %)	13 (57%)	21 (75%)	619 (61%)
8-10 (Mild) (n, %)	3 (13%)	3 (11%)	202 (20%)
11-21 (Moderate to severe) (n, %)	7 (30%)	4 (14%)	186 (18%)
<b>Depression (mean, 95% CI)*</b>	8.2 (6.0-10.3)	4.3 (2.7-5.8)	5.3 (5.0-5.5)
0-7 (none) (n, %)	12 (52%)	19 (68%)	729 (72%)
8-10 (mild) (n, %)	4 (17%)	6 (21%)	195 (19%)
11-21 (moderate to severe) (n, %)	7 (30%)	3 (11%)	83 (8%)

3



1     **Supplement 11: Comparison of Hospital Anxiety and Depression scale (HADS) scores in**  
2     **those with (n = 788) and without personal protective equipment (PPE) shortages (n = 270)**  
3     **(Total n = 1058).**

	PPE shortages	No PPE shortages
Anxiety (mean, 95% CI)	7.4 (7.1-7.7)	5.5 (5.0-6.0)
0-7 (none)	430 (55%)	190 (70%)
8-10 (mild)	183 (23%)	46 (17%)
11-21 (moderate to severe)	175 (22%)	34 (13%)
Depression (mean, 95% CI)	5.7 (5.5-6.0)	4.1 (3.7-4.5)
0-7 (none)	540 (69%)	220 (81%)
8-10 (mild)	167 (21%)	38 (14%)
11-21 (moderate to severe)	81 (10%)	12 (4%)

4