



# BMJ Open Practice variation across five European paediatric emergency departments: a prospective observational study

Fabienne Ropers <sup>1</sup>, Patrick Bossuyt,<sup>2</sup> Ian Maconochie,<sup>3</sup> Frank J Smit,<sup>4</sup> Claudio Alves,<sup>5</sup> Susanne Greber-Platzer,<sup>6</sup> Henriette A moll <sup>7</sup>, Joany Zachariasse<sup>7</sup>

**To cite:** Ropers F, Bossuyt P, Maconochie I, *et al.* Practice variation across five European paediatric emergency departments: a prospective observational study. *BMJ Open* 2022;**12**:e053382. doi:10.1136/bmjopen-2021-053382

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-053382>).

Received 20 May 2021

Accepted 12 February 2022

## ABSTRACT

**Objectives** To compare paediatric healthcare practice variation among five European emergency departments (EDs) by analysing variability in decisions about diagnostic testing, treatment and admission.

**Design and population** Consecutive paediatric visits in five European EDs in four countries (Austria, Netherlands, Portugal, UK) were prospectively collected during a study period of 9–36 months (2012–2015).

**Primary outcome measures** Practice variation was studied for the following management measures: lab testing, imaging, administration of intravenous medication and patient disposition after assessment at the ED.

**Analysis** Multivariable logistic regression was used to adjust for general patient characteristics and markers of disease severity. To assess whether ED was significantly associated with management, the goodness-of-fit of regression models based on all variables with and without ED as explanatory variable was compared. Management measures were analysed across different categories of presenting complaints.

**Results** Data from 111 922 children were included, with a median age of 4 years (IQR 1.7–9.4). There were large differences in frequencies of Manchester Triage System (MTS) urgency and selected MTS presentational flow charts. ED was a significant covariate for management measures. The variability in management among EDs was fairly consistent across different presenting complaints after adjustment for confounders. Adjusted OR (aOR) for laboratory testing were consistently higher in one hospital while aOR for imaging were consistently higher in another hospital. Iv administration of medication and fluids and admission was significantly more likely in two other hospitals, compared with others, for most presenting complaints.

**Conclusions** Distinctive hospital-specific patterns in variability of management could be observed in these five paediatric EDs, which were consistent across different groups of clinical presentations. This could indicate fundamental differences in paediatric healthcare practice, influenced by differences in factors such as organisation of primary care, diagnostic facilities and available beds, professional culture and patient expectations.

## INTRODUCTION

Variability in healthcare delivery can indicate appropriate use, overuse and underuse

## Strengths and limitations of this study

- Large European study on paediatric practice variation in emergency departments (EDs) including the entire range of paediatric presentations.
- Information on presenting complaint available.
- Adjustment for important patient characteristics and markers of disease severity.
- No data on differential diagnosis after assessment by ED physician or outcome.
- No specific data on referral status available.

of resources. Differences in patient characteristics, including severity and nature of presenting problems, result in differences in diagnostic and therapeutic management.<sup>1</sup> This resulting variation in management is warranted, because different clinical problems require different management to achieve the best patient outcome.<sup>2–4</sup>

Yet variation can also arise from other factors, like differences in practice guidelines and adherence, medical tradition, patient expectations or healthcare organisation.<sup>5–9</sup> In these instances, both deviations in management to the lower and higher end of the spectrum and higher and lower resource use can be associated with poorer outcomes or lower cost efficiency, depending on the underlying factors. Studying practice variation has therefore been acknowledged as an important tool to identify areas with potential for improvement of patient care.

Several studies have observed practice variation in the paediatric emergency setting, for specific presentations,<sup>10 11</sup> such as minor head injury or respiratory symptoms. Other studies have focused on variability in resource use in paediatric emergency departments (EDs) in low acuity presentations. These studies reported that physician training background was associated with resource use and that diagnostic testing and procedures were less frequent in the low acuity group.<sup>12 13</sup> Many



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Ms Joany Zachariasse;  
j.zachariasse@erasmusmc.nl

studies have been conducted in the North American setting and not all were able to adjust for differences in patient characteristics, such as disease severity.<sup>14</sup> Large scale European studies are scarce.

This large multicentre study aimed to compare paediatric healthcare practice among five European EDs. We wanted to analyse variability in decisions about diagnostic testing, treatment and admission, after adjustment for patient characteristics, across subgroups of presenting problems covering the broad spectrum of paediatric ED presentations.

## METHOD

### Study design, data source and study population

This study was embedded in the TRIAGE project (Triage Improvement Across General Emergency departments for paediatric patients), a prospective observational study and followed from observations in previous analyses. The study design has been described in detail elsewhere.<sup>15</sup> In brief, during this project electronic health record data of all ED visits of children <16 years were prospectively collected in five different hospitals in four different countries. The five participating hospitals were: Erasmus Medical Centre, the Netherlands; Maastad Hospital, the Netherlands; St. Mary's hospital Imperial College Healthcare National Health Service Trust, UK; Hospital Prof. Dr. Fernando Fonseca, Portugal; Vienna general hospital, Austria. In the latter ED, only low urgent trauma cases presented, because the majority of patients with trauma were seen in the traumatology department.

Study sites were diverse in their catchment area and complexity of the patient population, number of visits and organisation of healthcare. Data were obtained by questionnaires obtained from the participating EDs (online supplemental appendix 1). Four EDs were paediatric EDs, and one was mixed adult-paediatric. The supervising physician was a paediatrician in all EDs, and in one site a paediatric emergency physician. The enrolment period varied from 8 to 36 months between 2012 and 2015, during which 119 209 consecutive ED visits were included. The differences in patient load account for differences in enrollment time to include sufficient patients. Also practical reasons, such as availability of staff to help in high quality data collection, played a role.

Nurses at the participating EDs were informed about the study and encouraged to be complete in their registration of routine medical data.<sup>15</sup>

Children with incomplete triage data were excluded from the analysis. Complex comorbidity has been linked to a higher use of diagnostics and therapeutic interventions at the ED.<sup>16</sup> Children with known complex comorbidity were therefore excluded if patient-level information was available. This was the case for hospitals with high proportions of comorbidity: Erasmus MC, St Mary's and Vienna general hospital (10%–38% comorbidity). Maastad Hospital and Hospital Fernando Fonseca reported an estimated total comorbidity of less than 10%, and

much lower proportions of complex comorbidity, and did not provide patient-level information. Comorbidity was defined according to the Paediatric Medical Complexity Algorithm.<sup>17 18</sup>

### Main outcome measures

We evaluated ordering of diagnostic tests (laboratory testing and imaging at the ED), administration of intravenous medication or fluids and hospital admission. Laboratory testing included tests and cultures in blood, urine, faeces, and cerebrospinal fluid. Imaging included X-ray, ultrasound, CT and MRI. Admission was defined as admission from the ED to the general ward or paediatric intensive care unit.

### Confounders

Patient characteristics (age, gender), physiological parameters (heart rate, respiratory rate, oxygen saturation, temperature), presentational flow chart and urgency according to the Manchester triage system (MTS), and presentation during office hours or during out-of-office hours were considered as potential confounding variables. Office hours were defined as Monday until Friday, between 08:00 am and 05:59 pm, and all other time points were defined as out-of-office hours. Vital signs and age were included as continuous variables.

In all participating hospitals, the MTS was routinely used for triage of presenting children. The MTS consists of 53 presentational flow charts that cover almost all presentations to EDs.<sup>19</sup>

The triage nurses are trained to select the most specific presentational flow chart. Only if there is no defining symptom at presentation the nurse will select an aspecific flow chart, like unwell child or crying baby. To ensure sufficient standardisation of triage, triage nurses using the MTS are well trained.

Presentational flow charts in turn consist of signs and symptoms that classify patients into five urgency categories, indicating the time to first contact with the treating clinician. These categories were assigned to three groups: MTS emergent or very urgent (<10 min waiting time), MTS urgent (<60 min waiting time), and MTS standard (60–120 min) or non-urgent (120–240 min waiting time).

To create subgroups of comparable presenting symptoms, we used MTS presentational flow charts. These were grouped into nine categories as defined in our previous publications: cardiac, dermatologic, ear/nose/throat, gastrointestinal, neurologic/psychiatric/intoxications, respiratory, trauma/muscular, unwell and urinary/gynaecological.<sup>15 20</sup> Heterogeneous presentations with low frequency were grouped together as 'other' (online supplemental appendix 2).

In addition to the subgroups of presenting symptoms based on MTS presentational flow charts, we defined a subgroup of infectious presentations, because a suspected infection is an important reason for presentation at the ED. We defined this subgroup as children <5 years old, who had been assigned to the presentational flow

chart shortness of breath or vomiting/diarrhoea or had presented with fever (defined as temperature  $\geq 38.5^{\circ}\text{C}$  on presentation or MTS discriminator hot child).

### Statistical analysis

We evaluated ordering of diagnostic tests, initiation of treatment and hospital admission across centres, using multivariable logistic regression models, adjusting for identified confounders. In this analysis, the Maasstad hospital was (randomly) selected as the reference. Differences between EDs are expressed as adjusted ORs (aORs), relative to practice in the Maasstad hospital, with 95% CIs.

Patient characteristics and all other included variables are presented using descriptive statistics with absolute numbers, proportions, ranges and medians as appropriate. Vital signs are presented as proportion abnormal, based on the Advanced Paediatric Life Support reference values, with fever defined as a temperature  $\geq 38.5^{\circ}\text{C}$ .<sup>21</sup>

To assess whether ED was significantly associated with management when adjusted for confounding factors, the fit of regression models based on all variables with and without ED as explanatory variable was compared using the generalised likelihood ratio test statistic. Patients were then stratified according to categories of MTS presentational flow charts and separate regression analyses were performed within those strata. Because the ED of General hospital Vienna only treated a small proportion of trauma patients, this hospital was excluded from the analysis in the category trauma/muscular. Results of the presentational flow chart category 'other' are not presented, because of the inherent heterogeneity of this category.

Missing data for vital signs were imputed 25 times using the MICE algorithm in R (V.3.6.3). These missing data were assumed to be missing at random, conditional on other variables in the database. The imputation model included all predictors and outcome measures and additional descriptors of case mix: patient age and sex, date and time of arrival, and triage characteristics.<sup>15 22</sup> Analyses were performed with IBM SPSS statistics, V.25 (IBM).

### Patient and public involvement

Patients or the public were not involved in the design, or conduct, or reporting, or dissemination plans of our research.

## RESULTS

### Study Group

Of all 119 209 ED visits of patients 16 years or younger included in the TriAGE cohort, 5706 were excluded because of complex comorbidity, leaving 113 503 who met the inclusion criteria. A total of 1581 presentations had to be excluded because of missing presentational flow chart ( $n=1578$  presentations) or missing time of arrival ( $n=3$  presentations), resulting in a study group of 111 922 presentations (94%).

Across the 5 EDs, the median age at presentation ranged from 3.8 to 5.7 years, and 42%–48% of children were female (table 1). Most children presented with general malaise or because of parental concern, trauma or injuries, gastrointestinal or respiratory complaints. Between 11% and 33% of children had tachypnoea at presentation, 11%–18% tachycardia and 4%–9% had a recorded temperature of  $\geq 38.5^{\circ}\text{C}$ . In concordance with differences in frequency of abnormal vital signs, the casemix of patients differed among EDs with respect to MTS urgency and presentational complaint. In Erasmus and Maasstad hospital, for example, 46%–47% of patients were triaged as urgent, compared with 18%–24% of patients presenting at the three other hospitals (table 1).

### Management differences across EDs

Management also varied among EDs, with Vienna performing lab tests in 36% of presentations against 9.2% in St Mary's. Likewise, imaging was performed in 24%–37% of presentations in Maasstad, Erasmus and Fernando, while in only 7.2% of patients presenting in Vienna. Differences in therapy were less pronounced but, with regard to admission, high admission rates (20%–23%) were observed in Erasmus and Maasstad, while only 4.6%–9.6% of patients were admitted in the other hospitals (table 2). Inclusion of ED as confounding variable in the multivariable regression model improved model fit for all management measures ( $p<0.001$ ), indicating that management differed depending on the ED of presentation.

### Management differences within presentational flow chart categories

Because management will be guided by presenting complaint, we assessed differences in management across EDs in children with comparable presenting complaints. The size of presentational flow categories relative to total presentations varied per hospital. The MTS urgency within categories also differed, with higher MTS urgency in Maasstad and Erasmus, indicating differences in patient populations between EDs (table 1, figure 1).

In most presentational flow chart categories we observed, after adjusting for patient characteristics, time of presentation and markers disease severity, that patients presenting in Vienna and, for some categories, Erasmus MC, were more likely to receive lab testing. Patients presenting in Fernando were more likely to receive imaging in the majority of categories, followed by Maasstad and Erasmus MC (figure 1). Intravenous administration of medication or fluids was more likely in Maasstad hospital and, in some categories, in Erasmus MC and Fernando, compared with other hospitals. Admission was more likely in Maasstad hospital, followed by Erasmus MC. The chance of admission was consistently lower elsewhere after adjustment for other parameters, with the exception of smaller categories with broader CIs. One ED had an overall average or lower likelihood of medical interventions (St Mary's), but for other EDs, instead of

	Emergency department					
	Maasstad	Erasmus	Fernando	St Marys	Wien	Total
N	10 484	13 968	53 175	15 027	19 268	111 922
Patient characteristics						
Age in yrs (median, IQR)	5.7 (1.9–11.6)	4.1 (1.3–9.8)	4.7 (2.0–9.5)	3.8 (1.5–8.7)	3.9 (1.6–8.3)	4.4 (1.7–9.4)
Gender, n % female	43.3	42.2	47.9	44.2	47.5	46.2
Abnormal vital signs (95 th percentile APLS 2017)*						
Tachypnea (%)	32.9	20.3	10.8	16.9	22.3	16.9
Bradypnea (%)	1.9	5.2	7.5	1.3	4	5.3
Tachycardia (%)	18.2	12.3	12.9	14.1	10.8	13.1
Bradycardia (%)	4.4	7.9	6	4.3	10.3	6.6
Oxygen saturation <94% (%)	1.9	1.8	1.5	1.4	1	1.5
Fever (temp ≥38.5°C (%))	8	9.3	4	6.4	6.6	5.8
No of abnormal vital signs (%)						
0	53.9	61.7	67	69.9	59.4	64.2
1	33.8	30.1	27.8	23.1	33.3	28.9
2	11.6	7.7	4.9	6.4	7	6.4
3	0.7	0.6	0.3	0.6	0.3	0.4
MTS urgency (%)						
Emergent  very urgent	15.7	14	11.9	10.6	5.4	11.2
Urgent	47.4	45.7	20.4	24.3	18.1	26.2
Standard  non-urgent	36.8	40.3	67.7	65.1	76.5	62.5
Time of presentation (%)						
Office hours	39.8	47.3	42.3	36	43.6	42.1
Out of office hours	60.2	52.7	57.7	64	56.4	57.9
Presentational flow chart categories						
Cardiac	0.4	1	1.2	0.8	1.8	1.2
Dermatologic	8.5	11.8	14.3	9.9	14	12.8
ENT	1.6	3	14	4.4	14	10.2
Gastrointestinal	10	12.7	16.2	11.5	21.1	15.4
Neurologic/psychiatric	2.4	7.5	3.1	2.8	4	3.7
Respiratory	12.1	8.1	11.2	11.2	16.6	11.8

Continued



Table 1 Continued

## Emergency department

	Maasstad	Erasmus	Fernando	St Marys	Wien	Total
Trauma/muscular	44.3	29.9	14.7	23.2	3.3	18.6
Unwell	16.2	20.3	19	30.9	17.1	20.1
Urinary/gynaecological	1.2	2.8	2.3	1.5	2.3	2.1
Other	3.4	2.9	3.9	3.9	6	4.1
<b>Missing values</b>	<b>Maasstad</b>	<b>Erasmus</b>	<b>Fernando</b>	<b>St Marys</b>	<b>Wien</b>	<b>Total</b>
Heart rate	60.9% (n=6380)	51.1% (n=7138)	35.9% (n=19106)	19.6% (n=2940)	61.4% (n=11830)	42.3% (n=47394)
Respiratory rate	83.1% (n=8712)	68.2% (n=9531)	35.9% (n=19106)	23.6% (n=3544)	86.8% (n=16715)	51.5% (n= 57608)
Oxygen saturation	61.2% (N=6418)	69.4% (n=9694)	34.4% (n=18279)	19.8% (n=2973)	61.2% (n=11799)	43.9% (n=49163)
Temperature	57.9% (n=6069)	47.4% (n=6626)	12.1 % (n=6431)	32.4% (n=4872)	1% (n=194)	21.6% (n= 24192)

\*Presented as percentage of measured values. Percentage of missing values of vital signs is displayed above. ENT, ears, nose and throat; MTS, Manchester Triage System.

overall high or low resource use, there were specific interventions that were performed more or less likely within EDs (figures 1 and 2). The likelihood of administration of intravenous medication and admission seemed to vary in parallel directions.

### Subanalysis in infectious children

An additional regression analysis was performed in the subgroup of young children with suspected infectious diseases. Similar patterns of variability in management across EDs were observed (table 3). Lab testing was more likely in Vienna and in Erasmus MC, imaging more likely in Fernando, intravenous medication and admission more likely in Maasstad hospital, followed by Erasmus MC. This means that, in this more homogeneous group of children, there was no apparent lower variability in management among different EDs.

## DISCUSSION

In this large observational study of paediatric practice variation across five European EDs, management was associated with ED of presentation. We observed ED-related patterns of variability in the likelihood of diagnostic testing, intravenous medication and admission, which remained stable across groups of clinical presentations, after correcting for several general patient characteristics and markers of disease severity known to be associated with management. Though one ED had overall low resource use, there were large differences across other EDs in likelihood for imaging or laboratory testing, after adjusting for the differences in disease severity and presenting symptoms that were observed between hospitals.

Other unmeasured medical and non-medical factors are likely to play a role in hospital-specific patterns of variability. The proportion of self-referred patients differed greatly among hospitals (online supplemental appendix 1). Reasons for primary care physicians to refer to an ED include available diagnostic facilities, request for a professional opinion, or expected need for in-hospital treatment.<sup>6</sup> This means that disease characteristics of referred and non-referred presentations are likely to differ. These factors could partly be adjusted for by the measures of disease severity and presenting symptoms.

Prior out-of-hospital diagnostics and treatment will also influence management at the ED. The higher rate of referrals by primary care physicians in Maasstad hospital and Erasmus MC could account for the higher likelihood of admission to these hospitals, as has been reported previously.<sup>7 23</sup> Parent and patient expectations regarding management differ between self-referred and referred patients. Presentation at ED without prior consultation of the primary care physician can be triggered by parental perceptions of disease severity and the expectation that specific diagnostic facilities or treatment available at the ED are required.<sup>6 24-27</sup> This can also stimulate health-care providers to perform additional testing or influence their treatment decisions.<sup>28</sup> However, referral status

**Table 2** Management per ED

	Maasstad	Erasmus	Fernando	St Marys	Wien	Total
N	10 484	13 968	53 175	15 027	19 268	111 922
Diagnostic						
Lab any (%)	20	28.5	13.1	9.2	35.8	19.1
Imaging any (%)	37.2	24.9	23.7	14.2	7.2	21
Therapy						
Intravenous medication or fluids (%)	12.8	9.5	7.5	4.1	4	7.2
Admission						
General admission/ICU admission (%)	23.4	20.3	5.2	9.6	4.6	9.3
ICU admission (% of total)	0.2	2.2	0.3	0.1	0	0.4


ED, emergency department; ICU, intensive care unit.

only cannot explain the variability in management that was observed in the three hospitals with comparably low referral rates.


A myriad of other factors has been linked to clinical management. Financial incentives embedded in the organisation of healthcare systems could differ across EDs. National or local professional culture, standard of

care and facilities might partly account for the observed variability, such as preferences for lab testing, imaging, and the availability thereof.<sup>29–31</sup> Differences in practice guidelines, reflecting these differences in professional culture and diagnostic options, could also be of influence. These are neither harmonised across European countries, nor is adherence likely to be comparable


## Laboratory tests




## Imaging



## Iv medication or fluids



## Admission



OR are adjusted for age, gender, MTS urgency category, heart rate, respiratory rate, oxygen saturation, temperature and time of presentation

**Figure 1** aOR for management according to presentational flow chart categories. aOR, adjusted OR.



**Figure 2** Radar charts presenting aOR for management outcome measures in the five largest presentational flow chart categories. aOR, Adjusted OR.

across EDs. Holding varying guideline recommendations regarding lab tests and imaging partly responsible for the observed patterns would reflect international differences in the general value placed on specific diagnostic tests, regardless of disease presentation, as the differences in additional testing were rather consistent and independent of presenting complaint.

Parent and patient expectations and preferences regarding healthcare are affected by cultural and socio-economic factors. These, in turn, influence management

decisions and could represent another non-medical factor contributing to the observed variability.<sup>28 32</sup> Professional education and training have been reported to be associated with management, where paediatric specialty training was linked to a lower amount of diagnostic testing.<sup>5 33 34</sup> However, in our study there was no difference in respect to those factors among hospitals with higher and lower likelihood of testing.

**Table 3** aOR for infectious children <5 years

aOR for management in infectious children (n=23695)	Any lab tests		Any imaging		Intravenous medication or fluids		Admission (ICU and general)	
	aOR	95% CI	aOR	95% CI	aOR	95% CI	aOR	95% CI
Maasstad	Reference		Reference		Reference		Reference	
Erasmus	2.64	(2.33 to 2.99)	3.66	(2.93 to 4.56)	0.63	(0.53 to 0.73)	0.75	(0.65 to 0.85)
Fernando	0.89	(0.79 to 1.00)	6.91	(5.63 to 8.48)	0.43	(0.38 to 0.50)	0.19	(0.17 to 0.22)
St Marys	0.36	(0.30 to 0.41)	1.99	(1.57 to 2.52)	0.25	(0.20 to 0.30)	0.33	(0.29 to 0.38)
Wien	2.88	(2.54 to 3.27)	2.85	(2.28 to 3.57)	0.36	(0.30 to 0.43)	0.17	(0.15 to 0.20)

Based on MTS flow chart 'diarrhoea and vomiting' or 'shortness of breath', or based on presence of fever (MTS discriminator hot child/adult or temp  $\geq 38^{\circ}\text{C}$ ). ORs are adjusted for age, gender, MTS urgency category, heart rate, respiratory rate, oxygen saturation, temperature and time of presentation.

aOR, adjusted OR; ICU, intensive care unit; MTS, Manchester Triage System.

## Strengths/limitations

A major strength of this study is that we could adjust for several relevant patient characteristics and markers of disease severity, due to the availability of triage urgency data, presentational flow chart, vital signs and basic patient characteristics. We could include a large sample of patients from different European countries. This is an advantage, because these differences can help in identifying relevant factors responsible for practice variation, but also represents a limitation, since individual effects could not be disentangled. Hospitals differed in multiple characteristics, such as the availability of primary care physicians, rate of self-referrals, and patient casemix. Patient-specific data on referral were not available for all hospitals, and referral status could therefore not be included in the regression analyses. In addition, the availability of resources, including staffing and beds, could vary during the project, but exact data were missing for our analysis.

We used the selected MTS presentational flow chart as a proxy for presenting symptoms. In the course of the evaluation at the ED, the initial impression will have changed in a proportion of children, due to the elucidation of other signs and symptoms, which could lead to adjustments to the differential diagnosis and changes in subsequent management steps. Because we had no data on differential diagnosis and final diagnosis, we could only stratify according to presenting symptoms. The remaining heterogeneity of patients within categories and between EDs will have contributed to the observed variability in management. We did not have patient outcome measures available, therefore the consequences of deviations, compared with the benchmark, could not be assessed in terms of effects on outcomes.

## Implications

Our analysis revealed substantial variability in management, even after adjustment for relevant patient characteristics and markers of disease severity. We acknowledge that not all practice variation is unwarranted or problematic, because contextual and patient-related factors such as those described above can cause variation that is not associated with lower quality care.<sup>35</sup>

However, we believe that our findings of consistently higher likelihood of lab testing or imaging in some hospitals, compared with others, are sufficient reason to further study underlying reasons for these patterns. In that sense, ours can serve as a pilot study. As a starting point, deviations from the benchmark should prompt a general exploration of potential explanations, and how these deviations might affect patient outcome. In a second step, a review of recent guidelines and review syntheses, combined with an assessment of adherence to guidelines, could provide further insights. An accessible and feasible approach could be to increase awareness of practice guidelines during handover and rounds on a case level. Both by following recommendations with a strong evidence base for a well-defined population in favour of providing healthcare actions, and by following recommendations against certain practices

because of insufficient added value, quality of care will be improved and variation will be reduced.

A related study focusing on febrile children found that admission varied across European EDs, after adjusting for explanatory variables comparable to the ones in our study but also for management at the ED, pointing to other factors than disease characteristics.<sup>36</sup> Factors related to organisation of healthcare and local culture of care will likely play an important role. Though more difficult to influence, comparing and learning from differences in organisation and medical culture can be a first step to long-term changes, to ensure an sustainable healthcare system. The number of EDs required for a study searching to assess the importance of these factors depends on the heterogeneity of the EDs and healthcare systems, and on the research question. Such evaluation should preferentially involve patient important outcomes and prior out-of-hospital management, to assess the entire trajectory of care and to produce suggestions for improvements.

## CONCLUSION

In this analysis of paediatric healthcare practice among five European ED distinctive hospital-specific patterns in variability of management could be observed, which were consistent over different groups of clinical presentations. This pattern in variability could indicate fundamental differences in paediatric healthcare practice across countries, influenced by factors such as organisation of primary care, diagnostic facilities and available beds, professional culture and patient expectations.

## Author affiliations

<sup>1</sup>Willem-Alexander Children's Hospital, department of Pediatrics, Leiden University Medical Center, Leiden, The Netherlands

<sup>2</sup>Department of Clinical Epidemiology, Biostatistics and Bioinformatics, Amsterdam University Medical Centers, Amsterdam, Netherlands

<sup>3</sup>Department of Paediatric Emergency Medicine, Imperial College Healthcare NHS Trust, London, UK

<sup>4</sup>Department of Pediatrics, Maasstad Hospital, Rotterdam, Netherlands

<sup>5</sup>General Pediatrics, Emergency Unit, Hospital Professor Doutor Fernando Fonseca EPE, Amadora, Portugal

<sup>6</sup>Department of Pediatrics and Adolescent Medicine, Medical University of Vienna, Wien, Austria

<sup>7</sup>Department of General Paediatrics, Erasmus Medical Center, Rotterdam, The Netherlands

**Contributors** IM, FJS, CA, SG-P, HAM and JZ substantially contributed to the conception and design of the TriAGE study and data acquisition. FR, HAM, PB and JZ conceived the study idea. FR performed the analysis and R, PB, HAM and JZ interpreted the results. FR wrote the first draft of the manuscript. All authors revised it critically for important intellectual content and gave their approval of the final version. All authors had full access to all the data (including statistical reports and tables) in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis. JZ is guarantor.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not applicable.



**Ethics approval** The study was approved by the participating institutions' medical ethical committees: Medical Ethics Committee Erasmus MC (MEC-2013-567), Maasstad Ziekenhuis Board of Directors (Protocol L2013-103), Imperial College London Joint Research Compliance Office (Reference number: 14SM2164; Ethics reference number 14/WA/1051), Comissão de Ética para a Saúde do Hospital Prof. Dr. Fernando Fonseca EPE (Reunião de 06 de Dezembro de 2017), Ethik Kommission Medizinische der Medizinischen Universität Wien (EK Nr: 1405/2014). All waived the requirement for informed consent.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** No data are available.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Fabienne Ropers <http://orcid.org/0000-0001-7807-4753>

Henriette A moll <http://orcid.org/0000-0001-9304-3322>

#### REFERENCES

- Jain S, Elon LK, Johnson BA, *et al*. Physician practice variation in the pediatric emergency department and its impact on resource use and quality of care. *Pediatr Emerg Care* 2010;26:902–8.
- Wennberg JE. Unwarranted variations in healthcare delivery: implications for academic medical centres. *BMJ* 2002;325:961–4.
- Mercuri M, Gafni A. Medical practice variations: what the literature tells us (or does not) about what are warranted and unwarranted variations. *J Eval Clin Pract* 2011;17:671–7.
- Goodman DC. Unwarranted variation in pediatric medical care. *Pediatr Clin North Am* 2009;56:745–55.
- Weiner SG, Ruffing RP, Barnewolt BA. A comparison of resource utilization between emergency physicians and pediatric emergency physicians. *Pediatr Emerg Care* 2012;28:869–72.
- Cheek JA, Braitberg G, Craig S, *et al*. Why do children present to emergency departments? exploring motivators and measures of presentation appropriateness for children presenting to a paediatric emergency department. *J Paediatr Child Health* 2017;53:451–7.
- Emery DP, Milne T, Gilchrist CA, *et al*. The impact of primary care on emergency department presentation and hospital admission with pneumonia: a case-control study of preschool-aged children. *NPJ Prim Care Respir Med* 2015;25:14113.
- Cook DA, Pencille LJ, Dupras DM, *et al*. Practice variation and practice guidelines: attitudes of generalist and specialist physicians, nurse practitioners, and physician assistants. *PLoS One* 2018;13:e0191943.
- Rogers AJ, Kuppermann N, Anders J, *et al*. Practice Variation in the Evaluation and Disposition of Febrile Infants  $\leq 60$  Days of Age. *J Emerg Med* 2019;56:583–91.
- Neuman MI, Graham D, Bachur R. Variation in the use of chest radiography for pneumonia in pediatric emergency departments. *Pediatr Emerg Care* 2011;27:606–10.
- Stanley RM, Hoyle JD, Dayan PS, *et al*. Emergency department practice variation in computed tomography use for children with minor blunt head trauma. *J Pediatr* 2014;165:1201–6.
- Samuels-Kalow ME, Niedzwiecki M, Friedman AB, *et al*. Comparing resource use between paediatric emergency department visits by triage level. *Emerg Med J* 2018;35:681–4.
- Doctor K, Breslin K, Chamberlain JM, *et al*. Practice pattern variation in test ordering for low-acuity pediatric emergency department patients. *Pediatr Emerg Care* 2018.
- Florin TA, French B, Zorc JJ, *et al*. Variation in emergency department diagnostic testing and disposition outcomes in pneumonia. *Pediatrics* 2013;132:237–44.
- Zachariasse JM, Nieboer D, Maconochie IK, *et al*. Development and validation of a paediatric early warning score for use in the emergency department: a multicentre study. *Lancet Child Adolesc Health* 2020;4:583–91.
- Seiger N, van Veen M, Steyerberg EW, *et al*. Accuracy of triage for children with chronic illness and infectious symptoms. *Pediatrics* 2013;132:e1602–8.
- Zachariasse JM, Kuiper JW, de Hoog M, *et al*. Safety of the Manchester triage system to detect critically ill children at the emergency department. *J Pediatr* 2016;177:232–7.
- Simon TD, Cawthon ML, Stanford S, *et al*. Pediatric medical complexity algorithm: a new method to stratify children by medical complexity. *Pediatrics* 2014;133:e1647–54.
- Group, M.T. *Emergency triage*. 3rd ed. Wiley Blackwell, 2014: 1–204.
- Schinkelshoek G, Borensztajn DM, Zachariasse JM, *et al*. Management of children visiting the emergency department during out-of-office hours: an observational study. *BMJ Paediatr Open* 2020;4:e000687.
- Turner NML, Piet L. *Advanced pediatric life support*. 5th ed. Bohn Stafleu van Loghum, 2017.
- Donders ART, van der Heijden GJMG, Stijnen T, *et al*. Review: a gentle introduction to imputation of missing values. *J Clin Epidemiol* 2006;59:1087–91.
- Raven MC, Steiner F. A National study of outpatient health care providers' effect on emergency department visit acuity and likelihood of hospitalization. *Ann Emerg Med* 2018;71:728–36.
- Coster JE, Turner JK, Bradbury D, *et al*. Why do people choose emergency and urgent care services? a rapid review utilizing a systematic literature search and narrative synthesis. *Acad Emerg Med* 2017;24:1137–49.
- Lang T, Davido A, Diakité B, *et al*. Non-Urgent care in the hospital medical emergency department in France: how much and which health needs does it reflect? *J Epidemiol Community Health* 1996;50:456–62.
- Kraaijvanger N, van Leeuwen H, Rijpsma D, *et al*. Motives for self-referral to the emergency department: a systematic review of the literature. *BMC Health Serv Res* 2016;16:685.
- Nicholson E, McDonnell T, De Brún A, *et al*. Factors that influence family and parental preferences and decision making for unscheduled paediatric healthcare - systematic review. *BMC Health Serv Res* 2020;20:663.
- Lam JH, Pickles K, Stanaway FF, *et al*. Why clinicians overtest: development of a thematic framework. *BMC Health Serv Res* 2020;20:1011.
- Brown SR, Brown J. Why do physicians order unnecessary preoperative tests? A qualitative study. *Fam Med* 2011;43:338–43.
- van der Weijden T, van Bokhoven MA, Dinant G-J, *et al*. Understanding laboratory testing in diagnostic uncertainty: a qualitative study in general practice. *Br J Gen Pract* 2002;52:974–80.
- Pines JM, Isserman JA, Szyld D, *et al*. The effect of physician risk tolerance and the presence of an observation unit on decision making for ED patients with chest pain. *Am J Emerg Med* 2010;28:771–9.
- Kangovi S, Barg FK, Carter T, *et al*. Understanding why patients of low socioeconomic status prefer hospitals over ambulatory care. *Health Aff* 2013;32:1196–203.
- Niles LM, Goyal MK, Badolato GM, *et al*. Us emergency department trends in imaging for pediatric nontraumatic abdominal pain. *Pediatrics* 2017;140. doi:10.1542/peds.2017-0615. [Epub ahead of print: 15 Sep 2017].
- Chime NO, Katznelson J, Gangadharan S, *et al*. Comparing practice patterns between pediatric and general emergency medicine physicians: a scoping review. *Pediatr Emerg Care* 2017;33:278–86.
- Harrison R, Manias E, Mears S, *et al*. Addressing unwarranted clinical variation: a rapid review of current evidence. *J Eval Clin Pract* 2019;25:53–65.
- Borensztajn DM, Hagedoorn NN, Rivero Calle I, *et al*. Variation in hospital admission in febrile children evaluated at the emergency department (ED) in Europe: perform, a multicentre prospective observational study. *PLoS One* 2021;16:e0244810.

**Appendix 1. ED Characteristics**

	<b>Maastad Hospital, Rotterdam, the Netherlands</b>	<b>Erasmus MC, Rotterdam, the Netherlands</b>	<b>Hospital Fernando da Fonseca, Lisbon, Portugal</b>	<b>St Mary's Hospital, London, United Kingdom</b>	<b>General Hospital, Vienna, Austria</b>
Hospital characteristics	Teaching hospital 59 pediatric beds	University hospital 60 pediatric beds	Community hospital 91 pediatric beds	University hospital 46 pediatric beds	University hospital 74 pediatric beds
Catchment area	Urban  Generally low socio-economic status	Urban  Mixed high and low socio-economic status	Mixed urban and rural Generally low socio-economic status	Urban  Mixed high and low socio-economic status	Urban  Mixed high and low socio-economic status
Emergency department characteristics	Mixed adult-pediatric  9500 children/year	Pediatric only till October 2014, from then on mixed  6500 children/year	Pediatric only  60,000 children/year	Pediatric only  27,000 children/year	Pediatric only  22,000 children/year
Supervising physician	Pediatrician	Pediatrician	Pediatrician	Pediatric emergency physician	Pediatrician
Inclusion period	01-05-2014 to 31-10-2015	01-01-2012 to 31-12-2014	01-03-2014 to 28-02-2015	01-07-2014 to 28-02-2015	01-01-2014 to 31-12-2014
Number of patients included	10,484	13,968	53,175	15,027	19,268
Primary care availability	24/7	24/7	Daytime and evenings	Daytime and evenings	Daytime
Referral by emergency service	4.5%	8.9%	4.0%	5.6%	Not available
Self-referral	17%	27%	96%	82%	>90%*
Comorbidity in all children	<10%*	38%	<10%*	11%	10%

## Appendix 2. MTS presentational flow charts reclassified into 10 presentational flow chart categories

Category	MTS presentational flow charts[15]
Cardiac	Chest pain, palpitations
Dermatologic	Abscesses and local infections, bites and stings, burns and scalds, rashes, Wounds
Ear Nose Throat	Ear problems, facial problems, sore throat
Gastrointestinal	Abdominal pain in adults, abdominal pain in children, diarrhoea and vomiting, gastrointestinal bleeding
Neurologic, psychiatric and intoxications	Apparently drunk, behaving strangely, collapsed adult, fits, Headache, mental illness, overdose and poisoning, self-harm
Respiratory	Asthma, shortness of breath in adults, shortness of breath in children
Trauma/Muscular	Assault, back pain, falls, head injury, limping child, limb problems, major trauma, neck pain, torso injury
Unwell	Crying baby, irritable child, unwell adult, unwell child, worried parent
Urinary/gynaecological	Pregnancy, per vaginum bleeding, sexually acquired infection, testicular pain, urinary problems
Other	Allergy, dental problems, diabetes, eye problems, exposure to chemicals, foreign body, major incidents