

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (http://bmjopen.bmj.com).

If you have any questions on BMJ Open's open peer review process please email <a href="mailto:info.bmjopen@bmj.com">info.bmjopen@bmj.com</a>

# **BMJ Open**

# IDENTIFYING HIGH VALUE HOSPITAL CARE FOR MEDICARE BENEFICIARIES: AN OBSERVATIONAL STUDY

Journal:	BMJ Open
Manuscript ID	bmjopen-2021-053629
Article Type:	Original research
Date Submitted by the Author:	19-May-2021
Complete List of Authors:	Herrin, Jeph; Yale University School of Medicine; Yale University School of Medicine, Section of Cardiovascular Medicine Yu, Huihui; Yale University Venkatesh, Arjun; Yale University, Department of Emergency Medicine Desai, Sunita; New York University Grossman School of Medicine, Department of Population Health Thiel, Cassandra; New York University Grossman School of Medicine, Department of Population Health Lin, Zhenqiu; Yale-New Haven Hospital, Center for Outcomes Research and Evaluation Bernheim, Susannah M; Yale University School of Medicine, Department of Internal Medicine Horwitz, Leora; New York University Grossman School of Medicine, Center for Healthcare Innovation and Delivery Science
Keywords:	Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, STATISTICS & RESEARCH METHODS, Quality in health care < HEALTH SERVICES ADMINISTRATION & MANAGEMENT
	·

SCHOLARONE™ Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our licence.

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which Creative Commons licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

# IDENTIFYING HIGH VALUE HOSPITAL CARE FOR MEDICARE BENEFICIARIES: AN OBSERVATIONAL STUDY

Jeph Herrin, Ph.D.<sup>1</sup>

Huihui Yu, Ph.D.<sup>2</sup>

Arjun Venkatesh<sup>3</sup>

Sunita Desai4

Zhengui Lin<sup>1,5</sup>

Susannah Bernheim<sup>5,6</sup>

Leora I. Horwitz<sup>7,8,9</sup>

- 1. Section for Cardiovascular Medicine, Yale University School of Medicine, New Haven CT, USA
- 2. Yale University School of Medicine, New Haven CT, USA
- 3. Department of Emergency Medicine, Yale University School of Medicine, New Haven, CT, USA.
- 4. Department of Population Health, New York University Grossman School of Medicine, New York, NY, USA
- 5. Center for Outcomes Research and Evaluation, Yale New Haven Health, New Haven, CT.
- 6. Section of General Internal Medicine, Department of Medicine, Yale School of Medicine, New Haven, CT
- 7. Center for Healthcare Innovation and Delivery Science, NYU Langone Health, New York, New York.
- 8. Department of Population Health, NYU Grossman School of Medicine, New York, New York.
- 9. Department of Medicine, NYU Grossman School of Medicine, New York, New York.

# **Corresponding Author:**

Jeph Herrin

Jeph.herrin@yale.edu

**Keywords**: Healthcare Value, Quality of Care

Words: 3, 054

#### **ABSTRACT**

**Objectives** High value care is providing high quality care at low cost; we sought to define hospital value and identify the characteristics of hospitals which provide high value care.

**Design**: Retrospective observational study.

**Setting**: Acute care hospitals in the U.S.

**Participants**: All Medicare beneficiaries with claims included in Center for Medicare & Medicaid Services

Overall Star Ratings or in publicly available Medicare spending per beneficiary data.

Primary and secondary outcome measures: Our primary outcome was value defined as the difference between Star Ratings quality score and Medicare spending; the secondary outcome was classification as a 4 or 5 Star hospital with lowest quintile Medicare spending ("high value") or 1 or 2 Star hospital with highest quintile spending ("low value").

**Results**: 2,914 hospitals had both quality and spending data, and were included. ). The value score had a mean (SD) of 0.58 (1.79). A total of 286 hospitals were classified as high value; these represented 28.6% of 999 4 and 5-star hospitals and 46.8% of 611 low cost hospitals. A total of 258 hospitals were classified as low value; these represented 26.6% of 970 1 and 2-star hospitals and 49.3% of 523 high cost hospitals. In regression models ownership, non-teaching status, beds, urbanity, nurse to bed ratio, percentage of dual eligible Medicare patients, and smaller of DSH payments were associated with the primary value score.

**Conclusions**: There are high quality hospitals that are not high value, and a number of factors are strongly associated with being low or high value. These findings can inform efforts of policymakers and hospitals to increase the value of care.

## **ARTICLE SUMMARY**

# **Strengths and Limitations**

- This study incorporates a comprehensive measure of overall hospital quality derived from 5 distinct domains.
- This study uses payments as proxy for actual costs; however, these payments do include post discharge utilization.
- Two different measures of value are found to be consistent, and explained by factors that hospitals can modify.

# **Funding**

This work was funding by a grant from the Agency for Healthcare Quality and Research (R01HS022882).

#### INTRODUCTION

The past two decades have seen an increasing focus by United States policymakers and caregivers on improving hospital quality of care. The movement began in the early 2000s with public reporting of process measures, moved on to public reporting of outcomes measures such as readmissions, complications and mortality and, particularly since the passage of the Affordable Care Act in 2010, has since evolved to tying quality of care to payment.[1] There are now more than 100 quality measures on the CMS's Care Compare, and a summary of over 50 of these measures into an overall Hospital Quality Star Rating.[2] Patients and policy makers thus have available a range of quality metrics for comparing hospitals.

However, only recently have policymakers begun to take on the challenge of driving both improved quality and lower costs in tandem. The Value Based Purchasing Program (for hospitals) and the Quality Payment Program (for clinicians), for instance, both incorporate some measures of cost along with quality measures into their overall scoring system. Yet despite prior research indicating that people often perceive higher quality to be associated with higher costs and policymakers' concerns that cost reduction efforts will lower quality, the relationship between hospital quality and hospital cost, and in turn value, remains empirically underexplored. While in some cases high cost care is clearly of low quality (i.e., excess use of unnecessary care, higher complication rates), it may be that higher costs produce better outcomes in other situations. For example, higher spending may focus on high-value care interventions or services such as post-discharge planning or expensive, evidence based treatments such as PCI and ICDs.[3,4,5] Moreover, even if there is no systemic association between quality and cost, it is plausible that among hospitals that provide high quality care, some will be more cost effective at providing this care than others. Characteristics of hospitals that produce high value healthcare, therefore, as defined by quality accounting for cost, may not be the same as those that produce high quality alone.

We therefore sought to identify hospitals that provide 'high value' care and to examine how they differ from other hospitals, including specifically those which provide 'low value' care. Using the CMS Overall Hospital Quality Star Ratings to identify high and low quality care hospitals and the Medicare Spending per Beneficiary (MSPB) total cost of care scores to identify high and low cost hospitals, we created a value score as the difference between quality and cost scores. Using those hospitals that were particularly high or low value, we analyzed the characteristics of those hospitals, including anticipated drivers of costs (total expenses, supply costs) and factors historically associated with quality (beds, nurse staffing, teaching status, ownership, geographic region, and urban status).

## **METHODS**

Data and Cohort

We used publicly-reported Star Ratings data from Hospital Compare for 2018. The Star Rating system assigns each hospital an integer from 1 to 5, with 1 reflecting the lowest quality and 5 reflecting the highest quality. The Star Rating is a composite that reflects hospital performance across 57 measures, grouped into any of seven quality domains mortality, readmission, safety, patient experience, efficiency, effectiveness of care, and timeliness of care. Measures in each domain are used to estimate a latent variable model with a single latent variable; and the latent variables (group scores) for each domain are combined using a weighted average to generate an overall summary score. Finally, k-means clustering with 5 means is used to group the summary scores into five star categories.[6] A hospital was assigned an overall Star Rating if it reported at least 3 measures in at least 3 domains, one of which was safety, mortality or readmission.

Hospital-level Star Ratings data were linked to 2018 MSPB data using hospitals' CMS Certification numbers (CCNs). The MSPB score is calculated by the Center for Medicare & Medicaid Services (CMS) using Medicare Part A and Part B payments for all care provided from 3 days prior to, to 30 days

following an inpatient hospital stay (defined as an "episode" of care). MSPB episodes span all conditions, and MSPB scores are calculated for all Medicare fee-for-service hospital stays, excluding stays in psychiatric, rehabilitation, cancer, children's, critical access or long-stay hospitals; excluding hospitalizations resulting in transfers to another acute facility; and excluding readmissions that are within 30 days of an eligible hospitalization. Episodes in which the patient dies during the measurement period are also excluded, as are those with \$0 payments or involving transfers. The measure adjusts for the MS-DRG of the index hospitalization, age, use of long-term care, and 79 comorbidities. All payments are standardized across geographies for comparability; additional payments such as for indirect medical education or disproportionate share payments are omitted.[7] The MSPB scores is constructed as a ratio so that a value of 1 is "average" spending, and values lower or higher than 1 represent lower or higher spending than average.

Our study cohort comprised all hospitals which had an overall Star Rating; none were missing MSPB scores.

Value

We examined two related measures of value. For our first value outcome, we constructed a linear metric of value by subtracting the standardized MSPB score from the standardized Star Rating hospital summary score (used to create the 5 star categories). Though we initially considered a ratio of quality to spending, we were concerned that a ratio could allow hospitals to appear high value while having low quality, if their spending were low enough. Therefore, we used a difference measure, which ensures that hospitals categorized as high value were always of high quality. To facilitate interpretation, both spending and quality scores were standardized to range from 0 to 10; thus, higher scores indicate higher value, with the maximum value score being 10 and minimum -10.

For our second value outcome, we classified hospitals as 'high value' if they received a 4 or 5-star rating on Hospital Compare and had a MSPB score in the lowest quintile of all hospitals, and classified hospitals as 'low value' if they received a 1 or 2-star rating on Hospital Compare and had a MSPB score in the highest quintile of all hospitals; all other hospitals were classified as 'average' value. For this classification, we selected quintiles to categorize hospitals by their MSPB scores as providing meaningful cost distinctions while identifying adequate numbers of high and low value hospitals for analysis.

### Other Variables

Hospital characteristics were linked from the American Hospital Association (AHA) survey for 2018[8] and the CMS Provider of Services (POS) file of 2018. We examined hospital characteristics that have previously been associated with quality: geographic region (New England, Middle Atlantic, East North Central, West North Central, South Atlantic, East South Central, West South Central, Mountain, and Pacific); teaching status (teaching, residency program, non-teaching); number of beds(<100, 100-199,200-299,300-399, and 400+); nurse to bed ratio (≤0.75, 0.75-1, 1-1.5, > 1.5); urban location (urban, rural); ownership status (for profit, private not for profit, public, federal); percentage of Medicare admissions that are for patients dually eligible for both Medicare and Medicaid (≤ 10%, 11%-20%, 21%-40%, and > 40%); total expenses (quintiles); total cost of supplies (quintiles), ratio of supply expenses to total expenses (quintiles); and disproportionate share hospital (DSH) patient percentage (quintiles). [9]

# Patient and Public Involvement

There was no patient or public involvement in the design, conduct or reporting of this study.

Statistical Analysis

We summarized value score (mean and standard deviation) and value category (frequency and percent) overall and by hospital characteristics, testing for differences in category characteristic using ANOVA for the value score and chi-square tests for the value category. To assess the overall relationship between cost and quality of care, we graphed the distribution of MSPB scores over each Star rating using a box plot (Figure 1), and then plotted the value scores against the Star Rating summary scores to visualize the association of value with quality (Figure 2). Then, to identify the hospital characteristics independently associated with value, and to identify contrasts in associations with quality, we estimated a series of models. We first used bivariate linear regression models to estimate the relationship between each hospital characteristics and value, where the outcome was our continuous value score; all factors were then included in a final multivariate model. For comparison with factors associated with quality alone, we estimated identical models where the dependent variable was the Star Ratings summary score, rather than our value score. We next estimated a multinomial logit model with the 3-category value as the outcome using 'average' as the reference group, including all factors. Both analyses included the same set of hospital characteristics, and we reported overall Wald test p-values for each characteristic. We report numbers and percent of missing values, and use multiple imputation (with 20 imputations) to account for missing values in all models.

All analyses were done using SAS 9.4 and Stata 16.1 (2020, StataCorp, College Station TX). All statistical tests were two-tailed, and we used p<0.05 to determine statistical significance. The study was approved by the Yale and NYU Institutional Review Boards.

#### **RESULTS**

Our final study sample included 2,914 hospitals that received both a Star Rating and a MSPB score, all of which matched the AHA Survey file for 2018 (Table 1). The value score had a mean (SD) of 0.58 (1.79); see Figures 1 and 2. A total of 286 were classified as high value and 258 as low value by the categorical definition (Table 1). The 286 high value hospitals represented 28.6% of 999 4 and 5-star hospitals and 46.8% of 611 low cost (bottom quintile MSPB) hospitals; the 258 low value hospitals represented 26.6% of 970 1 and 2-star hospitals and 49.3% of 523 high cost hospitals.

In biviariate analyses (Table 1), we found not-for-profit private and government owned hospitals to have much higher values scores than for profit private hospitals, with score of 0.39 and 1.38 versus 0.08. Non-teaching hospitals and those with residency programs both had three times the value score of teaching hospitals, while hospitals with fewer than 100 beds (1.48) and less than 10% dual eligible patients (1.74) had the highest values scores. Hospitals in rural areas and smaller percentages of DSH payments also had higher value scores (all P-values < 0.05). For the categorical bivariate comparisons (Table 1), results were similar: high value hospitals were more likely to be not-for-profit, have no teaching program, have more beds and a higher nurse to bed ratio, and have a lower percentage of dual-eligible admissions compared to low- and average- value hospitals (all P values < 0.001). Regionally, lower value hospitals were more often in the Southern and Mid-Atlantic regions, while high value hospitals were in Northern Central and Pacific Regions (P<0.001).

In the multivariable model, all were significantly associated with value except those related to total and supply expenses (Table 2). The largest value effects were again for non-teaching, not-for-profit and government hospitals, hospitals with fewer than 100 beds and those with fewer than 10% dual eligible admissions. In equivalent model using the quality score instead of value as an outcome, there was no association with urban status, while, unlike with value, higher supply expenses were associated with higher quality; other patterns of effects were very similar.

In multinomial regression assessing the associations between characteristics and low, average, and high value classification (Table 3), we collapsed the region categories to account for sparse cells. In the final model Government owned and not-for-profit hospitals, hospitals with fewer than 100 beds, hospitals in the Western region, and those with lower percentages of dual eligible patients and DSH patients were more likely to be high value than average value (all P-values < 0.05). Correspondingly, public hospitals, those with the more beds, those in the Northeast region, and lowest nurse to bed ratio all had higher odds of being low-value (all P-values < 0.05).

# **DISCUSSION**

In this examination of hospital value of care, we identified hospitals that provided relatively higher quality of care to Medicare beneficiaries at lower comparative costs, as well as hospitals that provided lower quality care at a high level of spending. In bivariate analyses, both high and low value hospitals differed from other hospitals strongly in ownership, urban location, bed size and region, nurse-to-bed ratio, amount spent on supplies relative to total expenses, and percentage of dual-eligible patients. Most associations were significant in the multivariable primary analysis using the value score; of these, all but those related to urbanicity and expenses persisted in the multinomial logit analysis using a categorical outcome of low, average or high value.

The most notable finding, however, may not be the associations of hospital characteristics with value, but that value and quality of care are not synonymous: though generally, the MSPB declined with increasing Star ratings (Figure 1), less than a third of 4 and 5-star hospitals were high value, while more than a fourth of 1 and 2-star hospitals were lowest value. The observed trend in spending across overall quality (Figure 1) as well as the pattern of value classification is very similar to that found previously for patient experience Star Ratings [10,11]; though the patient experience Star Rating is a component of the overall Star Rating, it accounts for only 22% of the weight of the overall summary score, so it is unlikely

to drive the current findings. This finding was also reflected in our model of quality score, which found urban status was not related to quality while quality increased with more spending on supplies. That quality and value are not identical indicates that it will be possible to reduce spending among even the highest quality hospitals.

Most of the associations we found for value are similar to those seen for hospital characteristics and quality, both here with the Star Ratings summary score and in prior research; for example, Delancy et. al found that hospitals with fewer beds, no medical school affiliation, and lower proportions of DSH patients were more likely to have higher Star ratings [12]. This was similar to the results of our analysis of the star summary quality score (Table 2). However, in our categorical analysis, we found no relationship between teaching status and high value. Geographic region also explained some of the difference between low value and average value hospitals across all of our analyses, with hospitals in East South Central and Mid-Atlantic regions having lowest value and being more likely to be low value than those in any other regions, and Pacific region hospitals being consistently high value in all models. Since our spending measure is adjusted for local wage and cost variations, this geographic variation is likely driven substantially by variations in quality.

Of most relevance to hospitals and policy makers are the associations with factors that are within the control of the hospital. A key factor that was related to low value in all analyses was nurse to bed ratio, with lower ratios being more often found at the lowest value hospitals — only 9 (3.5%) low value hospitals had a ratio of 2 or more, while 43 (15.0%) of high value hospitals had a ratio of at least 2. This finding suggests that high levels of nurse staffing, while increasing the expense of inpatient care, may reduce overall Medicare payments while promoting high quality care. Importantly, this finding also highlights a perennial challenge for policymakers: the wrong pocket problem. Since the costs of increased staffing are borne by hospitals but the benefits of lower spending are reaped by payers and

patients, there is no intrinsic incentive for hospitals to pursue such interventions. Payment policies seeking to improve value should explicitly take these challenges into account.

Conversely, 66 (25.6%) of low value hospitals reported at least 40% of their Medicare patients were also eligible for Medicaid, while only 11 (3.8%) of high value hospitals reported such high rates of dual eligibility, with proportion of DSH patients exhibiting a similar but weaker pattern. This may indicate that lower income patients have unmeasured comorbidities or care complexities that are not captured by the risk adjustment used in Star Ratings' underlying measures or that these patients drive costs in ways unrelated to quality of care (e.g., limited social support may lead to greater post-discharge costs), that these patients may have higher cost needs that are not completely adjusted for, or possibly that DSH patients are more likely to receive care at lower quality hospitals.

This study has the limitations of any observational study, including that no causal inferences can be drawn; for example, it may be that lower value hospitals have increased costs of care because of lower quality, rather than independent of it. In addition, we have only examined a few key factors that may be associated with quality, in order to better focus on value of care. And, we have focused only on quality and spending metrics largely related to Medicare fee-for-service beneficiaries; other, broader measures of quality and costs that included the entire patient population could plausibly produce different results. Moreover, Medicare payments are distorted somewhat by the Inpatient Prospective Payment System. However, CMS Overall Star Ratings captures quality of care across 7 domains, and the MSPB is a measure of spending with cost-of-living adjustments that plausibly represents the overall spending per patient resulting from a hospitalization.

It is worth considering that the MSPB does not include admissions that lead to transfers, nor admissions in which the patient does not survive 30 days. Both exclusions are likely to confound the relationship between the spending score and the mortality domain, which accounts for 22% of the Star Rating summary score, since publicly reported mortality measures typically measure outcomes within

the same 30 days, and attribute them to the first hospital in the event of transfers. Excluding patients who do not survive and/or are transferred likely reduces the MSPB score (assuming these are more complex patients), and more so for hospitals with higher mortality or transfer rates. However, this would tend to bias our current findings toward the null, attenuating the relationship between spending value.

#### CONCLUSION

We have shown that care quality and care value are not identical; there are high quality hospitals that are not high value. In addition, there are a number of structural factors, some under the control of hospitals such as nurse to bed ratio, that are strongly associated with being low or high value. These findings can inform efforts of policymakers and hospitals to increase the value of care.

#### REFERENCES

- U.S. Centers for Medicare & Medicaid Services. Hospital Readmissions Reduction Program (HRRP).
   Accessed 2020 Dec 08. <a href="https://www.cms.gov/medicare/medicare-fee-for-service-payment/acuteinpatientpps/readmissions-reduction-program.html">https://www.cms.gov/medicare/medicare-fee-for-service-payment/acuteinpatientpps/readmissions-reduction-program.html</a>
- U.S. Centers for Medicare and Medicaid Services. Hospital Compare Overall Ratings. Accessed 2020
   Dec 08. https://qualitynet.cms.gov/inpatient/public-reporting/overall-ratings
- Doyle JJ, Graves JA, Gruber JA, Kleiner S. Do high-cost hospitals deliver better care? Evidence from ambulance referral patterns. National Bureau of Economic Research. http://www.nber.org/papers/w17936.pdf?new\_window=1. Accessed December 19, 2013.
- 4. Silber JH, Kaestner R, Even-Shoshan O, Wang Y, Bressler LJ. Aggressive treatment style and surgical outcomes. Health Serv Res. 2010;45(6 Pt 2):1872-1892.
- 5. Virtual Mentor. 2014;16(2):124-130.
- 6. QualityNet. https://www.qualitynet.org/inpatient/public-reporting/overall-ratings/resources
- 7. QualityNet. https://www.qualitynet.org/inpatient/measures/mspb/methodology
- American Hospital Association. American Hospital Association (AHA) Annual Survey Database –
   2018.
- 9. https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/dsh
- Liao JM, Zhou L, Navathe AS. Nationwide Hospital Performance on Publicly Reported Episode
   Spending Measures. J Hosp Med. 2020 Mar 18;15:E1-E7. doi: 10.12788/jhm.3377. Epub ahead of print. PMID: 32195657
- 11. Trzeciak S, Gaughan JP, Bosire J, Angelo M, Holzberg AS, Mazzarelli AJ. Association Between Medicare Star Ratings for Patient Experience and Medicare Spending per Beneficiary for US Hospitals. J Patient Exp. 2017 Mar;4(1):17-21. doi: 10.1177/2374373516685938. Epub 2017 Jan 12. PMID: 28725856; PMCID: PMC5513662

12. DeLancey JO, Softcheck J, Chung JW, Barnard C, Dahlke AR, Bilimoria KY. Associations Between Hospital Characteristics, Measure Reporting, and the Centers for Medicare & Medicaid Services Overall Hospital Quality Star Ratings. JAMA. 2017 May 16;317(19):2015-2017. doi: 10.1001/jama.2017.3148. PMID: 28510670; PMCID: PMC5815009.



**Table 1**. Characteristics of low, average, and high value hospitals.

Value					
Score Mean (SD)	P-value <sup>1</sup>	Low Value n(%) or mean(SD)	Average n(%) or mean(SD)	High Value n(%) or mean(SD)	P-value
0.58 (1.79)		258 (100.0)	2370 (100.0)	286 (100.0)	
	< 0.001				
-2.38 (1.26)		89 (34.5)	179 (7.6)	0 (0.0)	
-0.66 (1.02)		169 (65.5)	533 (22.5)	0 (0.0)	
0.71 (1.00)		0 (0.0)	945 (39.9)	0 (0.0)	
1.75 (1.00)		0 (0.0)	528 (22.3)		
3.16 (1.07)		0 (0.0)			
, ,	< 0.001	` ,	` '	` ,	< 0.001
0.39 (1.80)		43 (16.7)	356 (15.0)	33 (11.5)	
			, ,		
		• •			
2.00 (2.00)	< 0.001	0 (0.0)	20 (0.0)	_ (0)	< 0.001
0.87 (1.72)		22 (8.5)	381 (16.1)	48 (16.8)	
0.23 (2.02)	<0.001	112 (13.1)	033 (30.2)	70 (2 1.3)	< 0.001
1 48 (1 57)	10.001	21 (8 1)	575 (24 3)	155 (54 2)	10.001
, ,			, ,		
0.12 (1.74)	<0.001	07 (20.0)	477 (20.1)	14 (4.5)	< 0.001
1 08 (1 52)	\0.001	29 (11 2)	579 (24 4)	110 (38 5)	10.001
			· ·		
0.41 (1.04)	<0.001	223 (00.0)	1731 (73.0)	170 (01.5)	<0.001
0.70 (1.48)	<b>\0.001</b>	3 (1 2)	115 (4 9)	9 (3 1)	<b>\0.001</b>
			1 1		
0.81 (1.91)	10.001	32 (12.4)	291 (12.3)	53 (18.5)	10.001
0.54 (4.00)	<0.001	06 (22.2)	F2C (22 C)	70 (27 5)	<0.001
				• •	
		• •			
1.24 (1.67)		9 (3.5)	265 (11.2)	43 (15.0)	
	Mean (SD)  0.58 (1.79)  -2.38 (1.26) -0.66 (1.02) 0.71 (1.00)	Score Mean (SD)         P-value¹           0.58 (1.79)         <0.001	Score Mean (SD)         P-value¹         Low Value n(%) or mean(SD)           0.58 (1.79)         <0.001	Score Mean (SD)         P-value¹         Low Value n(%) or mean(SD)         Average n(%) or mean(SD)           0.58 (1.79)         258 (100.0)         2370 (100.0)           -2.38 (1.26)         89 (34.5)         179 (7.6)           -0.66 (1.02)         169 (65.5)         533 (22.5)           0.71 (1.00)         0 (0.0)         945 (39.9)           1.75 (1.00)         0 (0.0)         528 (22.3)           3.16 (1.07)         0 (0.0)         185 (7.8)           -0.39 (1.80)         43 (16.7)         356 (15.0)           0.77 (1.69)         128 (49.6)         1521 (64.2)           0.08 (1.96)         87 (33.7)         478 (20.2)           1.38 (1.60)         0 (0.0)         15 (0.6)           <0.001	Score Mean (SD)         P-value¹         Low Value n(%) or mean(SD)         Average mean(SD)         High Value n(%) or mean(SD)           0.58 (1.79)         258 (100.0)         2370 (100.0)         286 (100.0)           -2.38 (1.26)         89 (34.5)         179 (7.6)         0 (0.0)           -0.66 (1.02)         169 (65.5)         533 (22.5)         0 (0.0)           0.77 (1.00)         0 (0.0)         945 (39.9)         0 (0.0)           1.75 (1.00)         0 (0.0)         185 (7.8)         73 (25.5)           0.001         0 (0.0)         185 (7.8)         73 (25.5)           0.39 (1.80)         43 (16.7)         356 (15.0)         33 (11.5)           0.77 (1.69)         128 (49.6)         1521 (64.2)         215 (75.2)           0.08 (1.96)         87 (33.7)         478 (20.2)         36 (12.6)           1.38 (1.60)         0 (0.0)         15 (0.6)         2 (0.7)           0.72 (1.75)         122 (8.5)         381 (16.1)         48 (16.8)           0.72 (1.75)         121 (8.1)         575 (24.3)         155 (54.2)           0.65 (1.65)         58 (22.5)         645 (27.2)         67 (23.4)           0.21 (1.73)         54 (20.9)         399 (16.8)         33 (11.5)           0.08 (1.86)

Table 1 continued

% Duals		< 0.001				< 0.001
<=10%	1.74 (1.93)		7 (2.7)	176 (7.4)	33 (11.5)	
11%-20%	0.80 (1.58)		67 (26.0)	863 (36.4)	91 (31.8)	
21%-40%	0.51 (1.71)		118 (45.7)	1057 (44.6)	151 (52.8)	
>40%	-0.56 (1.94)		66 (25.6)	274 (11.6)	11 (3.8)	
Total Expenses		< 0.001				< 0.001
Quintile 1	1.09 (1.66)		28 (10.9)	459 (19.4)	95 (33.2)	
Quintile 2	0.79 (1.77)		54 (20.9)	446 (18.8)	81 (28.3)	
Quintile 3	0.44 (1.73)		57 (22.1)	475 (20.0)	50 (17.5)	
Quintile 4	0.31 (1.81)		71 (27.5)	473 (20.0)	37 (12.9)	
Quintile 5	0.24 (1.83)		48 (18.6)	511 (21.6)	22 (7.7)	
Missing	0.68 (1.61)		0 (0.0)	6 (0.3)	1 (0.3)	
Supplies \$		< 0.001				< 0.001
Quintile 1	1.11 (1.63)		23 (8.9)	301 (12.7)	72 (25.2)	
Quintile 2	0.83 (1.64)		31 (12.0)	311 (13.1)	53 (18.5)	
Quintile 3	0.50 (1.82)		35 (13.6)	326 (13.8)	35 (12.2)	
Quintile 4	0.59 (1.72)		37 (14.3)	334 (14.1)	24 (8.4)	
Quintile 5	0.27 (1.81)		34 (13.2)	349 (14.7)	12 (4.2)	
Missing	0.40 (1.86)		98 (38.0)	749 (31.6)	90 (31.5)	
Supplies/Total %		0.042				0.002
Quintile 1	0.83 (1.79)		29 (11.2)	311 (13.1)	56 (19.6)	
Quintile 2	0.75 (1.69)		23 (8.9)	329 (13.9)	43 (15.0)	
Quintile 3	0.51 (1.75)		38 (14.7)	317 (13.4)	41 (14.3)	
Quintile 4	0.53 (1.56)		28 (10.9)	344 (14.5)	23 (8.0)	
Quintile 5	0.68 (1.92)		42 (16.3)	320 (13.5)	33 (11.5)	
Missing	0.40 (1.86)		98 (38.0)	749 (31.6)	90 (31.5)	
% DSH		< 0.001				< 0.001
Quintile 1	1.35 (1.74)		32 (12.4)	468 (19.7)	84 (29.4)	
Quintile 2	0.85 (1.62)		33 (12.8)	480 (20.3)	70 (24.5)	
Quintile 3	0.63 (1.59)		42 (16.3)	488 (20.6)	52 (18.2)	
Quintile 4	0.30 (1.60)		55 (21.3)	484 (20.4)	44 (15.4)	
Quintile 5	-0.26 (1.96)		96 (37.2)	450 (19.0)	36 (12.6)	
	•					

Source: Center for Medicare and Medicaid Services Overall Star Ratings score and Medicare Spending per Beneficiary; American Hospital Association Survey.

# Notes

1. P-value based on ANOVA model estimated using multiple imputation for missing values.

**Table 2**. Results of multivariable linear models with value score and Star Ratings summary score as dependent variables.

Characteristic         Coeff (SE)         P-value           Ownership         <0.001         <0.001           Public         ref         ref           Not for profit         0.41 (0.22,0.60)         0.35 (0.21,0.49)           For profit         0.02 (-0.23,0.27)         0.19 (0.00,0.38)           Government         1.07 (-0.11,2.24)         0.43 (-0.45,1.31)           Teaching status         0.001         0.009           Nonteaching         ref         ref           Residency         -0.30 (-0.50,-0.10)         -0.20 (-0.35,-0.05)           Teaching         -0.25 (-0.45,-0.05)         -0.23 (-0.38,-0.08)           Beds         <0.001         <0.001           <100         ref         ref           100-199         -0.58 (-0.80,-0.36)         -0.34 (-0.50,-0.17)           200-299         -1.02 (-1.32,-0.73)         -0.61 (-0.83,-0.39)           300-399         -1.25 (-1.59,-0.92)         -0.71 (-0.96,-0.45)           400+         -1.47 (-1.84,-1.10)         -0.83 (-1.11,-0.56)           Urbanity         0.026         0.941           Rural         ref         ref           Urban         -0.21 (-0.40,-0.03)         0.01 (-0.13,0.15)           Region         <0.001<		Value		Quality	
Public         ref         ref           Not for profit         0.41 (0.22,0.60)         0.35 (0.21,0.49)           For profit         0.02 (-0.23,0.27)         0.19 (0.00,0.38)           Government         1.07 (-0.11,2.24)         0.43 (-0.45,1.31)           Teaching status         0.010         0.009           Nonteaching         ref         ref           Residency         -0.30 (-0.50,-0.10)         -0.20 (-0.35,-0.05)           Teaching         -0.25 (-0.45,-0.05)         -0.23 (-0.38,-0.08)           Beds         <0.001            *100         ref         ref           100-199         -0.58 (-0.80,-0.36)         -0.34 (-0.50,-0.17)           200-299         -1.02 (-1.32,-0.73)         -0.61 (-0.83,-0.39)           300-399         -1.25 (-1.59,-0.92)         -0.71 (-0.96,-0.45)           400+         -1.47 (-1.84,-1.10)         -0.83 (-1.11,-0.56)           Urbanity         0.026         0.941           Rural         ref         ref           Urban         -0.21 (-0.40,-0.03)         0.01 (-0.13,0.15)           Region         <0.001         <0.001           New England         ref         ref           Mid Atlantic         -0.67 (-1.02,-0.31)	Characteristic	Coeff (SE)	P-value		
Not for profit	Ownership		<0.001		<0.001
For profit         0.02 (-0.23,0.27)         0.19 (0.00,0.38)           Government         1.07 (-0.11,2.24)         0.43 (-0.45,1.31)           Teaching status         0.010         0.009           Nonteaching         ref         ref           Residency         -0.30 (-0.50,-0.10)         -0.20 (-0.35,-0.05)           Teaching         -0.25 (-0.45,-0.05)         -0.23 (-0.38,-0.08)           Beds         <0.0001	Public	ref		ref	
Government         1.07 (-0.11,2.24)         0.43 (-0.45,1.31)           Teaching status         0.010         0.009           Nonteaching         ref         ref           Residency         -0.30 (-0.50,-0.10)         -0.20 (-0.35,-0.05)           Teaching         -0.25 (-0.45,-0.05)         -0.23 (-0.38,-0.08)           Beds         <0.001	Not for profit	0.41 (0.22,0.60)		0.35 (0.21,0.49)	
Teaching status         0.010         0.009           Nonteaching         ref         ref           Residency         -0.30 (-0.50,-0.10)         -0.20 (-0.35,-0.05)           Teaching         -0.25 (-0.45,-0.05)         -0.23 (-0.38,-0.08)           Beds         <0.001	For profit	0.02 (-0.23,0.27)		0.19 (0.00,0.38)	
Nonteaching         ref         ref           Residency         -0.30 (-0.50,-0.10)         -0.20 (-0.35,-0.05)           Teaching         -0.25 (-0.45,-0.05)         -0.23 (-0.38,-0.08)           Beds         <0.001	Government	1.07 (-0.11,2.24)		0.43 (-0.45,1.31)	
Residency	Teaching status		0.010		0.009
Teaching         -0.25 (-0.45,-0.05)         -0.23 (-0.38,-0.08)           Beds         <0.001	Nonteaching	ref		ref	
Beds       <0.001       <0.001         <100	Residency	-0.30 (-0.50,-0.10)		-0.20 (-0.35,-0.05)	
<100	Teaching	-0.25 (-0.45,-0.05)		-0.23 (-0.38,-0.08)	
100-199       -0.58 (-0.80,-0.36)       -0.34 (-0.50,-0.17)         200-299       -1.02 (-1.32,-0.73)       -0.61 (-0.83,-0.39)         300-399       -1.25 (-1.59,-0.92)       -0.71 (-0.96,-0.45)         400+       -1.47 (-1.84,-1.10)       -0.83 (-1.11,-0.56)         Urbanity       0.026       0.941         Rural       ref       ref         Urban       -0.21 (-0.40,-0.03)       0.01 (-0.13,0.15)         Region       <0.001	Beds		< 0.001		< 0.001
200-299	<100	ref		ref	
300-399	100-199	-0.58 (-0.80,-0.36)		-0.34 (-0.50,-0.17)	
400+       -1.47 (-1.84,-1.10)       -0.83 (-1.11,-0.56)         Urbanity       0.026       0.941         Rural       ref       ref         Urban       -0.21 (-0.40,-0.03)       0.01 (-0.13,0.15)         Region       <0.001	200-299	-1.02 (-1.32,-0.73)		-0.61 (-0.83,-0.39)	
Urbanity         0.026         0.941           Rural         ref         ref           Urban         -0.21 (-0.40,-0.03)         0.01 (-0.13,0.15)           Region         <0.001	300-399	-1.25 (-1.59,-0.92)		-0.71 (-0.96,-0.45)	
Rural ref ref Urban -0.21 (-0.40,-0.03) 0.01 (-0.13,0.15)  Region <0.001 <0.001  New England ref ref Mid Atlantic -0.67 (-1.02,-0.31) -0.56 (-0.83,-0.30)  E North Central 0.36 (0.02,0.71) 0.27 (0.01,0.53)  W North Central 0.58 (0.21,0.95) 0.03 (-0.25,0.31)  South Atlantic -0.10 (-0.46,0.26) -0.26 (-0.53,0.01)  E South Central -0.70 (-1.13,-0.28) -0.55 (-0.87,-0.23)  W South Central -0.61 (-0.97,-0.24) -0.14 (-0.42,0.13)  Mountain 0.22 (-0.20,0.64) -0.03 (-0.35,0.29)  Pacific 0.84 (0.46,1.23) 0.35 (0.06,0.63)  Nurse/Beds 0.001 <0.001  <-0.75 ref ref 0.75-1 -0.18 (-0.40,0.03) -0.21 (-0.37,-0.05)  1-1.5 -0.14 (-0.33,0.05) -0.15 (-0.30,-0.01)  1.5-2 0.08 (-0.15,0.30) 0.04 (-0.13,0.21)	400+	-1.47 (-1.84,-1.10)		-0.83 (-1.11,-0.56)	
Urban -0.21 (-0.40,-0.03)	Urbanity		0.026		0.941
Region         <0.001         <0.001           New England         ref         ref           Mid Atlantic         -0.67 (-1.02,-0.31)         -0.56 (-0.83,-0.30)           E North Central         0.36 (0.02,0.71)         0.27 (0.01,0.53)           W North Central         0.58 (0.21,0.95)         0.03 (-0.25,0.31)           South Atlantic         -0.10 (-0.46,0.26)         -0.26 (-0.53,0.01)           E South Central         -0.70 (-1.13,-0.28)         -0.55 (-0.87,-0.23)           W South Central         -0.61 (-0.97,-0.24)         -0.14 (-0.42,0.13)           Mountain         0.22 (-0.20,0.64)         -0.03 (-0.35,0.29)           Pacific         0.84 (0.46,1.23)         0.35 (0.06,0.63)           Nurse/Beds         0.001         <0.001	Rural	ref		ref	
New England         ref         ref           Mid Atlantic         -0.67 (-1.02,-0.31)         -0.56 (-0.83,-0.30)           E North Central         0.36 (0.02,0.71)         0.27 (0.01,0.53)           W North Central         0.58 (0.21,0.95)         0.03 (-0.25,0.31)           South Atlantic         -0.10 (-0.46,0.26)         -0.26 (-0.53,0.01)           E South Central         -0.70 (-1.13,-0.28)         -0.55 (-0.87,-0.23)           W South Central         -0.61 (-0.97,-0.24)         -0.14 (-0.42,0.13)           Mountain         0.22 (-0.20,0.64)         -0.03 (-0.35,0.29)           Pacific         0.84 (0.46,1.23)         0.35 (0.06,0.63)           Nurse/Beds         0.001         <0.001	Urban	-0.21 (-0.40,-0.03)		0.01 (-0.13,0.15)	
Mid Atlantic       -0.67 (-1.02,-0.31)       -0.56 (-0.83,-0.30)         E North Central       0.36 (0.02,0.71)       0.27 (0.01,0.53)         W North Central       0.58 (0.21,0.95)       0.03 (-0.25,0.31)         South Atlantic       -0.10 (-0.46,0.26)       -0.26 (-0.53,0.01)         E South Central       -0.70 (-1.13,-0.28)       -0.55 (-0.87,-0.23)         W South Central       -0.61 (-0.97,-0.24)       -0.14 (-0.42,0.13)         Mountain       0.22 (-0.20,0.64)       -0.03 (-0.35,0.29)         Pacific       0.84 (0.46,1.23)       0.35 (0.06,0.63)         Nurse/Beds       0.001       <0.001	Region		<0.001		< 0.001
E North Central 0.36 (0.02,0.71) 0.27 (0.01,0.53)  W North Central 0.58 (0.21,0.95) 0.03 (-0.25,0.31)  South Atlantic -0.10 (-0.46,0.26) -0.26 (-0.53,0.01)  E South Central -0.70 (-1.13,-0.28) -0.55 (-0.87,-0.23)  W South Central -0.61 (-0.97,-0.24) -0.14 (-0.42,0.13)  Mountain 0.22 (-0.20,0.64) -0.03 (-0.35,0.29)  Pacific 0.84 (0.46,1.23) 0.35 (0.06,0.63)  Nurse/Beds 0.001 <0.001  <=0.75 ref ref 0.75-1 -0.18 (-0.40,0.03) -0.21 (-0.37,-0.05) 1-1.5 -0.14 (-0.33,0.05) -0.15 (-0.30,-0.01) 1.5-2 0.08 (-0.15,0.30) 0.04 (-0.13,0.21)	New England	ref		ref	
W North Central 0.58 (0.21,0.95) 0.03 (-0.25,0.31)  South Atlantic -0.10 (-0.46,0.26) -0.26 (-0.53,0.01)  E South Central -0.70 (-1.13,-0.28) -0.55 (-0.87,-0.23)  W South Central -0.61 (-0.97,-0.24) -0.14 (-0.42,0.13)  Mountain 0.22 (-0.20,0.64) -0.03 (-0.35,0.29)  Pacific 0.84 (0.46,1.23) 0.35 (0.06,0.63)  Nurse/Beds 0.001 <0.001  <=0.75 ref ref 0.75-1 -0.18 (-0.40,0.03) -0.21 (-0.37,-0.05)  1-1.5 -0.14 (-0.33,0.05) -0.15 (-0.30,-0.01)  1.5-2 0.08 (-0.15,0.30) 0.04 (-0.13,0.21)	Mid Atlantic	-0.67 (-1.02,-0.31)		-0.56 (-0.83,-0.30)	
South Atlantic       -0.10 (-0.46,0.26)       -0.26 (-0.53,0.01)         E South Central       -0.70 (-1.13,-0.28)       -0.55 (-0.87,-0.23)         W South Central       -0.61 (-0.97,-0.24)       -0.14 (-0.42,0.13)         Mountain       0.22 (-0.20,0.64)       -0.03 (-0.35,0.29)         Pacific       0.84 (0.46,1.23)       0.35 (0.06,0.63)         Nurse/Beds       0.001       <0.001	E North Central	0.36 (0.02,0.71)		0.27 (0.01,0.53)	
E South Central -0.70 (-1.13,-0.28) -0.55 (-0.87,-0.23)  W South Central -0.61 (-0.97,-0.24) -0.14 (-0.42,0.13)  Mountain 0.22 (-0.20,0.64) -0.03 (-0.35,0.29)  Pacific 0.84 (0.46,1.23) 0.35 (0.06,0.63)  Nurse/Beds 0.001 <0.001  <=0.75 ref ref 0.75-1 -0.18 (-0.40,0.03) -0.21 (-0.37,-0.05)  1-1.5 -0.14 (-0.33,0.05) -0.15 (-0.30,-0.01)  1.5-2 0.08 (-0.15,0.30) 0.04 (-0.13,0.21)	W North Central	0.58 (0.21,0.95)		0.03 (-0.25,0.31)	
W South Central -0.61 (-0.97,-0.24) -0.14 (-0.42,0.13)  Mountain 0.22 (-0.20,0.64) -0.03 (-0.35,0.29)  Pacific 0.84 (0.46,1.23) 0.35 (0.06,0.63)  Nurse/Beds 0.001 <0.001  <=0.75 ref ref 0.75-1 -0.18 (-0.40,0.03) -0.21 (-0.37,-0.05) 1-1.5 -0.14 (-0.33,0.05) -0.15 (-0.30,-0.01) 1.5-2 0.08 (-0.15,0.30) 0.04 (-0.13,0.21)	South Atlantic	-0.10 (-0.46,0.26)		-0.26 (-0.53,0.01)	
Mountain       0.22 (-0.20,0.64)       -0.03 (-0.35,0.29)         Pacific       0.84 (0.46,1.23)       0.35 (0.06,0.63)         Nurse/Beds       0.001       <0.001	E South Central	-0.70 (-1.13,-0.28)		-0.55 (-0.87,-0.23)	
Pacific       0.84 (0.46,1.23)       0.35 (0.06,0.63)         Nurse/Beds       0.001       <0.001         <=0.75       ref       ref         0.75-1       -0.18 (-0.40,0.03)       -0.21 (-0.37,-0.05)         1-1.5       -0.14 (-0.33,0.05)       -0.15 (-0.30,-0.01)         1.5-2       0.08 (-0.15,0.30)       0.04 (-0.13,0.21)	W South Central	-0.61 (-0.97,-0.24)		-0.14 (-0.42,0.13)	
Nurse/Beds       0.001       <0.001         <=0.75	Mountain	0.22 (-0.20,0.64)		-0.03 (-0.35,0.29)	
<=0.75       ref       ref         0.75-1       -0.18 (-0.40,0.03)       -0.21 (-0.37,-0.05)         1-1.5       -0.14 (-0.33,0.05)       -0.15 (-0.30,-0.01)         1.5-2       0.08 (-0.15,0.30)       0.04 (-0.13,0.21)	Pacific	0.84 (0.46,1.23)		0.35 (0.06,0.63)	
0.75-1       -0.18 (-0.40,0.03)       -0.21 (-0.37,-0.05)         1-1.5       -0.14 (-0.33,0.05)       -0.15 (-0.30,-0.01)         1.5-2       0.08 (-0.15,0.30)       0.04 (-0.13,0.21)	Nurse/Beds		0.001		< 0.001
1-1.5 -0.14 (-0.33,0.05) -0.15 (-0.30,-0.01) 1.5-2 0.08 (-0.15,0.30) 0.04 (-0.13,0.21)	<=0.75	ref		ref	
1.5-2 0.08 (-0.15,0.30) 0.04 (-0.13,0.21)	0.75-1	-0.18 (-0.40,0.03)		-0.21 (-0.37,-0.05)	
	1-1.5	-0.14 (-0.33,0.05)		-0.15 (-0.30,-0.01)	
2+ 0.28 (0.03,0.54) 0.14 (-0.06,0.33)	1.5-2	0.08 (-0.15,0.30)		0.04 (-0.13,0.21)	
	2+	0.28 (0.03,0.54)		0.14 (-0.06,0.33)	

Table 2. continued

	Value		Quality	
Characteristic	Coeff (SE)	P-value	Coeff (SE)	P-value
% Duals		<0.001		<0.001
<=10%	ref		ref	
11%-20%	-0.62 (-0.92,-0.33)		-0.37 (-0.59,-0.15)	
21%-40%	-0.95 (-1.28,-0.63)		-0.79 (-1.03,-0.55)	
>40%	-1.83 (-2.26,-1.41)		-1.32 (-1.64,-1.00)	
Total Expenses		0.758		0.674
Quintile 1	ref		ref	
Quintile 2	-0.06 (-0.36,0.24)		-0.12 (-0.35,0.10)	
Quintile 3	-0.21 (-0.60,0.17)		-0.20 (-0.49,0.09)	
Quintile 4	-0.18 (-0.66,0.29)		-0.21 (-0.57,0.14)	
Quintile 5	-0.29 (-0.86,0.29)		-0.31 (-0.74,0.13)	
Supplies \$		0.073		0.014
Quintile 1	ref		ref	
Quintile 2	-0.00 (-0.31,0.30)		0.12 (-0.11,0.34)	
Quintile 3	-0.00 (-0.40,0.40)		0.06 (-0.24,0.37)	
Quintile 4	0.34 (-0.15,0.84)		0.38 (0.01,0.75)	
Quintile 5	0.51 (-0.08,1.11)		0.52 (0.07,0.97)	
Supplies/Total %		0.393		0.623
Quintile 1	ref		ref	
Quintile 2	-0.08 (-0.30,0.13)		0.04 (-0.12,0.20)	
Quintile 3	-0.22 (-0.46,0.02)		0.06 (-0.12,0.24)	
Quintile 4	-0.23 (-0.49,0.03)		0.03 (-0.17,0.23)	
Quintile 5	-0.22 (-0.52,0.08)		-0.07 (-0.29,0.16)	
% DISH		0.004		<0.001
Quintile 1	ref		ref	
Quintile 2	-0.18 (-0.39,0.04)		-0.27 (-0.43,-0.11)	
Quintile 3	-0.32 (-0.55,-0.09)		-0.40 (-0.57,-0.23)	
Quintile 4	-0.38 (-0.62,-0.14)		-0.36 (-0.54,-0.18)	
Quintile 5	-0.53 (-0.82,-0.24)		-0.64 (-0.86,-0.43)	

Source: Center for Medicare and Medicaid Services Overall Star Ratings score and Medicare Spending per Beneficiary; American Hospital Association Survey.

**Table 3** Results of multilogit model for low and high value versus average value hospitals.

	Low Value		High Value	
Characteristic	OR (95% CI)	P-value	OR (95% CI)	P-value
Ownership		< 0.001		0.002
Public	ref		ref	
Not for profit	0.3 (0.2,0.6)		2.2 (1.3,3.7)	
For profit	0.8 (0.4,1.4)		1.0 (0.5,2.0)	
Government	-		2.9 (0.4,19.3)	
Teaching status		0.012		0.615
Nonteaching	ref		ref	
Residency	2.9 (1.4,5.8)		1.1 (0.7,1.8)	
Teaching	2.4 (1.2,4.8)		1.3 (0.8,2.2)	
Beds		< 0.001		0.003
<100	ref		ref	
100-199	2.7 (1.2,6.1)		0.5 (0.3,0.7)	
200-299	5.8 (2.2,15.0)		0.4 (0.2,0.8)	
300-399	11.8 (4.2,33.1)		0.4 (0.1,0.8)	
400+	10.1 (3.3,30.9)		0.2 (0.1,0.5)	
Urbanity		0.292		0.694
Rural	ref		ref	
Urban	1.4 (0.8,2.5)		1.1 (0.7,1.6)	
Region <sup>1</sup>		< 0.001		< 0.001
West	ref		ref	
Midwest	0.5 (0.2,1.2)		0.4 (0.2,0.6)	
Northeast	2.7 (1.4,5.3)		0.2 (0.1,0.3)	
South	1.3 (0.7,2.4)		0.3 (0.2,0.4)	
Nurse/Beds		0.021		0.618
<=0.75	ref		ref	
0.75-1	1.0 (0.6,1.6)		1.0 (0.6,1.7)	
1-1.5	0.8 (0.5,1.3)		0.8 (0.5,1.3)	
1.5-2	0.4 (0.2,0.9)		1.2 (0.7,2.1)	
2+	0.3 (0.1,0.7)		1.1 (0.6,2.0)	

Table 3 Continued

% Duals		0.304		0.016
<=10%	ref		ref	
11%-20%	2.5 (0.7,8.8)		1.0 (0.5,1.8)	
21%-40%	3.0 (0.8,11.2)		1.5 (0.8,3.1)	
>40%	3.8 (0.9,15.7)		0.5 (0.2,1.6)	
Total Expenses		0.573		0.666
Quintile 1	ref		ref	
Quintile 2	1.6 (0.6,4.1)		1.4 (0.8,2.6)	
Quintile 3	1.1 (0.3,3.5)		1.1 (0.5,2.5)	
Quintile 4	1.1 (0.3,4.2)		1.2 (0.4,3.6)	
Quintile 5	0.7 (0.1,3.6)		1.0 (0.2,4.3)	
Supplies \$	, , ,	0.609	, , ,	0.819
Quintile 1	ref		ref	
Quintile 2	0.5 (0.2,1.3)		0.8 (0.4,1.4)	
Quintile 3	0.4 (0.1,1.2)		0.6 (0.2,1.4)	
Quintile 4	0.3 (0.1,1.3)		0.5 (0.2,1.7)	
Quintile 5	0.3 (0.1,1.5)		0.4 (0.1,2.2)	
Supplies/Total %		0.012		0.309
Quintile 1	ref		ref	
Quintile 2	0.9 (0.5,1.8)		0.9 (0.6,1.5)	
Quintile 3	1.9 (1.0,3.7)		1.2 (0.7,2.0)	
Quintile 4	1.3 (0.6,2.9)		0.7 (0.4,1.3)	
Quintile 5	2.8 (1.2,6.2)		1.2 (0.6,2.4)	
% DISH		0.050		0.011
Quintile 1	ref		ref	
Quintile 2	0.7 (0.4,1.3)		0.9 (0.5,1.4)	
Quintile 3	1.0 (0.5,1.9)		0.5 (0.3,0.9)	
Quintile 4	0.6 (0.3,1.3)		0.4 (0.2,0.7)	
Quintile 5	1.4 (0.7,3.1)		0.6 (0.3,1.1)	

Source: Center for Medicare and Medicaid Services Overall Star Ratings score and Medicare Spending per Beneficiary; American Hospital Association Survey.

# Notes:

1. Region groups were collapsed due to sparse cells under original groupings.

Figure 1. Medicare Spending per Beneificiary by Overall Star Rating

Figure 2. Value score versus Star Rating summary score.





Figure 1. Medicare Spending per Beneificiary by Overall Star Rating 139x101mm (96 x 96 DPI)




Figure 2. Value score versus Star Rating summary score. 139x101mm~(96~x~96~DPI)

# **BMJ Open**

# IDENTIFYING HIGH VALUE HOSPITAL CARE FOR MEDICARE BENEFICIARIES: AN OBSERVATIONAL STUDY

Journal:	BMJ Open
Manuscript ID	bmjopen-2021-053629.R1
Article Type:	Original research
Date Submitted by the Author:	18-Nov-2021
Complete List of Authors:	Herrin, Jeph; Yale University School of Medicine; Yale University School of Medicine, Section of Cardiovascular Medicine Yu, Huihui; Yale University Venkatesh, Arjun; Yale University, Department of Emergency Medicine Desai, Sunita; New York University Grossman School of Medicine, Department of Population Health Thiel, Cassandra; New York University Grossman School of Medicine, Department of Population Health Lin, Zhenqiu; Yale-New Haven Hospital, Center for Outcomes Research and Evaluation Bernheim, Susannah M; Yale University School of Medicine, Department of Internal Medicine Horwitz, Leora; New York University Grossman School of Medicine, Center for Healthcare Innovation and Delivery Science
<b>Primary Subject Heading</b> :	Health policy
Secondary Subject Heading:	Health services research
Keywords:	Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, STATISTICS & RESEARCH METHODS, Quality in health care < HEALTH SERVICES ADMINISTRATION & MANAGEMENT
	·

SCHOLARONE™ Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our licence.

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which Creative Commons licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

# IDENTIFYING HIGH VALUE HOSPITAL CARE FOR MEDICARE BENEFICIARIES: AN OBSERVATIONAL STUDY

Jeph Herrin, Ph.D.<sup>1</sup>

Huihui Yu, Ph.D.<sup>2</sup>

Arjun Venkatesh<sup>3</sup>

Sunita Desai4

Cassandra Thiel<sup>5</sup>

Zhenqui Lin<sup>1,6</sup>

Susannah Bernheim<sup>6,7</sup>

Leora I. Horwitz<sup>5,8,9</sup>

- 1. Section for Cardiovascular Medicine, Yale University School of Medicine, New Haven CT, USA
- 2. Yale University School of Medicine, New Haven CT, USA
- 3. Department of Emergency Medicine, Yale University School of Medicine, New Haven, CT, USA.
- 4. Department of Population Health, New York University Grossman School of Medicine, New York, NY, USA
- 5. Department of Population Health, NYU Grossman School of Medicine, New York, New York.
- 6. Center for Outcomes Research and Evaluation, Yale New Haven Health, New Haven, CT.
- 7. Section of General Internal Medicine, Department of Medicine, Yale School of Medicine, New Haven, CT
- 8. Center for Healthcare Innovation and Delivery Science, NYU Langone Health, New York, New York.
- 9. Department of Medicine, NYU Grossman School of Medicine, New York, New York.

# **Corresponding Author:**

Jeph Herrin

Jeph.herrin@yale.edu

Keywords: Healthcare Value, Quality of Care

Words: 3, 054

#### **ABSTRACT**

**Objectives** High value care is providing high quality care at low cost; we sought to define hospital value and identify the characteristics of hospitals which provide high value care.

**Design**: Retrospective observational study.

**Setting**: Acute care hospitals in the U.S.

**Participants**: All Medicare beneficiaries with claims included in Center for Medicare & Medicaid Services

Overall Star Ratings or in publicly available Medicare spending per beneficiary data.

Primary and secondary outcome measures: Our primary outcome was value defined as the difference between Star Ratings quality score and Medicare spending; the secondary outcome was classification as a 4 or 5 Star hospital with lowest quintile Medicare spending ("high value") or 1 or 2 Star hospital with highest quintile spending ("low value").

**Results**: 2,914 hospitals had both quality and spending data, and were included. ). The value score had a mean (SD) of 0.58 (1.79). A total of 286 hospitals were classified as high value; these represented 28.6% of 999 4 and 5-star hospitals and 46.8% of 611 low cost hospitals. A total of 258 hospitals were classified as low value; these represented 26.6% of 970 1 and 2-star hospitals and 49.3% of 523 high cost hospitals. In regression models ownership, non-teaching status, beds, urbanity, nurse to bed ratio, percentage of dual eligible Medicare patients, and smaller of disproportionate share hospital payments were associated with the primary value score.

**Conclusions**: There are high quality hospitals that are not high value, and a number of factors are strongly associated with being low or high value. These findings can inform efforts of policymakers and hospitals to increase the value of care.

### **ARTICLE SUMMARY**

# **Strengths and Limitations**

- This study incorporates a comprehensive measure of overall hospital quality derived from 5 distinct domains.
- This study uses payments as proxy for actual costs; however, these payments do include post discharge utilization.
- Two different measures of value are found to be consistent, and explained by factors that hospitals can modify.

# **Funding**

This work was funding by a grant from the Agency for Healthcare Quality and Research (R01HS022882).

#### INTRODUCTION

The past two decades have seen an increasing focus by United States policymakers and caregivers on improving hospital quality of care. The movement began in the early 2000s with public reporting of process measures, moved on to public reporting of outcomes measures such as readmissions, complications and mortality and, particularly since the passage of the Affordable Care Act in 2010, has since evolved to tying quality of care to payment.[1] There are now more than 100 quality measures on the CMS's Care Compare, and a summary of over 50 of these measures into an overall Hospital Quality Star Rating.[2] Patients and policy makers thus have available a range of quality metrics for comparing hospitals.

However, only recently have policymakers begun to take on the challenge of driving both improved quality and lower costs in tandem. The Value Based Purchasing Program (for hospitals) and the Quality Payment Program (for clinicians), for instance, both incorporate some measures of cost along with quality measures into their overall scoring system. Yet despite prior research indicating that people often perceive higher quality to be associated with higher costs and policymakers' concerns that cost reduction efforts will lower quality, the relationship between hospital quality and hospital cost, and in turn value, remains empirically underexplored. While in some cases high cost care is clearly of low quality (i.e., excess use of unnecessary care, higher complication rates), it may be that higher costs produce better outcomes in other situations. For example, higher spending may focus on high-value care interventions or services such as post-discharge planning or expensive, evidence based treatments such as percutaneous coronary intervention (PCIs) and implantable cardioverter defibrillators (ICDs).[3,4,5] Moreover, even if there is no systemic association between quality and cost, it is plausible that among hospitals that provide high quality care, some will be more cost effective at providing this care than others. Characteristics of hospitals that produce high value healthcare, therefore, as defined by quality accounting for cost, may not be the same as those that produce high quality alone.

We therefore sought to identify hospitals that provide 'high value' care and to examine how they differ from other hospitals, including specifically those which provide 'low value' care. Using the CMS Overall Hospital Quality Star Ratings to identify high and low quality care hospitals and the Medicare Spending per Beneficiary (MSPB) total cost of care scores to identify high and low cost hospitals, we created a value score as the difference between quality and cost scores. Using those hospitals that were particularly high or low value, we analyzed the characteristics of those hospitals, including anticipated drivers of costs (total expenses, supply costs) and factors historically associated with quality (beds, nurse staffing, teaching status, ownership, geographic region, and urban status).

### **METHODS**

Data and Cohort

We used publicly-reported Star Ratings data from Hospital Compare for 2018. The Star Rating system assigns each hospital an integer from 1 to 5, with 1 reflecting the lowest quality and 5 reflecting the highest quality. The Star Rating is a composite that reflects hospital performance across 57 measures, grouped into any of seven quality domains mortality, readmission, safety, patient experience, efficiency, effectiveness of care, and timeliness of care. Measures in each domain are used to estimate a latent variable model with a single latent variable; and the latent variables (group scores) for each domain are combined using a weighted average to generate an overall summary score. Finally, k-means clustering with 5 means is used to group the summary scores into five star categories.[6] A hospital was assigned an overall Star Rating if it reported at least 3 measures in at least 3 domains, one of which was safety, mortality or readmission.

Hospital-level Star Ratings data were linked to 2018 MSPB data using hospitals' CMS Certification numbers (CCNs). The MSPB score is calculated by the Center for Medicare & Medicaid Services (CMS) using Medicare Part A and Part B payments for all care provided from 3 days prior to, to 30 days

following an inpatient hospital stay (defined as an "episode" of care). MSPB episodes span all conditions, and MSPB scores are calculated for all Medicare fee-for-service hospital stays, excluding stays in psychiatric, rehabilitation, cancer, children's, critical access or long-stay hospitals; excluding hospitalizations resulting in transfers to another acute facility; and excluding readmissions that are within 30 days of an eligible hospitalization. Episodes in which the patient dies during the measurement period are also excluded, as are those with \$0 payments or involving transfers. The measure adjusts for the Medicare Severity-Diagnosis Related Group (MS-DRG) of the index hospitalization, age, use of long-term care, and 79 comorbidities. All payments are standardized across geographies for comparability; additional payments such as for indirect medical education or disproportionate share payments are omitted.[7] The MSPB scores is constructed as a ratio so that a value of 1 is "average" spending, and values lower or higher than 1 represent lower or higher spending than average.

Our study cohort comprised all hospitals which had an overall Star Rating; none were missing MSPB scores.

Value

We examined two related measures of value. For our first value outcome, we constructed a linear metric of value by subtracting the standardized MSPB score from the standardized Star Rating hospital summary score (used to create the 5 star categories). Though we initially considered a ratio of quality to spending, we were concerned that a ratio could allow hospitals to appear high value while having low quality, if their spending were low enough. Therefore, we used a difference measure, which ensures that hospitals categorized as high value were always of high quality. To facilitate interpretation, both spending and quality scores were standardized to range from 0 to 10; thus, higher scores indicate higher value, with the maximum value score being 10 and minimum -10.

For our second value outcome, we classified hospitals as 'high value' if they received a 4 or 5-star rating on Hospital Compare and had a MSPB score in the lowest quintile of all hospitals, and classified hospitals as 'low value' if they received a 1 or 2-star rating on Hospital Compare and had a MSPB score in the highest quintile of all hospitals; all other hospitals were classified as 'average' value. For this classification, we selected quintiles to categorize hospitals by their MSPB scores as providing meaningful cost distinctions while identifying adequate numbers of high and low value hospitals for analysis.

### Other Variables

Hospital characteristics were linked from the American Hospital Association (AHA) survey for 2018[8] and the CMS Provider of Services (POS) file of 2018. We examined hospital characteristics that have previously been associated with quality: geographic region (New England, Middle Atlantic, East North Central, West North Central, South Atlantic, East South Central, West South Central, Mountain, and Pacific); teaching status (teaching, residency program, non-teaching); number of beds(<100, 100-199,200-299,300-399, and 400+); nurse to bed ratio ( $\leq$ 0.75, 0.75-1, 1-1.5, >1.5); urban location (urban, rural); ownership status (for profit, private not for profit, public, federal); percentage of Medicare admissions that are for patients dually eligible for both Medicare and Medicaid ( $\leq$  10%, 11%-20%, 21%-40%, and > 40%); total expenses (quintiles); total cost of supplies (quintiles), ratio of supply expenses to total expenses (quintiles); and disproportionate share hospital (DSH) patient percentage (quintiles). DSH is a Medicare payment adjustment based on the proportion of Medicare admissions at a hospital that are for patients enrolled in Medicaid [9]

# Patient and Public Involvement

There was no patient or public involvement in the design, conduct or reporting of this study.

Statistical Analysis

We summarized value score (mean and standard deviation) and value category (frequency and percent) overall and by hospital characteristics, testing for differences in category characteristic using ANOVA for the value score and chi-square tests for the value category. To assess the overall relationship between cost and quality of care, we graphed the distribution of MSPB scores over each Star rating using a box plot (Figure 1), and then plotted the value scores against the Star Rating summary scores to visualize the association of value with quality (Figure 2). Then, to identify the hospital characteristics independently associated with value, and to identify contrasts in associations with quality, we estimated a series of models. We first used bivariate linear regression models to estimate the relationship between each hospital characteristics and value, where the outcome was our continuous value score; all factors were then included in a final multivariate model. For comparison with factors associated with quality alone, we estimated identical models where the dependent variable was the Star Ratings summary score, rather than our value score. We next estimated a multinomial logit model with the 3-category value as the outcome using 'average' as the reference group, including all factors. We used a multinomial rather than ordinal model to avoid assuming that associations were monotone; that is, we anticipated that some factors might be associated with low value and not high, and some with high value but not low. Both analyses included the same set of hospital characteristics, and all models included indicators for each domain, equal to 1 if the domain was included in the hospital's overall star rating score. We reported overall Wald test p-values for each characteristic. We report numbers and percent of missing values, and use multiple imputation (with 20 imputations) to account for missing values in all models.

All analyses were done using SAS 9.4 and Stata 16.1 (2020, StataCorp, College Station TX). All statistical tests were two-tailed, and we used p<0.05 to determine statistical significance. The study was approved by the Yale and NYU Institutional Review Boards.

# **RESULTS**

Our final study sample included 2,914 hospitals that received both a Star Rating and a MSPB score, all of which matched the AHA Survey file for 2018 (Table 1). The value score had a mean (SD) of 0.58 (1.79); see Figures 1 and 2. A total of 286 were classified as high value and 258 as low value by the categorical definition (Table 1). The 286 high value hospitals represented 28.6% of 999 4 and 5-star hospitals and 46.8% of 611 low cost (bottom quintile MSPB) hospitals; the 258 low value hospitals represented 26.6% of 970 1 and 2-star hospitals and 49.3% of 523 high cost hospitals.

In biviariate analyses (Table 1), we found not-for-profit private and government owned hospitals to have much higher values scores than for profit private hospitals, with score of 0.39 and 1.38 versus 0.08. Non-teaching hospitals and those with residency programs both had three times the value score of teaching hospitals, while hospitals with fewer than 100 beds (1.48) and less than 10% dual eligible patients (1.74) had the highest values scores. Hospitals in rural areas and smaller percentages of DSH payments also had higher value scores (all P-values < 0.05). For the categorical bivariate comparisons (Table 1), results were similar: high value hospitals were more likely to be not-for-profit, have no teaching program, have more beds and a higher nurse to bed ratio, and have a lower percentage of dual-eligible admissions compared to low- and average- value hospitals (all P values < 0.001). Regionally, lower value hospitals were more often in the Southern and Mid-Atlantic regions, while high value hospitals were in Northern Central and Pacific Regions (P<0.001).

In the multivariable model, all were significantly associated with value except those related to total and supply expenses (Table 2). The largest value effects were again for non-teaching, not-for-profit and government hospitals, hospitals with fewer than 100 beds and those with fewer than 10% dual eligible admissions. In equivalent model using the quality score instead of value as an outcome, there was no

association with urban status, while, unlike with value, higher supply expenses were associated with higher quality; other patterns of effects were very similar.

In multinomial regression assessing the associations between characteristics and low, average, and high value classification (Table 3), we collapsed the region categories to account for sparse cells. In the final model Government owned and not-for-profit hospitals, hospitals with fewer than 100 beds, hospitals in the Western region, and those with lower percentages of dual eligible patients and DSH patients were more likely to be high value than average value (all P-values < 0.05). Correspondingly, public hospitals, those with the more beds, those in the Northeast region, and lowest nurse to bed ratio all had higher odds of being low-value (all P-values < 0.05).

# **DISCUSSION**

In this examination of hospital value of care, we identified hospitals that provided relatively higher quality of care to Medicare beneficiaries at lower comparative costs, as well as hospitals that provided lower quality care at a high level of spending. In bivariate analyses, both high and low value hospitals differed from other hospitals strongly in ownership, urban location, bed size and region, nurse-to-bed ratio, amount spent on supplies relative to total expenses, and percentage of dual-eligible patients.

Most associations were significant in the multivariable primary analysis using the value score; of these, all but those related to urbanicity and expenses persisted in the multinomial logit analysis using a categorical outcome of low, average or high value.

The most notable finding, however, may not be the associations of hospital characteristics with value, but that value and quality of care are not synonymous: though generally, the MSPB declined with increasing Star ratings (Figure 1), less than a third of 4 and 5-star hospitals were high value, while more than a fourth of 1 and 2-star hospitals were lowest value. The observed trend in spending across overall quality (Figure 1) as well as the pattern of value classification is very similar to that found previously for

patient experience Star Ratings [10,11]; though the patient experience Star Rating is a component of the overall Star Rating, it accounts for only 22% of the weight of the overall summary score, so it is unlikely to drive the current findings. This finding was also reflected in our model of quality score, which found urban status was not related to quality while quality increased with more spending on supplies. That quality and value are not identical indicates that it will be possible to reduce spending among even the highest quality hospitals; however there is clearly a tension between the two, for if quality improvements driven by increased supply spending (or other additional resources) that translate to greater spending, then only offsets in other areas can increase value.

Most of the associations we found for value are similar to those seen for hospital characteristics and quality, both here with the Star Ratings summary score and in prior research; for example, Delancy et. al found that hospitals with fewer beds, no medical school affiliation, and lower proportions of DSH patients were more likely to have higher Star ratings [12]. This was similar to the results of our analysis of the star summary quality score (Table 2). However, in our categorical analysis, we found no relationship between teaching status and high value. Geographic region also explained some of the difference between low value and average value hospitals across all of our analyses, with hospitals in East South Central and Mid-Atlantic regions having lowest value and being more likely to be low value than those in any other regions, and Pacific region hospitals being consistently high value in all models. Since our spending measure is adjusted for local wage and cost variations, this geographic variation is likely driven substantially by variations in quality.

Of most relevance to hospitals and policy makers are the associations with factors that are within the control of the hospital. A key factor that was related to low value in all analyses was nurse to bed ratio, with lower ratios being more often found at the lowest value hospitals – only 9 (3.5%) low value hospitals had a ratio of 2 or more, while 43 (15.0%) of high value hospitals had a ratio of at least 2. This finding suggests that high levels of nurse staffing, while increasing the expense of inpatient care, may

reduce overall Medicare payments while promoting high quality care. Importantly, this finding also highlights a perennial challenge for policymakers: the wrong pocket problem. Since the costs of increased staffing are borne by hospitals but the benefits of lower spending are reaped by payers and patients, there is no intrinsic incentive for hospitals to pursue such interventions. Payment policies seeking to improve value should explicitly take these challenges into account.

Conversely, 66 (25.6%) of low value hospitals reported at least 40% of their Medicare patients were also eligible for Medicaid, while only 11 (3.8%) of high value hospitals reported such high rates of dual eligibility, with proportion of DSH patients exhibiting a similar but weaker pattern. This may indicate that lower income patients have unmeasured comorbidities or care complexities that are not captured by the risk adjustment used in Star Ratings' underlying measures or that these patients drive costs in ways unrelated to quality of care (e.g., limited social support may lead to greater post-discharge costs), that these patients may have higher cost needs that are not completely adjusted for, or possibly that DSH patients are more likely to receive care at lower quality hospitals.

This study has the limitations of any observational study, including that no causal inferences can be drawn; for example, it may be that lower value hospitals have increased costs of care because of lower quality, rather than independent of it. In addition, we have only examined a few key factors that may be associated with quality, in order to better focus on value of care. And, we have focused only on quality and spending metrics largely related to Medicare fee-for-service beneficiaries; other, broader measures of quality and costs that included the entire patient population could plausibly produce different results. Moreover, Medicare payments are distorted somewhat by the Inpatient Prospective Payment System. However, CMS Overall Star Ratings captures quality of care across 7 domains, and the MSPB is a measure of spending with cost-of-living adjustments that plausibly represents the overall spending per patient resulting from a hospitalization.

It is worth considering that the MSPB does not include admissions that lead to transfers, nor admissions in which the patient does not survive 30 days. Both exclusions are likely to confound the relationship between the spending score and the mortality domain, which accounts for 22% of the Star Rating summary score, since publicly reported mortality measures typically measure outcomes within the same 30 days, and attribute them to the first hospital in the event of transfers. Excluding patients who do not survive and/or are transferred likely reduces the MSPB score (assuming these are more complex patients), and more so for hospitals with higher mortality or transfer rates. However, this would tend to bias our current findings toward the null, attenuating the relationship between spending value.

# **CONCLUSION**

We have shown that care quality and care value are not identical; there are high quality hospitals that are not high value. In addition, there are a number of structural factors, some under the control of hospitals such as nurse to bed ratio, that are strongly associated with being low or high value. These findings can inform efforts of policymakers and hospitals to increase the value of care.

#### **Contributions**

Drs Herrin, Venkatesh, Bernheim and Horwitz contributed to the conception and design of the study. Drs Herrin and Yu performed all analyses. Dr Herrin drafted the manuscript. Drs Venkatesh, Desai, Thiel, Lin, Bernheim and Horwitz contributed critical additions and conceptual revisions to the manuscript.

# **Competing Interests**

Drs. Bernheim, Herrin, Lin and Venkatesh recieve salary support from the Centers for Medicare and Medicaid Services to develop, implement, and maintain hospital performance outcome measures, including the methodology for the Overall Hospital Star Ratings, that are publicly reported. Drs. Horwitz and Yu have worked under contract to the Centers for Medicare and Medicaid Services to develop quality measures, including some used in the Overall Hospital Star Ratings program.

# **Funding**

This work was funded by the Agency for Healthcare Research and Quality, grant R01HS022882.

# **Data Sharing**

Star Ratings and Medicare spending data are publicly available, and an analytic file including those data can be provided by the authors. Data collected by the American Hospital Association are proprietary, and will not be available for open access sharing.

# **Ethics Approval**

This study was approved by the Institutional Review Boards of Yale University (ID 0903004927) and New York University (ID S19-01460).

#### REFERENCES

- U.S. Centers for Medicare & Medicaid Services. Hospital Readmissions Reduction Program (HRRP).
   Accessed 2020 Dec 08. <a href="https://www.cms.gov/medicare/medicare-fee-for-service-payment/acuteinpatientpps/readmissions-reduction-program.html">https://www.cms.gov/medicare/medicare-fee-for-service-payment/acuteinpatientpps/readmissions-reduction-program.html</a>
- U.S. Centers for Medicare and Medicaid Services. Hospital Compare Overall Ratings. Accessed 2020
   Dec 08. https://qualitynet.cms.gov/inpatient/public-reporting/overall-ratings
- Doyle JJ, Graves JA, Gruber JA, Kleiner S. Do high-cost hospitals deliver better care? Evidence from ambulance referral patterns. National Bureau of Economic Research. http://www.nber.org/papers/w17936.pdf?new\_window=1. Accessed December 19, 2013.
- 4. Silber JH, Kaestner R, Even-Shoshan O, Wang Y, Bressler LJ. Aggressive treatment style and surgical outcomes. Health Serv Res. 2010;45(6 Pt 2):1872-1892.
- 5. Virtual Mentor. 2014;16(2):124-130.
- 6. QualityNet. https://www.qualitynet.org/inpatient/public-reporting/overall-ratings/resources
- 7. QualityNet. https://www.qualitynet.org/inpatient/measures/mspb/methodology
- American Hospital Association. American Hospital Association (AHA) Annual Survey Database –
   2018.
- 9. https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/dsh
- Liao JM, Zhou L, Navathe AS. Nationwide Hospital Performance on Publicly Reported Episode
   Spending Measures. J Hosp Med. 2020 Mar 18;15:E1-E7. doi: 10.12788/jhm.3377. Epub ahead of print. PMID: 32195657
- 11. Trzeciak S, Gaughan JP, Bosire J, Angelo M, Holzberg AS, Mazzarelli AJ. Association Between Medicare Star Ratings for Patient Experience and Medicare Spending per Beneficiary for US Hospitals. J Patient Exp. 2017 Mar;4(1):17-21. doi: 10.1177/2374373516685938. Epub 2017 Jan 12. PMID: 28725856; PMCID: PMC5513662

12. DeLancey JO, Softcheck J, Chung JW, Barnard C, Dahlke AR, Bilimoria KY. Associations Between Hospital Characteristics, Measure Reporting, and the Centers for Medicare & Medicaid Services Overall Hospital Quality Star Ratings. JAMA. 2017 May 16;317(19):2015-2017. doi: 10.1001/jama.2017.3148. PMID: 28510670; PMCID: PMC5815009.



**Table 1**. Characteristics of low, average, and high value hospitals.

	Value	0,	,			
Characteristic	Score		Low Value	Average	High Value	P-value
	Mean (SD)	P-value <sup>1</sup>	n(%) or	n(%) or	n(%) or	
			mean(SD)	mean(SD)	mean(SD)	
N	0.58 (1.79)		258 (100.0)	2370 (100.0)	286 (100.0)	
Star Rating		<0.001				
1	-2.38 (1.26)		89 (34.5)	179 (7.6)	0 (0.0)	
2	-0.66 (1.02)		169 (65.5)	533 (22.5)	0 (0.0)	
3	0.71 (1.00)		0 (0.0)	945 (39.9)	0 (0.0)	
4	1.75 (1.00)		0 (0.0)	528 (22.3)	213 (74.5)	
5	3.16 (1.07)		0 (0.0)	185 (7.8)	73 (25.5)	
Ownership		< 0.001				< 0.001
Public	0.39 (1.80)		43 (16.7)	356 (15.0)	33 (11.5)	
Not for profit	0.77 (1.69)		128 (49.6)	1521 (64.2)	215 (75.2)	
For profit	0.08 (1.96)		87 (33.7)	478 (20.2)	36 (12.6)	
Government	1.38 (1.60)		0 (0.0)	15 (0.6)	2 (0.7)	
Teaching status		< 0.001				< 0.001
Nonteaching	0.87 (1.72)		22 (8.5)	381 (16.1)	48 (16.8)	
Residency	0.72 (1.75)		124 (48.1)	1130 (47.7)	168 (58.7)	
Teaching	0.25 (1.82)		112 (43.4)	859 (36.2)	70 (24.5)	
Beds		< 0.001				< 0.001
<100	1.48 (1.57)		21 (8.1)	575 (24.3)	155 (54.2)	
100-199	0.65 (1.65)		58 (22.5)	645 (27.2)	67 (23.4)	
200-299	0.21 (1.73)		54 (20.9)	399 (16.8)	33 (11.5)	
300-399	0.08 (1.86)		58 (22.5)	274 (11.6)	17 (5.9)	
400+	-0.12 (1.74)		67 (26.0)	477 (20.1)	14 (4.9)	
Urbanity	, ,	< 0.001	, ,	, ,	, ,	< 0.001
Rural	1.08 (1.52)		29 (11.2)	579 (24.4)	110 (38.5)	
Urban	0.41 (1.84)		229 (88.8)	1791 (75.6)	176 (61.5)	
Region	` ,	< 0.001	, ,	,	, ,	< 0.001
New England	0.70 (1.48)		3 (1.2)	115 (4.9)	9 (3.1)	
Mid Atlantic	-0.27 (1.92)		62 (24.0)	260 (11.0)	13 (4.5)	
E North Central	1.02 (1.58)		22 (8.5)	382 (16.1)	57 (19.9)	
W North Central	1.60 (1.48)		1 (0.4)	179 (7.6)	50 (17.5)	
South Atlantic	0.40 (1.71)		43 (16.7)	411 (17.3)	42 (14.7)	
E South Central	0.15 (1.46)		23 (8.9)	236 (10.0)	11 (3.8)	
W South Central	0.17 (1.80)		60 (23.3)	341 (14.4)	20 (7.0)	
Mountain	1.13 (1.78)		12 (4.7)	155 (6.5)	31 (10.8)	
Pacific	0.81 (1.91)		32 (12.4)	291 (12.3)	53 (18.5)	
Nurse/Beds	0.01 (1.51)	< 0.001	32 (12.4)	231 (12.3)	33 (18.3)	< 0.001
<=0.75	0.51 (1.88)	<b>\0.001</b>	86 (33.3)	536 (22.6)	79 (27.6)	<b>\0.001</b>
0.75-1	0.35 (1.73)		52 (20.2)	405 (17.1)	47 (16.4)	
1-1.5	0.33 (1.73)		86 (33.3)	783 (33.0)	65 (22.7)	
1.5-2	0.41 (1.72)		25 (9.7)	381 (16.1)	52 (18.2)	
1.5-2 2+	1.24 (1.67)		9 (3.5)	265 (11.2)	43 (15.0)	
<b>4</b> T	1.24 (1.07)		3 (3.3)	203 (11.2)	43 (13.0)	

Table 1 continued

% Duals		< 0.001				< 0.001
<=10%	1.74 (1.93)		7 (2.7)	176 (7.4)	33 (11.5)	
11%-20%	0.80 (1.58)		67 (26.0)	863 (36.4)	91 (31.8)	
21%-40%	0.51 (1.71)		118 (45.7)	1057 (44.6)	151 (52.8)	
>40%	-0.56 (1.94)		66 (25.6)	274 (11.6)	11 (3.8)	
Total Expenses		< 0.001				< 0.001
Quintile 1	1.09 (1.66)		28 (10.9)	459 (19.4)	95 (33.2)	
Quintile 2	0.79 (1.77)		54 (20.9)	446 (18.8)	81 (28.3)	
Quintile 3	0.44 (1.73)		57 (22.1)	475 (20.0)	50 (17.5)	
Quintile 4	0.31 (1.81)		71 (27.5)	473 (20.0)	37 (12.9)	
Quintile 5	0.24 (1.83)		48 (18.6)	511 (21.6)	22 (7.7)	
Missing	0.68 (1.61)		0 (0.0)	6 (0.3)	1 (0.3)	
Supplies \$		< 0.001				< 0.001
Quintile 1	1.11 (1.63)		23 (8.9)	301 (12.7)	72 (25.2)	
Quintile 2	0.83 (1.64)		31 (12.0)	311 (13.1)	53 (18.5)	
Quintile 3	0.50 (1.82)		35 (13.6)	326 (13.8)	35 (12.2)	
Quintile 4	0.59 (1.72)		37 (14.3)	334 (14.1)	24 (8.4)	
Quintile 5	0.27 (1.81)		34 (13.2)	349 (14.7)	12 (4.2)	
Missing	0.40 (1.86)		98 (38.0)	749 (31.6)	90 (31.5)	
Supplies/Total %		0.042				0.002
Quintile 1	0.83 (1.79)		29 (11.2)	311 (13.1)	56 (19.6)	
Quintile 2	0.75 (1.69)		23 (8.9)	329 (13.9)	43 (15.0)	
Quintile 3	0.51 (1.75)		38 (14.7)	317 (13.4)	41 (14.3)	
Quintile 4	0.53 (1.56)		28 (10.9)	344 (14.5)	23 (8.0)	
Quintile 5	0.68 (1.92)		42 (16.3)	320 (13.5)	33 (11.5)	
Missing	0.40 (1.86)		98 (38.0)	749 (31.6)	90 (31.5)	
% DSH		< 0.001				< 0.001
Quintile 1	1.35 (1.74)		32 (12.4)	468 (19.7)	84 (29.4)	
Quintile 2	0.85 (1.62)		33 (12.8)	480 (20.3)	70 (24.5)	
Quintile 3	0.63 (1.59)		42 (16.3)	488 (20.6)	52 (18.2)	
Quintile 4	0.30 (1.60)		55 (21.3)	484 (20.4)	44 (15.4)	
Quintile 5	-0.26 (1.96)		96 (37.2)	450 (19.0)	36 (12.6)	
	• •		· · · · · ·		· · · · · · · · · · · · · · · · · · ·	

Source: Center for Medicare and Medicaid Services Overall Star Ratings score and Medicare Spending per Beneficiary; American Hospital Association Survey.

# Notes

1. P-value based on ANOVA model estimated using multiple imputation for missing values.

**Table 2**. Results of multivariable linear models with value score and Star Ratings summary score as dependent variables.

	Value		Quality	
Characteristic	Coeff (SE)	P-value		
Ownership		<0.001		<0.001
Public	ref		ref	
Not for profit	0.40 (0.21,0.59)		0.34 (0.20,0.48)	
For profit	-0.04 (-0.29,0.21)		0.14 (-0.05,0.33)	
Government	0.96 (-0.23,2.15)		0.47 (-0.42,1.36)	
Teaching status		0.008		0.006
Nonteaching	ref		ref	
Residency	-0.30 (-0.50,-0.10)		-0.19 (-0.34,-0.05)	
Teaching	-0.27 (-0.47,-0.07)		-0.24 (-0.39,-0.09)	
Beds		<0.001		<0.001
<100	ref		ref	
100-199	-0.55 (-0.77,-0.33)		-0.31 (-0.47,-0.14)	
200-299	-0.99 (-1.29,-0.70)		-0.57 (-0.79,-0.36)	
300-399	-1.24 (-1.57,-0.90)		-0.67 (-0.93,-0.42)	
400+	-1.45 (-1.81,-1.08)		-0.80 (-1.07,-0.52)	
Urbanity		0.015		0.894
Rural	ref		ref	
Urban	-0.23 (-0.42,-0.04)		-0.01 (-0.15,0.13)	
Region		< 0.001		<0.001
New England	ref		ref	
Mid Atlantic	-0.67 (-1.03,-0.31)		-0.56 (-0.83,-0.29)	
E North Central	0.35 (0.01,0.70)		0.27 (0.02,0.53)	
W North Central	0.57 (0.20,0.95)		0.03 (-0.25,0.31)	
South Atlantic	-0.09 (-0.45,0.27)		-0.25 (-0.52,0.02)	
E South Central	-0.70 (-1.13,-0.28)		-0.55 (-0.86,-0.23)	
W South Central	-0.62 (-0.99,-0.26)		-0.15 (-0.42,0.12)	
Mountain	0.23 (-0.19,0.65)		-0.02 (-0.34,0.29)	
Pacific	0.83 (0.45,1.22)		0.35 (0.06,0.64)	
Nurse/Beds		0.006		0.001
<=0.75	ref		ref	
0.75-1	-0.19 (-0.40,0.03)		-0.21 (-0.37,-0.05)	
1-1.5	-0.14 (-0.34,0.05)		-0.16 (-0.30,-0.01)	
1.5-2	0.05 (-0.18,0.28)		0.03 (-0.14,0.20)	
2+	0.22 (-0.04,0.48)		0.10 (-0.10,0.29)	

Table 2. continued

	Value		Quality	
Characteristic	Coeff (SE)	P-value	Coeff (SE)	P-value
% Duals		<0.001		<0.001
<=10%	ref		ref	
11%-20%	-0.48 (-0.78,-0.18)		-0.26 (-0.48,-0.03)	
21%-40%	-0.81 (-1.14,-0.48)		-0.68 (-0.93,-0.43)	
>40%	-1.66 (-2.10,-1.23)		-1.20 (-1.53,-0.88)	
Total Expenses		0.821		0.790
Quintile 1	ref		ref	
Quintile 2	-0.01 (-0.31,0.29)		-0.09 (-0.32,0.13)	
Quintile 3	-0.15 (-0.54,0.24)		-0.15 (-0.44,0.14)	
Quintile 4	-0.12 (-0.60,0.35)		-0.17 (-0.52,0.18)	
Quintile 5	-0.24 (-0.81,0.34)		-0.27 (-0.71,0.16)	
Supplies \$		0.034		0.008
Quintile 1	ref		ref	
Quintile 2	-0.06 (-0.37,0.24)		0.07 (-0.16,0.30)	
Quintile 3	-0.06 (-0.47,0.34)		0.01 (-0.29,0.31)	
Quintile 4	0.30 (-0.20,0.79)		0.34 (-0.03,0.71)	
Quintile 5	0.50 (-0.09,1.10)		0.50 (0.05,0.94)	
Supplies/Total %		0.424		0.479
Quintile 1	ref		ref	
Quintile 2	-0.10 (-0.32,0.12)		0.03 (-0.14,0.19)	
Quintile 3	-0.21 (-0.45,0.03)		0.06 (-0.12,0.24)	
Quintile 4	-0.23 (-0.49,0.03)		0.02 (-0.17,0.22)	
Quintile 5	-0.24 (-0.54,0.06)		-0.09 (-0.31,0.13)	
% DISH		0.003		< 0.001
Quintile 1	ref		ref	
Quintile 2	-0.17 (-0.39,0.04)		-0.27 (-0.43,-0.10)	
Quintile 3	-0.32 (-0.55,-0.09)		-0.40 (-0.57,-0.23)	
Quintile 4	-0.38 (-0.63,-0.14)		-0.37 (-0.55,-0.18)	
Quintile 5	-0.56 (-0.85,-0.27)		-0.66 (-0.88,-0.44)	

Source: Center for Medicare and Medicaid Services Overall Star Ratings score and Medicare Spending per Beneficiary; American Hospital Association Survey.

**Table 3** Results of multilogit model for low and high value versus average value hospitals.

	Low Value		High Value	
Characteristic	OR (95% CI)	P-value	OR (95% CI)	P-value
Ownership		<0.001		0.002
Public	ref		ref	
Not for profit	0.4 (0.2,0.6)		2.1 (1.2,3.5)	
For profit	0.8 (0.5,1.5)		0.8 (0.4,1.8)	
Government	0.0 (0.0,.)		2.0 (0.3,13.3)	
Teaching status		0.012		0.698
Nonteaching	ref		ref	
Residency	2.9 (1.4,5.8)		1.1 (0.7,1.7)	
Teaching	2.4 (1.2,4.9)		1.2 (0.7,2.1)	
Beds		<0.001		0.003
<100	ref		ref	
100-199	2.9 (1.2,6.8)		0.5 (0.3,0.7)	
200-299	5.9 (2.2,16.1)		0.4 (0.2,0.8)	
300-399	11.9 (4.1,34.5)		0.3 (0.1,0.8)	
400+	10.0 (3.2,31.7)		0.2 (0.1,0.5)	
Urbanity		0.300		0.758
Rural	ref		ref	
Urban	1.4 (0.8,2.5)		1.1 (0.7,1.6)	
Region <sup>1</sup>		<0.001		< 0.001
West	ref		ref	
Midwest	0.5 (0.2,1.2)		0.4 (0.2,0.6)	
Northeast	2.7 (1.4,5.3)		0.1 (0.1,0.3)	
South	1.3 (0.7,2.4)		0.3 (0.2,0.4)	
Nurse/Beds		0.025		0.673
<=0.75	ref		ref	
0.75-1	1.0 (0.6,1.7)		1.0 (0.6,1.7)	
1-1.5	0.8 (0.5,1.3)		0.8 (0.5,1.2)	
1.5-2	0.5 (0.2,0.9)		1.1 (0.6,1.9)	
2+	0.3 (0.1,0.7)		1.0 (0.5,1.8)	

Table 3 Continued

% Duals		0.427		0.009
<=10%	ref		ref	
11%-20%	2.2 (0.6,7.8)		1.1 (0.6,2.2)	
21%-40%	2.7 (0.7,9.8)		1.9 (0.9,4.0)	
>40%	3.2 (0.8,13.2)		0.6 (0.2,2.0)	
Total Expenses	, , ,	0.536	, , ,	0.626
Quintile 1	ref		ref	
Quintile 2	1.7 (0.7,4.6)		1.4 (0.8,2.6)	
Quintile 3	1.2 (0.4,4.0)		1.1 (0.5,2.4)	
Quintile 4	1.3 (0.3,5.1)		1.2 (0.4,3.5)	
Quintile 5	0.9 (0.2,4.4)		0.9 (0.2,4.1)	
Supplies \$		0.660		0.845
Quintile 1	ref		ref	
Quintile 2	0.6 (0.2,1.4)		0.7 (0.4,1.4)	
Quintile 3	0.4 (0.1,1.3)		0.6 (0.2,1.4)	
Quintile 4	0.4 (0.1,1.4)		0.5 (0.2,1.7)	
Quintile 5	0.3 (0.1,1.5)		0.5 (0.1,2.6)	
Supplies/Total %		0.010		0.273
Quintile 1	ref		ref	
Quintile 2	1.0 (0.5,2.0)		0.9 (0.5,1.4)	
Quintile 3	2.0 (1.0,4.0)		1.2 (0.7,2.1)	
Quintile 4	1.4 (0.7,3.1)		0.6 (0.3,1.3)	
Quintile 5	3.0 (1.3,6.8)		1.1 (0.5,2.3)	
% DISH		0.048		0.006
Quintile 1	ref		ref	
Quintile 2	0.7 (0.4,1.4)		0.8 (0.5,1.4)	
Quintile 3	1.1 (0.6,2.0)		0.5 (0.3,0.9)	
Quintile 4	0.6 (0.3,1.3)		0.4 (0.2,0.7)	
Quintile 5	1.6 (0.7,3.3)		0.5 (0.2,1.0)	

Source: Center for Medicare and Medicaid Services Overall Star Ratings score and Medicare Spending per Beneficiary; American Hospital Association Survey.

# Notes:

1. Region groups were collapsed due to sparse cells under original groupings.

Figure 1. Medicare Spending per Beneificiary by Overall Star Rating

Figure 2. Value score versus Star Rating summary score.





Figure 1. Medicare Spending per Beneificiary by Overall Star Rating  $139 \times 101 \text{mm} \ (300 \times 300 \ \text{DPI})$ 




Figure 2. Value score versus Star Rating summary score.  $139x101mm (300 \times 300 DPI)$ 

# **BMJ Open**

# IDENTIFYING HIGH VALUE CARE FOR MEDICARE BENEFICIARIES: A CROSS-SECTIONAL STUDY OF ACUTE CARE HOSPITALS IN THE U.S.

Journal:	BMJ Open
Manuscript ID	bmjopen-2021-053629.R2
Article Type:	Original research
Date Submitted by the Author:	09-Feb-2022
Complete List of Authors:	Herrin, Jeph; Yale University School of Medicine; Yale University School of Medicine, Section of Cardiovascular Medicine Yu, Huihui; Yale University Venkatesh, Arjun; Yale University, Department of Emergency Medicine Desai, Sunita; New York University Grossman School of Medicine, Department of Population Health Thiel, Cassandra; New York University Grossman School of Medicine, Department of Population Health Lin, Zhenqiu; Yale-New Haven Hospital, Center for Outcomes Research and Evaluation Bernheim, Susannah M; Yale University School of Medicine, Department of Internal Medicine Horwitz, Leora; New York University Grossman School of Medicine, Center for Healthcare Innovation and Delivery Science
 b>Primary Subject Heading:	Health policy
Secondary Subject Heading:	Health services research
Keywords:	Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, STATISTICS & RESEARCH METHODS, Quality in health care < HEALTH SERVICES ADMINISTRATION & MANAGEMENT

SCHOLARONE™ Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our licence.

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which Creative Commons licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

# IDENTIFYING HIGH VALUE CARE FOR MEDICARE BENEFICIARIES: A CROSS-SECTIONAL STUDY OF ACUTE CARE HOSPITALS IN THE U.S.

Jeph Herrin, Ph.D.<sup>1</sup>

Huihui Yu, Ph.D.<sup>2</sup>

Arjun Venkatesh<sup>3</sup>

Sunita Desai4

Cassandra Thiel<sup>5</sup>

Zhenqui Lin<sup>1,6</sup>

Susannah Bernheim<sup>6,7</sup>

Leora I. Horwitz<sup>5,8,9</sup>

- 1. Section for Cardiovascular Medicine, Yale University School of Medicine, New Haven CT, USA
- 2. Yale University School of Medicine, New Haven CT, USA
- 3. Department of Emergency Medicine, Yale University School of Medicine, New Haven, CT, USA.
- 4. Department of Population Health, New York University Grossman School of Medicine, New York, NY, USA
- 5. Department of Population Health, NYU Grossman School of Medicine, New York, New York.
- 6. Center for Outcomes Research and Evaluation, Yale New Haven Health, New Haven, CT.
- Section of General Internal Medicine, Department of Medicine, Yale School of Medicine, New Haven, CT
- 8. Center for Healthcare Innovation and Delivery Science, NYU Langone Health, New York, New York
- 9. Department of Medicine, NYU Grossman School of Medicine, New York, New York.

# **Corresponding Author:**

Jeph Herrin

Jeph.herrin@yale.edu

**Keywords**: Healthcare Value, Quality of Care

Words: 3, 054

#### **ABSTRACT**

**Objectives** High value care is providing high quality care at low cost; we sought to define hospital value and identify the characteristics of hospitals which provide high value care.

**Design**: Retrospective observational study.

Setting: Acute care hospitals in the U.S.

**Participants**: All Medicare beneficiaries with claims included in Center for Medicare & Medicaid Services

Overall Star Ratings or in publicly available Medicare spending per beneficiary data.

Primary and secondary outcome measures: Our primary outcome was value defined as the difference between Star Ratings quality score and Medicare spending; the secondary outcome was classification as a 4 or 5 Star hospital with lowest quintile Medicare spending ("high value") or 1 or 2 Star hospital with highest quintile spending ("low value").

**Results**: 2,914 hospitals had both quality and spending data, and were included. ). The value score had a mean (SD) of 0.58 (1.79). A total of 286 hospitals were classified as high value; these represented 28.6% of 999 4 and 5-star hospitals and 46.8% of 611 low cost hospitals. A total of 258 hospitals were classified as low value; these represented 26.6% of 970 1 and 2-star hospitals and 49.3% of 523 high cost hospitals. In regression models ownership, non-teaching status, beds, urbanity, nurse to bed ratio, percentage of dual eligible Medicare patients, and smaller of disproportionate share hospital payments were associated with the primary value score.

**Conclusions**: There are high quality hospitals that are not high value, and a number of factors are strongly associated with being low or high value. These findings can inform efforts of policymakers and hospitals to increase the value of care.

#### **ARTICLE SUMMARY**

# **Strengths and Limitations**

- This study incorporates a comprehensive measure of overall hospital quality derived from 5 distinct domains.
- This study uses payments as proxy for actual costs; however, these payments do include post discharge utilization.
- The quality and payment measures include only patients enrolled in a single insurance program,
   Medicare Fee for Service, and excludes most patients under 65 years of age.

# **Funding**

This work was funding by a grant from the Agency for Healthcare Quality and Research (R01HS022882).

#### INTRODUCTION

The past two decades have seen an increasing focus by United States policymakers and caregivers on improving hospital quality of care. The movement began in the early 2000s with public reporting of process measures, moved on to public reporting of outcomes measures such as readmissions, complications and mortality and, particularly since the passage of the Affordable Care Act in 2010, has since evolved to tying quality of care to payment.[1] There are now more than 100 quality measures on the CMS's Care Compare, and a summary of over 50 of these measures into an overall Hospital Quality Star Rating.[2] Patients and policy makers thus have available a range of quality metrics for comparing hospitals.

However, only recently have policymakers begun to take on the challenge of driving both improved quality and lower costs in tandem. The Value Based Purchasing Program (for hospitals) and the Quality Payment Program (for clinicians), for instance, both incorporate some measures of cost along with quality measures into their overall scoring system. Yet despite prior research indicating that people often perceive higher quality to be associated with higher costs and policymakers' concerns that cost reduction efforts will lower quality, the relationship between hospital quality and hospital cost, and in turn value, remains empirically underexplored. While in some cases high cost care is clearly of low quality (i.e., excess use of unnecessary care, higher complication rates), it may be that higher costs produce better outcomes in other situations. For example, higher spending may focus on high-value care interventions or services such as post-discharge planning or expensive, evidence based treatments such as percutaneous coronary intervention (PCIs) and implantable cardioverter defibrillators (ICDs).[3,4,5] Moreover, even if there is no systemic association between quality and cost, it is plausible that among hospitals that provide high quality care, some will be more cost effective at providing this care than others. Characteristics of hospitals that produce high value healthcare, therefore, as defined by quality accounting for cost, may not be the same as those that produce high quality alone.

We therefore sought to identify hospitals that provide 'high value' care and to examine how they differ from other hospitals, including specifically those which provide 'low value' care. Using the CMS Overall Hospital Quality Star Ratings to identify high and low quality care hospitals and the Medicare Spending per Beneficiary (MSPB) total cost of care scores to identify high and low cost hospitals, we created a value score as the difference between quality and cost scores. Using those hospitals that were particularly high or low value, we analyzed the characteristics of those hospitals, including anticipated drivers of costs (total expenses, supply costs) and factors historically associated with quality (beds, nurse staffing, teaching status, ownership, geographic region, and urban status).

#### **METHODS**

Data and Cohort

We used publicly-reported Star Ratings data from Hospital Compare for 2018. The Star Rating system assigns each hospital an integer from 1 to 5, with 1 reflecting the lowest quality and 5 reflecting the highest quality. The Star Rating is a composite that reflects hospital performance across 57 measures, grouped into any of seven quality domains mortality, readmission, safety, patient experience, efficiency, effectiveness of care, and timeliness of care. Measures in each domain are used to estimate a latent variable model with a single latent variable; and the latent variables (group scores) for each domain are combined using a weighted average to generate an overall summary score. Finally, k-means clustering with 5 means is used to group the summary scores into five star categories.[6] A hospital was assigned an overall Star Rating if it reported at least 3 measures in at least 3 domains, one of which was safety, mortality or readmission.

Hospital-level Star Ratings data were linked to 2018 MSPB data using hospitals' CMS Certification numbers (CCNs). The MSPB score is calculated by the Center for Medicare & Medicaid Services (CMS) using Medicare Part A and Part B payments for all care provided from 3 days prior to, to 30 days

following an inpatient hospital stay (defined as an "episode" of care). MSPB episodes span all conditions, and MSPB scores are calculated for all Medicare fee-for-service hospital stays, excluding stays in psychiatric, rehabilitation, cancer, children's, critical access or long-stay hospitals; excluding hospitalizations resulting in transfers to another acute facility; and excluding readmissions that are within 30 days of an eligible hospitalization. Episodes in which the patient dies during the measurement period are also excluded, as are those with \$0 payments or involving transfers. The measure adjusts for the Medicare Severity-Diagnosis Related Group (MS-DRG) of the index hospitalization, age, use of long-term care, and 79 comorbidities. All payments are standardized across geographies for comparability; additional payments such as for indirect medical education or disproportionate share payments are omitted.[7] The MSPB scores is constructed as a ratio so that a value of 1 is "average" spending, and values lower or higher than 1 represent lower or higher spending than average.

Our study cohort comprised all hospitals which had an overall Star Rating; none were missing MSPB scores.

Value

We examined two related measures of value. For our first value outcome, we constructed a linear metric of value by subtracting the standardized MSPB score from the standardized Star Rating hospital summary score (used to create the 5 star categories). Though we initially considered a ratio of quality to spending, we were concerned that a ratio could allow hospitals to appear high value while having low quality, if their spending were low enough. Therefore, we used a difference measure, which ensures that hospitals categorized as high value were always of high quality. To facilitate interpretation, both spending and quality scores were standardized to range from 0 to 10; thus, higher scores indicate higher value, with the maximum value score being 10 and minimum -10.

For our second value outcome, we classified hospitals as 'high value' if they received a 4 or 5-star rating on Hospital Compare and had a MSPB score in the lowest quintile of all hospitals, and classified hospitals as 'low value' if they received a 1 or 2-star rating on Hospital Compare and had a MSPB score in the highest quintile of all hospitals; all other hospitals were classified as 'average' value. For this classification, we selected quintiles to categorize hospitals by their MSPB scores as providing meaningful cost distinctions while identifying adequate numbers of high and low value hospitals for analysis.

#### Other Variables

Hospital characteristics were linked from the American Hospital Association (AHA) survey for 2018[8] and the CMS Provider of Services (POS) file of 2018. We examined hospital characteristics that have previously been associated with quality: geographic region (New England, Middle Atlantic, East North Central, West North Central, South Atlantic, East South Central, West South Central, Mountain, and Pacific); teaching status (teaching, residency program, non-teaching); number of beds(<100, 100-199,200-299,300-399, and 400+); nurse to bed ratio (≤0.75, 0.75-1, 1-1.5, > 1.5); urban location (urban, rural); ownership status (for profit, private not for profit, public, federal); percentage of Medicare admissions that are for patients dually eligible for both Medicare and Medicaid (≤ 10%, 11%-20%, 21%-40%, and > 40%); total expenses (quintiles); total cost of supplies (quintiles), ratio of supply expenses to total expenses (quintiles); and disproportionate share hospital (DSH) patient percentage (quintiles). DSH is a Medicare payment adjustment based on the proportion of Medicare admissions at a hospital that are for patients enrolled in Medicaid [9]

# Patient and Public Involvement

There was no patient or public involvement in the design, conduct or reporting of this study.

Statistical Analysis

We summarized value score (mean and standard deviation) and value category (frequency and percent) overall and by hospital characteristics, testing for differences in category characteristic using ANOVA for the value score and chi-square tests for the value category. To assess the overall relationship between cost and quality of care, we graphed the distribution of MSPB scores over each Star rating using a box plot (Figure 1), and then plotted the value scores against the Star Rating summary scores to visualize the association of value with quality (Figure 2). Then, to identify the hospital characteristics independently associated with value, and to identify contrasts in associations with quality, we estimated a series of models. We first used bivariate linear regression models to estimate the relationship between each hospital characteristics and value, where the outcome was our continuous value score; all factors were then included in a final multivariate model. For comparison with factors associated with quality alone, we estimated identical models where the dependent variable was the Star Ratings summary score, rather than our value score. We next estimated a multinomial logit model with the 3-category value as the outcome using 'average' as the reference group, including all factors. We used a multinomial rather than ordinal model to avoid assuming that associations were monotone; that is, we anticipated that some factors might be associated with low value and not high, and some with high value but not low. Both analyses included the same set of hospital characteristics, and all models included indicators for each domain, equal to 1 if the domain was included in the hospital's overall star rating score. We reported overall Wald test p-values for each characteristic. We report numbers and percent of missing values, and use multiple imputation (with 20 imputations) to account for missing values in all models.

All analyses were done using SAS 9.4 and Stata 16.1 (2020, StataCorp, College Station TX). All statistical tests were two-tailed, and we used p<0.05 to determine statistical significance. The study was approved by the Yale and NYU Institutional Review Boards.

# **RESULTS**

Our final study sample included 2,914 hospitals that received both a Star Rating and a MSPB score, all of which matched the AHA Survey file for 2018 (Table 1). The value score had a mean (SD) of 0.58 (1.79); see Figures 1 and 2. A total of 286 were classified as high value and 258 as low value by the categorical definition (Table 1). The 286 high value hospitals represented 28.6% of 999 4 and 5-star hospitals and 46.8% of 611 low cost (bottom quintile MSPB) hospitals; the 258 low value hospitals represented 26.6% of 970 1 and 2-star hospitals and 49.3% of 523 high cost hospitals.

In biviariate analyses (Table 1), we found not-for-profit private and government owned hospitals to have much higher values scores than for profit private hospitals, with score of 0.39 and 1.38 versus 0.08. Non-teaching hospitals and those with residency programs both had three times the value score of teaching hospitals, while hospitals with fewer than 100 beds (1.48) and less than 10% dual eligible patients (1.74) had the highest values scores. Hospitals in rural areas and smaller percentages of DSH payments also had higher value scores (all P-values < 0.05). For the categorical bivariate comparisons (Table 1), results were similar: high value hospitals were more likely to be not-for-profit, have no teaching program, have more beds and a higher nurse to bed ratio, and have a lower percentage of dual-eligible admissions compared to low- and average- value hospitals (all P values < 0.001). Regionally, lower value hospitals were more often in the Southern and Mid-Atlantic regions, while high value hospitals were in Northern Central and Pacific Regions (P<0.001).

In the multivariable model, all were significantly associated with value except those related to total and supply expenses (Table 2). The largest value effects were again for non-teaching, not-for-profit and government hospitals, hospitals with fewer than 100 beds and those with fewer than 10% dual eligible admissions. In equivalent model using the quality score instead of value as an outcome, there was no

association with urban status, while, unlike with value, higher supply expenses were associated with higher quality; other patterns of effects were very similar.

In multinomial regression assessing the associations between characteristics and low, average, and high value classification (Table 3), we collapsed the region categories to account for sparse cells. In the final model Government owned and not-for-profit hospitals, hospitals with fewer than 100 beds, hospitals in the Western region, and those with lower percentages of dual eligible patients and DSH patients were more likely to be high value than average value (all P-values < 0.05). Correspondingly, public hospitals, those with the more beds, those in the Northeast region, and lowest nurse to bed ratio all had higher odds of being low-value (all P-values < 0.05).

# **DISCUSSION**

In this examination of hospital value of care, we identified hospitals that provided relatively higher quality of care to Medicare beneficiaries at lower comparative costs, as well as hospitals that provided lower quality care at a high level of spending. In bivariate analyses, both high and low value hospitals differed from other hospitals strongly in ownership, urban location, bed size and region, nurse-to-bed ratio, amount spent on supplies relative to total expenses, and percentage of dual-eligible patients.

Most associations were significant in the multivariable primary analysis using the value score; of these, all but those related to urbanicity and expenses persisted in the multinomial logit analysis using a categorical outcome of low, average or high value.

The most notable finding, however, may not be the associations of hospital characteristics with value, but that value and quality of care are not synonymous: though generally, the MSPB declined with increasing Star ratings (Figure 1), less than a third of 4 and 5-star hospitals were high value, while more than a fourth of 1 and 2-star hospitals were lowest value. The observed trend in spending across overall quality (Figure 1) as well as the pattern of value classification is very similar to that found previously for

patient experience Star Ratings [10,11]; though the patient experience Star Rating is a component of the overall Star Rating, it accounts for only 22% of the weight of the overall summary score, so it is unlikely to drive the current findings. This finding was also reflected in our model of quality score, which found urban status was not related to quality while quality increased with more spending on supplies. That quality and value are not identical indicates that it will be possible to reduce spending among even the highest quality hospitals; however there is clearly a tension between the two, for if quality improvements driven by increased supply spending (or other additional resources) that translate to greater spending, then only offsets in other areas can increase value.

Most of the associations we found for value are similar to those seen for hospital characteristics and quality, both here with the Star Ratings summary score and in prior research; for example, Delancy et. al found that hospitals with fewer beds, no medical school affiliation, and lower proportions of DSH patients were more likely to have higher Star ratings [12]. This was similar to the results of our analysis of the star summary quality score (Table 2). However, in our categorical analysis, we found no relationship between teaching status and high value. Geographic region also explained some of the difference between low value and average value hospitals across all of our analyses, with hospitals in East South Central and Mid-Atlantic regions having lowest value and being more likely to be low value than those in any other regions, and Pacific region hospitals being consistently high value in all models. Since our spending measure is adjusted for local wage and cost variations, this geographic variation is likely driven substantially by variations in quality.

Of most relevance to hospitals and policy makers are the associations with factors that are within the control of the hospital. A key factor that was related to low value in all analyses was nurse to bed ratio, with lower ratios being more often found at the lowest value hospitals – only 9 (3.5%) low value hospitals had a ratio of 2 or more, while 43 (15.0%) of high value hospitals had a ratio of at least 2. This finding suggests that high levels of nurse staffing, while increasing the expense of inpatient care, may

reduce overall Medicare payments while promoting high quality care. Importantly, this finding also highlights a perennial challenge for policymakers: the wrong pocket problem. Since the costs of increased staffing are borne by hospitals but the benefits of lower spending are reaped by payers and patients, there is no intrinsic incentive for hospitals to pursue such interventions. Payment policies seeking to improve value should explicitly take these challenges into account.

Conversely, 66 (25.6%) of low value hospitals reported at least 40% of their Medicare patients were also eligible for Medicaid, while only 11 (3.8%) of high value hospitals reported such high rates of dual eligibility, with proportion of DSH patients exhibiting a similar but weaker pattern. This may indicate that lower income patients have unmeasured comorbidities or care complexities that are not captured by the risk adjustment used in Star Ratings' underlying measures or that these patients drive costs in ways unrelated to quality of care (e.g., limited social support may lead to greater post-discharge costs), that these patients may have higher cost needs that are not completely adjusted for, or possibly that DSH patients are more likely to receive care at lower quality hospitals.

This study has the limitations of any observational study, including that no causal inferences can be drawn; for example, it may be that lower value hospitals have increased costs of care because of lower quality, rather than independent of it. In addition, we have only examined a few key factors that may be associated with quality, in order to better focus on value of care. And, we have focused only on quality and spending metrics largely related to Medicare fee-for-service beneficiaries; other, broader measures of quality and costs that included the entire patient population could plausibly produce different results. Moreover, Medicare payments are distorted somewhat by the Inpatient Prospective Payment System. However, CMS Overall Star Ratings captures quality of care across 7 domains, and the MSPB is a measure of spending with cost-of-living adjustments that plausibly represents the overall spending per patient resulting from a hospitalization.

It is worth considering that the MSPB does not include admissions that lead to transfers, nor admissions in which the patient does not survive 30 days. Both exclusions are likely to confound the relationship between the spending score and the mortality domain, which accounts for 22% of the Star Rating summary score, since publicly reported mortality measures typically measure outcomes within the same 30 days, and attribute them to the first hospital in the event of transfers. Excluding patients who do not survive and/or are transferred likely reduces the MSPB score (assuming these are more complex patients), and more so for hospitals with higher mortality or transfer rates. However, this would tend to bias our current findings toward the null, attenuating the relationship between spending value.

# **CONCLUSION**

We have shown that care quality and care value are not identical; there are high quality hospitals that are not high value. In addition, there are a number of structural factors, some under the control of hospitals such as nurse to bed ratio, that are strongly associated with being low or high value. These findings can inform efforts of policymakers and hospitals to increase the value of care.

#### **Contributions**

Drs Herrin, Venkatesh, Bernheim and Horwitz contributed to the conception and design of the study. Drs Herrin and Yu performed all analyses. Dr Herrin drafted the manuscript. Drs Venkatesh, Desai, Thiel, Lin, Bernheim and Horwitz contributed critical additions and conceptual revisions to the manuscript.

# **Competing Interests**

Drs. Bernheim, Herrin, Lin and Venkatesh recieve salary support from the Centers for Medicare and Medicaid Services to develop, implement, and maintain hospital performance outcome measures, including the methodology for the Overall Hospital Star Ratings, that are publicly reported. Drs. Horwitz and Yu have worked under contract to the Centers for Medicare and Medicaid Services to develop quality measures, including some used in the Overall Hospital Star Ratings program.

# **Funding**

This work was funded by the Agency for Healthcare Research and Quality, grant R01HS022882.

# **Data Sharing**

Star Ratings and Medicare spending data are publicly available, and an analytic file including those data can be provided by the authors. Data collected by the American Hospital Association are proprietary, and will not be available for open access sharing.

# **Ethics Approval**

This study was approved by the Institutional Review Boards of Yale University (ID 0903004927) and New York University (ID S19-01460).

#### **REFERENCES**

- U.S. Centers for Medicare & Medicaid Services. Hospital Readmissions Reduction Program (HRRP).
   Accessed 2020 Dec 08. <a href="https://www.cms.gov/medicare/medicare-fee-for-service-payment/acuteinpatientpps/readmissions-reduction-program.html">https://www.cms.gov/medicare/medicare-fee-for-service-payment/acuteinpatientpps/readmissions-reduction-program.html</a>
- U.S. Centers for Medicare and Medicaid Services. Hospital Compare Overall Ratings. Accessed 2020
   Dec 08. <a href="https://qualitynet.cms.gov/inpatient/public-reporting/overall-ratings">https://qualitynet.cms.gov/inpatient/public-reporting/overall-ratings</a>
- Doyle JJ, Graves JA, Gruber JA, Kleiner S. Do high-cost hospitals deliver better care? Evidence from ambulance referral patterns. National Bureau of Economic Research. http://www.nber.org/papers/w17936.pdf?new\_window=1. Accessed December 19, 2013.
- 4. Silber JH, Kaestner R, Even-Shoshan O, Wang Y, Bressler LJ. Aggressive treatment style and surgical outcomes. Health Serv Res. 2010;45(6 Pt 2):1872-1892.
- 5. Virtual Mentor. 2014;16(2):124-130.
- 6. QualityNet. https://www.qualitynet.org/inpatient/public-reporting/overall-ratings/resources
- 7. QualityNet. https://www.qualitynet.org/inpatient/measures/mspb/methodology
- American Hospital Association. American Hospital Association (AHA) Annual Survey Database –
   2018.
- 9. https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/dsh
- Liao JM, Zhou L, Navathe AS. Nationwide Hospital Performance on Publicly Reported Episode
   Spending Measures. J Hosp Med. 2020 Mar 18;15:E1-E7. doi: 10.12788/jhm.3377. Epub ahead of print. PMID: 32195657
- 11. Trzeciak S, Gaughan JP, Bosire J, Angelo M, Holzberg AS, Mazzarelli AJ. Association Between Medicare Star Ratings for Patient Experience and Medicare Spending per Beneficiary for US Hospitals. J Patient Exp. 2017 Mar;4(1):17-21. doi: 10.1177/2374373516685938. Epub 2017 Jan 12. PMID: 28725856; PMCID: PMC5513662

12. DeLancey JO, Softcheck J, Chung JW, Barnard C, Dahlke AR, Bilimoria KY. Associations Between Hospital Characteristics, Measure Reporting, and the Centers for Medicare & Medicaid Services Overall Hospital Quality Star Ratings. JAMA. 2017 May 16;317(19):2015-2017. doi: 10.1001/jama.2017.3148. PMID: 28510670; PMCID: PMC5815009.



**Table 1**. Characteristics of low, average, and high value hospitals.

	Value		,p			
Characteristic	Score		Low Value	Average	High Value	P-value
	Mean (SD)	P-value <sup>1</sup>	n(%) or	n(%) or	n(%) or	
			mean(SD)	mean(SD)	mean(SD)	
N	0.58 (1.79)		258 (100.0)	2370 (100.0)	286 (100.0)	
Star Rating		< 0.001				
1	-2.38 (1.26)		89 (34.5)	179 (7.6)	0 (0.0)	
2	-0.66 (1.02)		169 (65.5)	533 (22.5)	0 (0.0)	
3	0.71 (1.00)		0 (0.0)	945 (39.9)	0 (0.0)	
4	1.75 (1.00)		0 (0.0)	528 (22.3)	213 (74.5)	
5	3.16 (1.07)		0 (0.0)	185 (7.8)	73 (25.5)	
Ownership		< 0.001				< 0.001
Public	0.39 (1.80)		43 (16.7)	356 (15.0)	33 (11.5)	
Not for profit	0.77 (1.69)		128 (49.6)	1521 (64.2)	215 (75.2)	
For profit	0.08 (1.96)		87 (33.7)	478 (20.2)	36 (12.6)	
Government	1.38 (1.60)		0 (0.0)	15 (0.6)	2 (0.7)	
Teaching status		< 0.001				< 0.001
Nonteaching	0.87 (1.72)		22 (8.5)	381 (16.1)	48 (16.8)	
Residency	0.72 (1.75)		124 (48.1)	1130 (47.7)	168 (58.7)	
Teaching	0.25 (1.82)		112 (43.4)	859 (36.2)	70 (24.5)	
Beds		< 0.001				< 0.001
<100	1.48 (1.57)		21 (8.1)	575 (24.3)	155 (54.2)	
100-199	0.65 (1.65)		58 (22.5)	645 (27.2)	67 (23.4)	
200-299	0.21 (1.73)		54 (20.9)	399 (16.8)	33 (11.5)	
300-399	0.08 (1.86)		58 (22.5)	274 (11.6)	17 (5.9)	
400+	-0.12 (1.74)		67 (26.0)	477 (20.1)	14 (4.9)	
Urbanity	, ,	< 0.001	, ,			< 0.001
Rural	1.08 (1.52)		29 (11.2)	579 (24.4)	110 (38.5)	
Urban	0.41 (1.84)		229 (88.8)	1791 (75.6)	176 (61.5)	
Region	. ,	< 0.001	` ,	, ,	, ,	< 0.001
New England	0.70 (1.48)		3 (1.2)	115 (4.9)	9 (3.1)	
Mid Atlantic	-0.27 (1.92)		62 (24.0)	260 (11.0)	13 (4.5)	
E North Central	1.02 (1.58)		22 (8.5)	382 (16.1)	57 (19.9)	
W North Central	1.60 (1.48)		1 (0.4)	179 (7.6)	50 (17.5)	
South Atlantic	0.40 (1.71)		43 (16.7)	411 (17.3)	42 (14.7)	
E South Central	0.15 (1.46)		23 (8.9)	236 (10.0)	11 (3.8)	
W South Central	0.17 (1.80)		60 (23.3)	341 (14.4)	20 (7.0)	
Mountain	1.13 (1.78)		12 (4.7)	155 (6.5)	31 (10.8)	
Pacific	0.81 (1.91)		32 (12.4)	291 (12.3)	53 (18.5)	
Nurse/Beds	0.00 (0.00 0)	< 0.001	()		()	< 0.001
<=0.75	0.51 (1.88)		86 (33.3)	536 (22.6)	79 (27.6)	
0.75-1	0.35 (1.73)		52 (20.2)	405 (17.1)	47 (16.4)	
1-1.5	0.41 (1.72)		86 (33.3)	783 (33.0)	65 (22.7)	
1.5-2	0.79 (1.81)		25 (9.7)	381 (16.1)	52 (18.2)	
2+	1.24 (1.67)		9 (3.5)	265 (11.2)	43 (15.0)	
	(=)		- (3.5)	\/	- ()	

Table 1 continued

% Duals		< 0.001				< 0.001
<=10%	1.74 (1.93)		7 (2.7)	176 (7.4)	33 (11.5)	
11%-20%	0.80 (1.58)		67 (26.0)	863 (36.4)	91 (31.8)	
21%-40%	0.51 (1.71)		118 (45.7)	1057 (44.6)	151 (52.8)	
>40%	-0.56 (1.94)		66 (25.6)	274 (11.6)	11 (3.8)	
Total Expenses		< 0.001				< 0.001
Quintile 1	1.09 (1.66)		28 (10.9)	459 (19.4)	95 (33.2)	
Quintile 2	0.79 (1.77)		54 (20.9)	446 (18.8)	81 (28.3)	
Quintile 3	0.44 (1.73)		57 (22.1)	475 (20.0)	50 (17.5)	
Quintile 4	0.31 (1.81)		71 (27.5)	473 (20.0)	37 (12.9)	
Quintile 5	0.24 (1.83)		48 (18.6)	511 (21.6)	22 (7.7)	
Missing	0.68 (1.61)		0 (0.0)	6 (0.3)	1 (0.3)	
Supplies \$		< 0.001				< 0.001
Quintile 1	1.11 (1.63)		23 (8.9)	301 (12.7)	72 (25.2)	
Quintile 2	0.83 (1.64)		31 (12.0)	311 (13.1)	53 (18.5)	
Quintile 3	0.50 (1.82)		35 (13.6)	326 (13.8)	35 (12.2)	
Quintile 4	0.59 (1.72)		37 (14.3)	334 (14.1)	24 (8.4)	
Quintile 5	0.27 (1.81)		34 (13.2)	349 (14.7)	12 (4.2)	
Missing	0.40 (1.86)		98 (38.0)	749 (31.6)	90 (31.5)	
Supplies/Total %		0.042				0.002
Quintile 1	0.83 (1.79)		29 (11.2)	311 (13.1)	56 (19.6)	
Quintile 2	0.75 (1.69)		23 (8.9)	329 (13.9)	43 (15.0)	
Quintile 3	0.51 (1.75)		38 (14.7)	317 (13.4)	41 (14.3)	
Quintile 4	0.53 (1.56)		28 (10.9)	344 (14.5)	23 (8.0)	
Quintile 5	0.68 (1.92)		42 (16.3)	320 (13.5)	33 (11.5)	
Missing	0.40 (1.86)		98 (38.0)	749 (31.6)	90 (31.5)	
% DSH	, ,	< 0.001				< 0.001
Quintile 1	1.35 (1.74)		32 (12.4)	468 (19.7)	84 (29.4)	
Quintile 2	0.85 (1.62)		33 (12.8)	480 (20.3)	70 (24.5)	
Quintile 3	0.63 (1.59)		42 (16.3)	488 (20.6)	52 (18.2)	
Quintile 4	0.30 (1.60)		55 (21.3)	484 (20.4)	44 (15.4)	
Quintile 5	-0.26 (1.96)		96 (37.2)	450 (19.0)	36 (12.6)	
	` '		•		•	

Source: Center for Medicare and Medicaid Services Overall Star Ratings score and Medicare Spending per Beneficiary; American Hospital Association Survey.

## Notes

1. P-value based on ANOVA model estimated using multiple imputation for missing values.

**Table 2**. Results of multivariable linear models with value score and Star Ratings summary score as dependent variables.

Ownership		Value		Quality	
Public         ref         ref           Not for profit         0.40 (0.21,0.59)         0.34 (0.20,0.48)           For profit         -0.04 (-0.29,0.21)         0.14 (-0.05,0.33)           Government         0.96 (-0.23,2.15)         0.47 (-0.42,1.36)           Teaching status         0.008         0.006           Nonteaching         ref         ref           Residency         -0.30 (-0.50,-0.10)         -0.19 (-0.34,-0.05)           Teaching         -0.27 (-0.47,-0.07)         -0.24 (-0.39,-0.09)           Beds         <0.001         <0.001           <100         ref         ref           100-199         -0.55 (-0.77,-0.33)         -0.31 (-0.47,-0.14)           200-299         -0.99 (-1.29,-0.70)         -0.57 (-0.79,-0.36)           300-399         -1.24 (-1.57,-0.90)         -0.67 (-0.93,-0.42)           400+         -1.45 (-1.81,-1.08)         -0.80 (-1.07,-0.52)           Urbanity         0.015         0.894           Rural         ref         ref           Urban         -0.23 (-0.42,-0.04)         -0.01 (-0.15,0.13)           Region         <0.001         <0.001           New England         ref         ref           Mid Atlantic         -0.67 (-1.03,-	Characteristic	Coeff (SE)	P-value		
Not for profit         0.40 (0.21,0.59)         0.34 (0.20,0.48)           For profit         -0.04 (-0.29,0.21)         0.14 (-0.05,0.33)           Government         0.96 (-0.23,2.15)         0.47 (-0.42,1.36)           Teaching status         0.008         0.006           Nonteaching         ref         ref           Residency         -0.30 (-0.50,-0.10)         -0.19 (-0.34,-0.05)           Teaching         -0.27 (-0.47,-0.07)         -0.24 (-0.39,-0.09)           Beds         <0.001	Ownership		<0.001	<0.	001
For profit	Public	ref		ref	
Government         0.96 (-0.23,2.15)         0.47 (-0.42,1.36)           Teaching status         0.008         0.006           Nonteaching         ref         ref           Residency         -0.30 (-0.50,-0.10)         -0.19 (-0.34,-0.05)           Teaching         -0.27 (-0.47,-0.07)         -0.24 (-0.39,-0.09)           Beds         <0.001	Not for profit	0.40 (0.21,0.59)		0.34 (0.20,0.48)	
Teaching status         0.008         0.006           Nonteaching         ref         ref           Residency         -0.30 (-0.50,-0.10)         -0.19 (-0.34,-0.05)           Teaching         -0.27 (-0.47,-0.07)         -0.24 (-0.39,-0.09)           Beds         <0.001	For profit	-0.04 (-0.29,0.21)		0.14 (-0.05,0.33)	
Nonteaching         ref         ref           Residency         -0.30 (-0.50,-0.10)         -0.19 (-0.34,-0.05)           Teaching         -0.27 (-0.47,-0.07)         -0.24 (-0.39,-0.09)           Beds         <0.001	Government	0.96 (-0.23,2.15)		0.47 (-0.42,1.36)	
Residency         -0.30 (-0.50,-0.10)         -0.19 (-0.34,-0.05)           Teaching         -0.27 (-0.47,-0.07)         -0.24 (-0.39,-0.09)           Beds         <0.001	Teaching status		0.008	0.0	006
Teaching         -0.27 (-0.47,-0.07)         -0.24 (-0.39,-0.09)           Beds         <0.001	Nonteaching	ref		ref	
Seds	Residency	-0.30 (-0.50,-0.10)		-0.19 (-0.34,-0.05)	
<100	Teaching	-0.27 (-0.47,-0.07)		-0.24 (-0.39,-0.09)	
100-199	Beds		< 0.001	<0.	001
200-299	<100	ref		ref	
300-399	100-199	-0.55 (-0.77,-0.33)		-0.31 (-0.47,-0.14)	
400+       -1.45 (-1.81,-1.08)       -0.80 (-1.07,-0.52)         Urbanity       0.015       0.894         Rural       ref       ref         Urban       -0.23 (-0.42,-0.04)       -0.01 (-0.15,0.13)         Region       <0.001	200-299	-0.99 (-1.29,-0.70)		-0.57 (-0.79,-0.36)	
Urbanity       0.015       0.894         Rural       ref       ref         Urban       -0.23 (-0.42,-0.04)       -0.01 (-0.15,0.13)         Region       <0.001	300-399	-1.24 (-1.57,-0.90)		-0.67 (-0.93,-0.42)	
Rural ref ref Urban -0.23 (-0.42,-0.04) -0.01 (-0.15,0.13)  Region <0.001 <0.001  New England ref ref Mid Atlantic -0.67 (-1.03,-0.31) -0.56 (-0.83,-0.29)  E North Central 0.35 (0.01,0.70) 0.27 (0.02,0.53)  W North Central 0.57 (0.20,0.95) 0.03 (-0.25,0.31)  South Atlantic -0.09 (-0.45,0.27) -0.25 (-0.52,0.02)  E South Central -0.70 (-1.13,-0.28) -0.55 (-0.86,-0.23)  W South Central -0.62 (-0.99,-0.26) -0.15 (-0.42,0.12)  Mountain 0.23 (-0.19,0.65) -0.02 (-0.34,0.29)  Pacific 0.83 (0.45,1.22) 0.35 (0.06,0.64)  Nurse/Beds 0.006 0.001  <=0.75 ref ref 0.75-1 -0.19 (-0.40,0.03) -0.21 (-0.37,-0.05) 1-1.5 -0.14 (-0.34,0.05) -0.16 (-0.30,-0.01)	400+	-1.45 (-1.81,-1.08)		-0.80 (-1.07,-0.52)	
Urban       -0.23 (-0.42,-0.04)       -0.01 (-0.15,0.13)         Region       <0.001	Urbanity		0.015	0.8	394
Region       <0.001       <0.001         New England       ref       ref         Mid Atlantic       -0.67 (-1.03,-0.31)       -0.56 (-0.83,-0.29)         E North Central       0.35 (0.01,0.70)       0.27 (0.02,0.53)         W North Central       0.57 (0.20,0.95)       0.03 (-0.25,0.31)         South Atlantic       -0.09 (-0.45,0.27)       -0.25 (-0.52,0.02)         E South Central       -0.70 (-1.13,-0.28)       -0.55 (-0.86,-0.23)         W South Central       -0.62 (-0.99,-0.26)       -0.15 (-0.42,0.12)         Mountain       0.23 (-0.19,0.65)       -0.02 (-0.34,0.29)         Pacific       0.83 (0.45,1.22)       0.35 (0.06,0.64)         Nurse/Beds       0.006       0.001         <=0.75	Rural	ref		ref	
New England       ref       ref         Mid Atlantic       -0.67 (-1.03,-0.31)       -0.56 (-0.83,-0.29)         E North Central       0.35 (0.01,0.70)       0.27 (0.02,0.53)         W North Central       0.57 (0.20,0.95)       0.03 (-0.25,0.31)         South Atlantic       -0.09 (-0.45,0.27)       -0.25 (-0.52,0.02)         E South Central       -0.70 (-1.13,-0.28)       -0.55 (-0.86,-0.23)         W South Central       -0.62 (-0.99,-0.26)       -0.15 (-0.42,0.12)         Mountain       0.23 (-0.19,0.65)       -0.02 (-0.34,0.29)         Pacific       0.83 (0.45,1.22)       0.35 (0.06,0.64)         Nurse/Beds       0.006       0.001         <=0.75	Urban	-0.23 (-0.42,-0.04)		-0.01 (-0.15,0.13)	
Mid Atlantic       -0.67 (-1.03,-0.31)       -0.56 (-0.83,-0.29)         E North Central       0.35 (0.01,0.70)       0.27 (0.02,0.53)         W North Central       0.57 (0.20,0.95)       0.03 (-0.25,0.31)         South Atlantic       -0.09 (-0.45,0.27)       -0.25 (-0.52,0.02)         E South Central       -0.70 (-1.13,-0.28)       -0.55 (-0.86,-0.23)         W South Central       -0.62 (-0.99,-0.26)       -0.15 (-0.42,0.12)         Mountain       0.23 (-0.19,0.65)       -0.02 (-0.34,0.29)         Pacific       0.83 (0.45,1.22)       0.35 (0.06,0.64)         Nurse/Beds       0.006       0.001         <=0.75	Region		<0.001	<0.	001
E North Central 0.35 (0.01,0.70) 0.27 (0.02,0.53)  W North Central 0.57 (0.20,0.95) 0.03 (-0.25,0.31)  South Atlantic -0.09 (-0.45,0.27) -0.25 (-0.52,0.02)  E South Central -0.70 (-1.13,-0.28) -0.55 (-0.86,-0.23)  W South Central -0.62 (-0.99,-0.26) -0.15 (-0.42,0.12)  Mountain 0.23 (-0.19,0.65) -0.02 (-0.34,0.29)  Pacific 0.83 (0.45,1.22) 0.35 (0.06,0.64)  Nurse/Beds 0.006 0.001  <=0.75 ref ref 0.75-1 -0.19 (-0.40,0.03) -0.21 (-0.37,-0.05) 1-1.5 -0.14 (-0.34,0.05) -0.16 (-0.30,-0.01)	New England	ref		ref	
W North Central 0.57 (0.20,0.95) 0.03 (-0.25,0.31)  South Atlantic -0.09 (-0.45,0.27) -0.25 (-0.52,0.02)  E South Central -0.70 (-1.13,-0.28) -0.55 (-0.86,-0.23)  W South Central -0.62 (-0.99,-0.26) -0.15 (-0.42,0.12)  Mountain 0.23 (-0.19,0.65) -0.02 (-0.34,0.29)  Pacific 0.83 (0.45,1.22) 0.35 (0.06,0.64)  Nurse/Beds 0.006 0.001  <=0.75 ref ref 0.75-1 -0.19 (-0.40,0.03) -0.21 (-0.37,-0.05) 1-1.5 -0.14 (-0.34,0.05) -0.16 (-0.30,-0.01)	Mid Atlantic	-0.67 (-1.03,-0.31)		-0.56 (-0.83,-0.29)	
South Atlantic       -0.09 (-0.45,0.27)       -0.25 (-0.52,0.02)         E South Central       -0.70 (-1.13,-0.28)       -0.55 (-0.86,-0.23)         W South Central       -0.62 (-0.99,-0.26)       -0.15 (-0.42,0.12)         Mountain       0.23 (-0.19,0.65)       -0.02 (-0.34,0.29)         Pacific       0.83 (0.45,1.22)       0.35 (0.06,0.64)         Nurse/Beds       0.006       0.001         <=0.75	E North Central	0.35 (0.01,0.70)		0.27 (0.02,0.53)	
E South Central -0.70 (-1.13,-0.28) -0.55 (-0.86,-0.23)  W South Central -0.62 (-0.99,-0.26) -0.15 (-0.42,0.12)  Mountain 0.23 (-0.19,0.65) -0.02 (-0.34,0.29)  Pacific 0.83 (0.45,1.22) 0.35 (0.06,0.64)  Nurse/Beds 0.006 0.001  <=0.75 ref ref 0.75-1 -0.19 (-0.40,0.03) -0.21 (-0.37,-0.05) 1-1.5 -0.14 (-0.34,0.05) -0.16 (-0.30,-0.01)	W North Central	0.57 (0.20,0.95)		0.03 (-0.25,0.31)	
W South Central -0.62 (-0.99,-0.26) -0.15 (-0.42,0.12)  Mountain 0.23 (-0.19,0.65) -0.02 (-0.34,0.29)  Pacific 0.83 (0.45,1.22) 0.35 (0.06,0.64)  Nurse/Beds 0.006 0.001  <=0.75 ref ref 0.75-1 -0.19 (-0.40,0.03) -0.21 (-0.37,-0.05) 1-1.5 -0.14 (-0.34,0.05) -0.16 (-0.30,-0.01)	South Atlantic	-0.09 (-0.45,0.27)		-0.25 (-0.52,0.02)	
Mountain       0.23 (-0.19,0.65)       -0.02 (-0.34,0.29)         Pacific       0.83 (0.45,1.22)       0.35 (0.06,0.64)         Nurse/Beds       0.006       0.001         <=0.75	E South Central	-0.70 (-1.13,-0.28)		-0.55 (-0.86,-0.23)	
Pacific       0.83 (0.45,1.22)       0.35 (0.06,0.64)         Nurse/Beds       0.006       0.001         <=0.75       ref       ref         0.75-1       -0.19 (-0.40,0.03)       -0.21 (-0.37,-0.05)         1-1.5       -0.14 (-0.34,0.05)       -0.16 (-0.30,-0.01)	W South Central	-0.62 (-0.99,-0.26)		-0.15 (-0.42,0.12)	
Nurse/Beds       0.006       0.001         <=0.75	Mountain	0.23 (-0.19,0.65)		-0.02 (-0.34,0.29)	
<=0.75 ref ref 0.75-1 -0.19 (-0.40,0.03) -0.21 (-0.37,-0.05) 1-1.5 -0.14 (-0.34,0.05) -0.16 (-0.30,-0.01)	Pacific	0.83 (0.45,1.22)		0.35 (0.06,0.64)	
0.75-1       -0.19 (-0.40,0.03)       -0.21 (-0.37,-0.05)         1-1.5       -0.14 (-0.34,0.05)       -0.16 (-0.30,-0.01)	Nurse/Beds		0.006	0.0	001
1-1.5 -0.14 (-0.34,0.05) -0.16 (-0.30,-0.01)	<=0.75	ref		ref	
	0.75-1	-0.19 (-0.40,0.03)		-0.21 (-0.37,-0.05)	
1.5-2 0.05 (-0.18,0.28) 0.03 (-0.14,0.20)	1-1.5	-0.14 (-0.34,0.05)		-0.16 (-0.30,-0.01)	
	1.5-2	0.05 (-0.18,0.28)		0.03 (-0.14,0.20)	
2+ 0.22 (-0.04,0.48) 0.10 (-0.10,0.29)	2+	0.22 (-0.04,0.48)		0.10 (-0.10,0.29)	

Table 2. continued

	Value		Quality		
Characteristic	Coeff (SE)	P-value	Coeff (SE)	P-value	
% Duals		< 0.001		<0.001	
<=10%	ref		ref		
11%-20%	-0.48 (-0.78,-0.18)		-0.26 (-0.48,-0.03)		
21%-40%	-0.81 (-1.14,-0.48)		-0.68 (-0.93,-0.43)		
>40%	-1.66 (-2.10,-1.23)		-1.20 (-1.53,-0.88)		
Total Expenses		0.821		0.790	
Quintile 1	ref		ref		
Quintile 2	-0.01 (-0.31,0.29)		-0.09 (-0.32,0.13)		
Quintile 3	-0.15 (-0.54,0.24)		-0.15 (-0.44,0.14)		
Quintile 4	-0.12 (-0.60,0.35)		-0.17 (-0.52,0.18)		
Quintile 5	-0.24 (-0.81,0.34)		-0.27 (-0.71,0.16)		
Supplies \$		0.034		0.008	
Quintile 1	ref		ref		
Quintile 2	-0.06 (-0.37,0.24)		0.07 (-0.16,0.30)		
Quintile 3	-0.06 (-0.47,0.34)		0.01 (-0.29,0.31)		
Quintile 4	0.30 (-0.20,0.79)		0.34 (-0.03,0.71)		
Quintile 5	0.50 (-0.09,1.10)		0.50 (0.05,0.94)		
Supplies/Total %		0.424		0.479	
Quintile 1	ref		ref		
Quintile 2	-0.10 (-0.32,0.12)		0.03 (-0.14,0.19)		
Quintile 3	-0.21 (-0.45,0.03)		0.06 (-0.12,0.24)		
Quintile 4	-0.23 (-0.49,0.03)		0.02 (-0.17,0.22)		
Quintile 5	-0.24 (-0.54,0.06)		-0.09 (-0.31,0.13)		
% DISH		0.003		<0.001	
Quintile 1	ref		ref		
Quintile 2	-0.17 (-0.39,0.04)		-0.27 (-0.43,-0.10)		
Quintile 3	-0.32 (-0.55,-0.09)		-0.40 (-0.57,-0.23)		
Quintile 4	-0.38 (-0.63,-0.14)		-0.37 (-0.55,-0.18)		
Quintile 5	-0.56 (-0.85,-0.27)		-0.66 (-0.88,-0.44)		

Source: Center for Medicare and Medicaid Services Overall Star Ratings score and Medicare Spending per Beneficiary; American Hospital Association Survey.

**Table 3** Results of multilogit model for low and high value versus average value hospitals.

	Low Value		High Value	
Characteristic	OR (95% CI)	P-value	OR (95% CI)	P-value
Ownership		<0.001		0.002
Public	ref		ref	
Not for profit	0.4 (0.2,0.6)		2.1 (1.2,3.5)	
For profit	0.8 (0.5,1.5)		0.8 (0.4,1.8)	
Government	0.0 (0.0,.)		2.0 (0.3,13.3)	
Teaching status		0.012		0.698
Nonteaching	ref		ref	
Residency	2.9 (1.4,5.8)		1.1 (0.7,1.7)	
Teaching	2.4 (1.2,4.9)		1.2 (0.7,2.1)	
Beds		<0.001		0.003
<100	ref		ref	
100-199	2.9 (1.2,6.8)		0.5 (0.3,0.7)	
200-299	5.9 (2.2,16.1)		0.4 (0.2,0.8)	
300-399	11.9 (4.1,34.5)		0.3 (0.1,0.8)	
400+	10.0 (3.2,31.7)		0.2 (0.1,0.5)	
Urbanity		0.300		0.758
Rural	ref		ref	
Urban	1.4 (0.8,2.5)		1.1 (0.7,1.6)	
Region <sup>1</sup>		<0.001		< 0.001
West	ref		ref	
Midwest	0.5 (0.2,1.2)		0.4 (0.2,0.6)	
Northeast	2.7 (1.4,5.3)		0.1 (0.1,0.3)	
South	1.3 (0.7,2.4)		0.3 (0.2,0.4)	
Nurse/Beds		0.025		0.673
<=0.75	ref		ref	
0.75-1	1.0 (0.6,1.7)		1.0 (0.6,1.7)	
1-1.5	0.8 (0.5,1.3)		0.8 (0.5,1.2)	
1.5-2	0.5 (0.2,0.9)		1.1 (0.6,1.9)	
2+	0.3 (0.1,0.7)		1.0 (0.5,1.8)	

Table 3 Continued

% Duals		0.427		0.009
<=10%	ref		ref	
11%-20%	2.2 (0.6,7.8)		1.1 (0.6,2.2)	
21%-40%	2.7 (0.7,9.8)		1.9 (0.9,4.0)	
>40%	3.2 (0.8,13.2)		0.6 (0.2,2.0)	
Total Expenses	, , ,	0.536	, , ,	0.626
Quintile 1	ref		ref	
Quintile 2	1.7 (0.7,4.6)		1.4 (0.8,2.6)	
Quintile 3	1.2 (0.4,4.0)		1.1 (0.5,2.4)	
Quintile 4	1.3 (0.3,5.1)		1.2 (0.4,3.5)	
Quintile 5	0.9 (0.2,4.4)		0.9 (0.2,4.1)	
Supplies \$		0.660		0.845
Quintile 1	ref		ref	
Quintile 2	0.6 (0.2,1.4)		0.7 (0.4,1.4)	
Quintile 3	0.4 (0.1,1.3)		0.6 (0.2,1.4)	
Quintile 4	0.4 (0.1,1.4)		0.5 (0.2,1.7)	
Quintile 5	0.3 (0.1,1.5)		0.5 (0.1,2.6)	
Supplies/Total %		0.010		0.273
Quintile 1	ref		ref	
Quintile 2	1.0 (0.5,2.0)		0.9 (0.5,1.4)	
Quintile 3	2.0 (1.0,4.0)		1.2 (0.7,2.1)	
Quintile 4	1.4 (0.7,3.1)		0.6 (0.3,1.3)	
Quintile 5	3.0 (1.3,6.8)		1.1 (0.5,2.3)	
% DISH		0.048		0.006
Quintile 1	ref		ref	
Quintile 2	0.7 (0.4,1.4)		0.8 (0.5,1.4)	
Quintile 3	1.1 (0.6,2.0)		0.5 (0.3,0.9)	
Quintile 4	0.6 (0.3,1.3)		0.4 (0.2,0.7)	
Quintile 5	1.6 (0.7,3.3)		0.5 (0.2,1.0)	

Source: Center for Medicare and Medicaid Services Overall Star Ratings score and Medicare Spending per Beneficiary; American Hospital Association Survey.

## Notes:

1. Region groups were collapsed due to sparse cells under original groupings.

Figure 1. Medicare Spending per Beneificiary by Overall Star Rating

Figure 2. Value score versus Star Rating summary score.






Figure 1. Medicare Spending per Beneificiary by Overall Star Rating  $139 \times 101 \text{mm}$  (600  $\times$  600 DPI)




Figure 2. Value score versus Star Rating summary score. 139x101mm~(600~x~600~DPI)

## IDENTIFYING HIGH VALUE CARE FOR MEDICARE BENEFICIARIES: A CROSS-SECTIONAL STUDY OF ACUTE CARE HOSPITALS IN THE U.S.

## **STROBE Statement**

Checklist of items that should be included in reports of *cross-sectional studies* 

Source: https://www.strobe-statement.org/download/strobe-checklist-cross-sectional-studies-doc

Item	Item No		
	NO	Recommendation	Status
Title and abstract	1	(a) Indicate the study's design with a commonly used	Done
		term in the title or the abstract	
		(b) Provide in the abstract an informative and	Done
		balanced summary of what was done and what was	
		found	
ntroduction			
Background/rationale	2	Explain the scientific background and rationale for	Done
		the investigation being reported	
Objectives	3	State specific objectives, including any prespecified	Done
		hypotheses	
Methods			
Study design	4	Present key elements of study design early in the	Brief overview in Introduction.
		paper	Details in Methods
Setting	5	Describe the setting, locations, and relevant dates,	Done
		including periods of recruitment, exposure, follow-	
		up, and data collection	
Participants	6	(a) Give the eligibility criteria, and the sources and	Done
		methods of selection of participants	
Variables	7	Clearly define all outcomes, exposures, predictors,	Done
		potential confounders, and effect modifiers. Give	
		diagnostic criteria, if applicable	
Data sources/	8*	For each variable of interest, give sources of data	Done
measurement		and details of methods of assessment	
		(measurement). Describe comparability of	
		assessment methods if there is more than one group	
Bias	9	Describe any efforts to address potential sources of	NA
		bias	
Study size	10	Explain how the study size was arrived at	Done
Quantitative variables	11	Explain how quantitative variables were handled in	Done
Quantitative variables		the analyses. If applicable, describe which groupings	Done
		were chosen and why	
Statistical methods	12	(a) Describe all statistical methods, including those	Done
Judistical methods	12	used to control for confounding	Donc
		(b) Describe any methods used to examine	Done
		subgroups and interactions	DOILE
			Dono
		(c) Explain how missing data were addressed	Done
		(d) If applicable, describe analytical methods taking	Done
		account of sampling strategy	

		$(\underline{e})$ Describe any sensitivity analyses	NA
Results			
Participants	13*	(a) Report numbers of individuals at each stage of	NA
		study—eg numbers potentially eligible, examined for	
		eligibility, confirmed eligible, included in the study,	
		completing follow-up, and analysed	
		(b) Give reasons for non-participation at each stage	NA
		(c) Consider use of a flow diagram	NA
Descriptive data	14*	(a) Give characteristics of study participants (eg	Done
		demographic, clinical, social) and information on	
		exposures and potential confounders	
		(b) Indicate number of participants with missing data	Done
		for each variable of interest	
Outcome data	15*	Report numbers of outcome events or summary	Done
		measures	
Main results	16	(a) Give unadjusted estimates and, if applicable,	Done
		confounder-adjusted estimates and their precision	
		(eg, 95% confidence interval). Make clear which	
		confounders were adjusted for and why they were	
		included	
		(b) Report category boundaries when continuous	Done
		variables were categorized	
		(c) If relevant, consider translating estimates of	NA
		relative risk into absolute risk for a meaningful time	
		period	
Other analyses	17	Report other analyses done—eg analyses of	Done
		subgroups and interactions, and sensitivity analyses	
Discussion			
Key results	18	Summarise key results with reference to study	Done
		objectives	
Limitations	19	Discuss limitations of the study, taking into account	Done
		sources of potential bias or imprecision. Discuss both	
		direction and magnitude of any potential bias	
Interpretation	20	Give a cautious overall interpretation of results	Done
		considering objectives, limitations, multiplicity of	
		analyses, results from similar studies, and other	
		relevant evidence	
Generalisability	21	Discuss the generalisability (external validity) of the	Done
		study results	
Other information			
Funding	22	Give the source of funding and the role of the	Done
		funders for the present study and, if applicable, for	
		the original study on which the present article is	
		based	

<sup>\*</sup>Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web Annals
AOBE Initiativ sites of PLoS Medicine at http://www.plosmedicine.org/, Annals of Internal Medicine at http://www.annals.org/, and Epidemiology at http://www.epidem.com/). Information on the STROBE Initiative is available at www.strobe-statement.org.