

# BMJ Open Food insecurity and associated depression among older adults in India: evidence from a population-based study

Muhammad T <sup>1</sup>, KM Sulaiman,<sup>2</sup> Drishti Drishti,<sup>3</sup> Shobhit Srivastava <sup>4</sup>

**To cite:** T M, Sulaiman KM, Drishti D, *et al.* Food insecurity and associated depression among older adults in India: evidence from a population-based study. *BMJ Open* 2022;**12**:e052718. doi:10.1136/bmjopen-2021-052718

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-052718>).

Received 24 April 2021  
Accepted 23 March 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Department of Family & Generations, International Institute for Population Sciences, Mumbai, India

<sup>2</sup>Department of Migration & Urban Studies, International Institute for Population Sciences, Mumbai, Maharashtra, India

<sup>3</sup>Department of Public Health & Mortality Studies, International Institute for Population Sciences, Mumbai, India

<sup>4</sup>Department of Survey Research & Data Analytics, International Institute for Population Sciences, Mumbai, India

## Correspondence to

Shobhit Srivastava;  
shobhitsrivastava889@gmail.com

## ABSTRACT

**Objective** The present study aimed to examine the associations of several indicators of food insecurity with depression among older adults in India.

**Design** A cross-sectional study was conducted using country-representative survey data.

**Setting and participants** The present study uses data of the Longitudinal Aging Study in India conducted during 2017–2018. The effective sample size for the present study was 31 464 older adults aged 60 years and above.

**Primary and secondary outcome measures** The outcome variable was major depression among older adults. Descriptive statistics along with bivariate analysis was presented. Additionally, binary logistic regression analysis was used to establish the association between the depression and food security factors along with other covariates.

**Results** The overall prevalence of major depression was 8.4% among older adults in India. A proportion of 6.3% of the older adults reduced the size of meals, 40% reported that they did not eat enough food of their choice, 5.6% mentioned that they were hungry but did not eat, 4.2% reported that they did not eat for a whole day and 5.6% think that they have lost weight due to lack of enough food in the household. Older adults who reported to have reduced the size of meals due to lack of enough food (adjusted OR (AOR): 1.76, CI 1.44 to 2.15) were hungry but did not eat (AOR: 1.35, CI 1.06 to 1.72) did not eat food for a whole day (AOR: 1.33; CI 1.03 to 1.71), lost weight due to lack of food (AOR: 1.57; CI 1.30 to 1.89) had higher odds of being depressed in reference to their respective counterparts.

**Conclusion** The findings suggest that self-reported food insecurity indicators were strongly associated with major depression among older Indian adults. The national food security programmes should be enhanced as an effort to improve mental health status and quality of life among older population.

## INTRODUCTION

Food insecurity is defined as not having physical, social and economic access to sufficient, safe and nutritious food that satisfies their dietary needs and food choices for a productive and healthy life.<sup>1 2</sup> About 815 million people live in this situation globally.<sup>3</sup> To support this population, sustainable development goals, targets 2.1 and 2.2, emphasise


## Strengths and limitations of this study

- The study uses a large nationally representative sample of older population.
- Cross-sectional design is a limitation of the study as it is impossible to establish the observed directions of the relationships.
- The food security indicators were self-reported which may result in recall and reporting biases.

ending hunger and all forms of malnutrition.<sup>4</sup> Food insecurity incorporates more than just the current nutritional state, capturing as well vulnerability to anticipated disturbances in access to adequate and appropriate food.<sup>5–7</sup> After the economic liberalisations, developing countries struggle to meet global nutritional standards and ensure food security.<sup>8</sup> Food security has been a policy priority in India for a long time, mainly focusing on its vulnerable populations like children and older adults.<sup>9–11</sup>

In adult populations, food insecurity is associated with insufficient dietary consumption, nutritional status and poor physical and mental well-being.<sup>12</sup> A couple of studies found that food insecurity is related to poor social and functional health, hypertension, diabetes and anxiety.<sup>12–16</sup> Empirical evidence pointed out that the prevalence of food insecurity is exceptionally high among older adults<sup>17–19</sup> due to physical limitations, poor heart conditions, social isolation and lack of transportation.<sup>20–23</sup> Similarly, food insecure older adults have been reported to spend less on healthcare<sup>24</sup> and to show higher levels of non-adherence to medical treatments due to financial limitations.<sup>25</sup> Therefore, among older adults, food insecurity has been linked with poor health status,<sup>26</sup> lower cognitive performance<sup>27</sup> and, notably, higher risk of depression.<sup>28</sup>

The WHO defines depression as a mental disorder characterised by sadness, lack of interest or pleasure, guilt or low self-worth,



**Figure 1** Conceptual framework of major depression. ADL, Activities of Daily Living; IADL, Instrumental ADL; MPCE, monthly per capita consumption expenditure.

disordered sleep or appetite, feelings of tiredness and reduced concentration.<sup>29</sup> Research has shown a relationship between depression and various socioeconomic variables such as old age, low level of education, hunger and physical labour.<sup>30–31</sup> Depressive disorders are the most common psychiatric condition among older people.<sup>32–33</sup> Recent evidence recognised several factors associated with depression in older adults, including comorbid physical disease, pain and disability, cognitive impairment, neuroticism, education level, loneliness and lack of social support.<sup>34–37</sup> Also, multiple studies have suggested that food insecurity is connected with poor mental health, especially depressive symptoms among older adults.<sup>38–40</sup>

The majority of the research articulates that food insecurity positively associates with depressive symptoms in older adults, and there is a dearth of studies in low-income and middle-income countries. This study aimed to examine the associations of specific indicators of food insecurity, including reduction in meal size, not eating food of one's choice, not eating enough food, remaining hungry for a whole day and body weight loss with depression among older adults in India. Furthermore, we analysed the association of food insecurity indicators after adjusting for socioeconomic and health attributes of older Indian adults with their depressive symptoms. Based on the conceptual framework provided (figure 1) and the past research, this study hypothesised that those who reduced meal size due to food shortage did not have food of own choice, remained hungry for a whole day or lost weight due to food shortage would be more likely to be depressed compared with those who did not experience these.

## DATA, VARIABLES AND METHODS

### Data source

This study uses data from India's first nationally representative longitudinal Ageing survey (Longitudinal Ageing

Study in India; LASI, 2017–2018), which investigates into the health, economics and social determinants and consequences of population ageing in India.<sup>41</sup> The present study was cross-sectional in nature. The representative sample included 72 250 individuals aged 45 and above and their spouses across all states and union territories of India except Sikkim. The LASI adopts a multistage stratified area probability cluster sampling design to select the eventual units of observation. Households with at least one member aged 45 and above were taken as the eventual unit of observation. This study provides scientific evidence on demographics, household economic status, chronic health conditions, symptom-based health condition, functional and mental health, biomarkers, health-care utilisation, work and employment, etc. It enables the cross-state analyses and the cross-national analyses of ageing, health, economic status and social behaviours and has been designed to evaluate the effect of changing policies and behavioural outcomes in India. The LASI was interviewer (face to face)-administered survey during household visits using computer-assisted personal interview (CAPI) technology. The interview was conducted in the local language of the area administered.<sup>41</sup> The total response rate at individual level was 95.6%. Detailed information on the sampling frame is available on the LASI wave-1 report.<sup>41</sup> The effective sample size for the present study was 31 464 older adults aged 60 years and above.<sup>41</sup>

## Variable description

### Outcome variable

The outcome variable for the study was major depression which was coded as 0 for 'not diagnosed with depression' and 1 for 'diagnosed with depression'.<sup>41</sup> Major depression among older adults with symptoms of dysphoria was calculated using the Short Form Composite International Diagnostic Interview (CIDI-SF) (Cronbach alpha: 0.70). Persons with a score of 3 or more were considered being depressed. This scale is used for probable psychiatric diagnosis of major depression and has been validated in field settings and widely used in population-based health surveys.<sup>42–43</sup>

### Explanatory variables

The explanatory variables were divided into three sections, namely, food security indicators, individual factors and household/community factors.

### Food security indicators

The food security indicators in the current study were adapted from similar items established in food security questionnaires of the US Household Food Security Survey Module adult scale,<sup>22</sup> and the items are validated in Indian settings.<sup>44</sup> The items are:

1. In the last 12 months, did you ever reduce the size of your meals or skip meals because there was not enough food at your household? The variable generated using this question was 'reduced the size of meals' and it was coded as 0 'no' and 1 'yes'.

2. In the last 12 months, did you eat enough food of your choice? Please exclude fasting/food-related restrictions due to religious or health-related reason. The variable generated using this question was 'did not eat enough food of once choice' and it was coded as 0 'no' and 1 'yes'.

In the last 12 months, were you hungry but did not eat because there was not enough food at your household? Please exclude fasting/food-related restrictions due to religious or health-related reasons. The variable generated using this question was 'hungry but did not eat' and it was coded as 0 'no' and 1 'yes'.

3. In the past 12 months, did you ever not eat for a whole day because there was not enough food at your household? Please exclude fasting/food-related restrictions due to religious or health-related reasons. The variable generated using this question was 'did not eat for a whole day' and it was coded as 0 'no' and 1 'yes'.
4. Do you think that you have lost weight in the last 12 months because there was not enough food at your household? The variable generated using this question was 'lost weight due to lack of food' as it was coded as 0 'no' and 1 'yes'.

#### Individual factors

1. Age was categorised as young old (60–69 years), old old (70–79 years) and oldest old (80+years).
2. Sex was categorised as male and female.
3. Educational status was categorised as no education/primary not completed, primary, secondary and higher.
4. Working status was categorised as currently working, not working/retired and never worked.
5. Social participation was categorised as no and yes. Social participation was measured through the question 'Are you a member of any of the organisations, religious groups, clubs or societies?' The response was categorised as no and yes.<sup>45</sup>
6. Self-rated health was coded as good, which includes excellent, very good and good whereas poor includes fair and poor.<sup>46</sup>
7. Difficulty in Activities of Daily Living (ADL) was coded as no and yes. ADL is a term used to refer to normal daily self-care activities (such as movement in bed, changing position from sitting to standing, feeding, bathing, dressing, grooming, personal hygiene, etc) The ability or inability to perform ADLs is used to measure a person's functional status, especially in the case of people with disabilities and the older adults.<sup>47 48</sup>
8. Difficulty in Instrumental ADL (IADL) was coded as no and yes. ADL are not necessarily related to fundamental functioning of a person, but they let an individual live independently in a community. The set ask were necessary for independent functioning in the community. Respondents were asked whether they were having any difficulties that were expected to last more than 3 months, such as preparing a hot meal, shopping for groceries, making a telephone call, taking medica-

tions, doing work around the house or garden, managing money (such as paying bills and keeping track of expenses) and getting around or finding an address in unfamiliar places.<sup>47 48</sup>

9. Morbidity status was categorised as 0 'no morbidity', 1 'any one morbid condition' and 2+ 'comorbidity'.<sup>49</sup>

#### Household/community factors

1. The monthly per capita consumption expenditure (MPCE) quintile was assessed using household consumption data. Sets of 11 and 29 questions on the expenditures on food and non-food items, respectively, were used to canvas the sample households. Food expenditure was collected based on a reference period of 7 days, and non-food expenditure was collected based on reference periods of 30 days and 365 days. Food and non-food expenditures have been standardised to the 30-day reference period. The MPCE is computed and used as the summary measure of consumption. The variable was then divided into five quintiles, that is, from poorest to richest.<sup>41</sup>
2. Religion was coded as Hindu, Muslim, Christian and Others.
3. Caste was recoded as Scheduled Tribe (ST), Scheduled Caste (SC), Other Backward Class (OBC) and others. The Scheduled Caste includes 'untouchables'; a group of the population that is socially segregated and financially/economically by their low status as per Hindu caste hierarchy. The SCs and STs are among the most disadvantaged socioeconomic groups in India. The OBC is the group of people who were identified as 'educationally, economically and socially backward'. The OBCs are considered low in the traditional caste hierarchy but are not considered untouchables. The 'other' caste category is identified as having higher social status.<sup>50</sup>
4. Place of residence was categorised as rural and urban.
5. The region was coded as North, Central, East, Northeast, West and South.<sup>51</sup>

#### Statistical analysis

In this study, descriptive statistics and bivariate analysis have been performed to determine the prevalence of major depression by food security factors along with individual and household factors.  $\chi^2$  test was used to check for intergroup differences in the prevalence of depression among older adults.<sup>52 53</sup> Furthermore, binary logistic regression analysis<sup>54</sup> was used to fulfil the aims and objective of the study. The results are presented in the form of OR with a 95% CI. There were two models in the present analysis. Model 1 represents the unadjusted OR (UOR). Model 2 represents the adjusted OR (AOR), that is, adjusted for individual (age, sex, education, living arrangements, work status, social participation, self-rated health, ADL/IADL difficulty and chronic morbidity) and household/community factors (wealth quintiles, religion, caste, place of residence and regions).

The equation for logistic regression is as follows:



$$\ln \left( \frac{P_i}{1-P_i} \right) = \beta_0 + \beta_1 x_1 + \dots + \beta_M x_{m-1}$$

where  $\beta_0, \dots, \beta_M$  are regression coefficients indicating the relative effect of a particular explanatory variable on the outcome variable. Variance inflation factor<sup>55–57</sup> was used to check multicollinearity among the variables used and it was found that there was no evidence of multicollinearity. Syvset command was used in STATA V.14<sup>58</sup> to account for complex survey design. Further, individual weights were used to make the estimates nationally representative.

## PATIENT AND PUBLIC INVOLVEMENT

No patient involved.

## RESULTS

### Sociodemographic and health profile of older adults

Table 1 depicts the socioeconomic profile of older adults in India. A proportion of 6.3% of the older adults reduced the size of meals due to lack of enough food in the household. About 40% of the older adults reported that they did not eat enough food of their choice. Of 5.6% of the older adults reported that they were hungry but did not eat because there was not enough food at their household. Of 4.2% of the older adults reported that they did not eat for a whole day because there was not enough food at their household. About 5.6% of the older adults think that they have lost weight due to lack of food at their household. Around 11% of the older adults belonged to oldest old age group; 68% of older adults had no education or their primary education was incomplete; 5.7% of older adults lived alone and 26.4% of older adults never worked in their lifetime. The share of older adults who had any social participation was 4.5%. Nearly 48.6% of older adults had poor self-rated health; 24.4% and 48.7% of older adults had difficulty in ADL and IADL, respectively; and 23.9% of older adults had two or more chronic diseases.

### Percentage of older adults suffering from major depression

Table 2 presents the share of older adults suffering from major depression in India. The overall prevalence of major depression was 8.4% among older adults in India. Higher percentage of older adults who reduced their size of meal suffered from major depression (23.6%) compared with those who did not reduce their meal (7.4%). Older adults who were hungry but did not eat had higher prevalence of major depression (25.3%) than those who had enough food (7.4%). Higher percentage of older adults who did not eat for a whole day suffered from major depression (24.8%) in compared with those who had food daily (7.7%). Older adults who lost their weight due to lack of food had higher prevalence of major depression (24.1%) in reference to their counterparts with no weight loss (7.5%).

## Logistic regression estimates of older adults suffering from major depression

Table 3 presents the results from logistic regression analysis of older adults suffering from major depression. Model 1 represents the unadjusted estimates, whereas model 2 represents the adjusted estimates. In model 1, older adults who reported to have reduced the size of meals due to lack of enough food had higher odds of being depressed in comparison to those who did not reduce the size of meals due to lack of enough food (UOR: 1.95, CI 1.61 to 2.37). Older adults who reported that they were hungry but did not eat because there was not enough food in their household had higher odds of being depressed in comparison to their counterparts with adequate food availability (UOR: 1.46, CI 1.16 to 1.85). Older adults who reported that they have lost weight due to lack of food had higher odds of being depressed compared with their counterparts with no weight loss (UOR: 2.17, CI 1.80 to 2.6).

Model 2 reveals that older adults who reported to have reduced the size of meals due to lack of enough food had higher odds of being depressed in comparison to those who did not reduce the size of meals due to lack of enough food (AOR: 1.76, CI 1.44 to 2.15). The choice of food did not have any significant association with major depression among older adults. Older adults who reported that they were hungry but did not eat because there was not enough food in their household had higher odds of being depressed compared with their counterparts with adequate food availability (AOR: 1.35, CI 1.06 to 1.72). Older adults who did not eat food for the whole day had higher odds of suffering from major depression in comparison to their counterparts who had food (AOR: 1.33; CI 1.03 to 1.71). Older adults who reported that they have lost weight due to lack of food had higher odds of being depressed compared with their counterparts with no weight loss (AOR: 1.57; CI 1.30 to 1.89).

The odds of major depression were higher among older women than men (AOR: 1.15; CI 1.02 to 1.28). The odds of major depression were lower among older adults who were living with children in comparison to those who were living with others (AOR: 0.80, CI 0.66 to 0.96). Older adults who had poor self-rated health had 2.38 times higher odds of suffering from major depression in comparison to older adults who had good self-rated health (AOR: 2.38, CI 2.15 to 2.64). The odds of major depression were significantly higher among the older adults who had difficulty in ADL and IADL in reference to the older adults who did not had difficulty in ADL and IADL, respectively (AOR: 1.56, CI 1.4 to 1.74) (AOR: 1.54, CI 1.38 to 1.72). Older adults who belonged to the poorest MPCE quintile had lower odds of suffering from major depression in comparison to older adults from richest wealth quintile (AOR: 0.75; CI 0.65 to 0.88). Older adults from urban areas had significantly lower odds of suffering from major depression compared with older adults who were from rural areas (AOR: 0.82; CI 0.73, to 0.92). Table S1 (online

**Table 1** Socioeconomic and health profile of older adults in India, 2017–2018

Background characteristics	Total		Not depressed		Depressed	
	Sample	Percentage	Sample	Percentage	Sample	Percentage
<b>Food security factors</b>						
Reduced the size of meals						
No	29 471	93.7	27 746	94.7	1784	82.2
Yes	1993	6.3	1548	5.3	386	17.8
Did not eat enough food of one's choice						
No	18 922	60.1	17 712	60.5	1229	56.6
Yes	12 542	39.9	11 582	39.5	941	43.4
Hungry but did not eat						
No	29 711	94.4	27 963	95.5	1806	83.2
Yes	1753	5.6	1331	4.5	364	16.8
Did not eat for a whole day						
No	30 152	95.8	28 291	96.6	1903	87.7
Yes	1312	4.2	1003	3.4	267	12.3
Lost weight due to lack of food						
No	29 695	94.4	27 928	95.3	1820	83.9
Yes	1769	5.6	1366	4.7	350	16.1
<b>Individual factors</b>						
Age						
Young old	18 410	58.5	17 165	58.6	1250	57.6
Old old	9501	30.2	8878	30.3	630	29.0
Oldest old	3553	11.3	3251	11.1	290	13.4
Sex						
Male	14 931	47.5	14 079	48.1	886	40.8
Female	16 533	52.6	15 215	51.9	1284	59.2
Education						
No education/primary not completed	21 381	68.0	19 710	67.3	1633	75.2
Primary completed	3520	11.2	3299	11.3	225	10.4
Secondary completed	4371	13.9	4186	14.3	208	9.6
Higher and above	2191	7.0	2098	7.2	105	4.8
Living arrangements						
Alone	1787	5.7	1576	5.4	194	8.9
With spouse	6397	20.3	5977	20.4	424	19.5
With children	21 475	68.3	20 102	68.6	1394	64.2
Others	1805	5.7	1639	5.6	158	7.3
Working status						
Working	9680	30.8	9079	31.0	613	28.3
Not working/retired	13 470	42.8	12 386	42.3	1054	48.6
Never worked	8314	26.4	7829	26.7	502	23.2
Social participation						
No	30 053	95.5	27 955	95.4	2093	96.5
Yes	1411	4.5	1339	4.6	77	3.5
Self-rated health*						
Good	15 850	51.4	16 108	55.0	603	27.8
Poor	14 961	48.6	13 186	45.0	1567	72.2

Continued

Table 1 Continued

Background characteristics	Total		Not depressed		Depressed	
	Sample	Percentage	Sample	Percentage	Sample	Percentage
Difficulty in ADL*						
No	23 802	75.7	22 594	77.1	1291	59.5
Yes	7662	24.4	6700	22.9	879	40.5
Difficulty in IADL*						
No	16 130	51.3	15 489	52.9	732	33.7
Yes	15 334	48.7	13 805	47.1	1438	66.3
Morbidity status						
0	14 773	47.0	13 981	47.7	835	38.5
1	9171	29.2	8540	29.2	632	29.1
2+	7520	23.9	6773	23.1	703	32.4
Household/community factors						
MPCE quintile						
Poorest	6829	21.7	6343	21.7	483	22.3
Poorer	6831	21.7	6411	21.9	430	19.8
Middle	6590	21.0	6174	21.1	424	19.5
Richer	6038	19.2	5613	19.2	424	19.5
Richest	5175	16.5	4753	16.2	409	18.9
Religion						
Hindu	25 871	82.2	24 091	82.2	1780	82.0
Muslim	3548	11.3	3286	11.2	259	12.0
Christian	900	2.9	850	2.9	52	2.4
Others	1145	3.6	1067	3.6	78	3.6
Caste						
Scheduled Caste	5949	18.9	5458	18.6	475	21.9
Scheduled Tribe	2556	8.1	2475	8.5	99	4.6
Other Backward Class	14 231	45.2	13 168	45.0	1048	48.3
Others	8729	27.7	8193	28.0	548	25.3
Place of residence						
Rural	22 196	70.6	20 446	69.8	1708	78.7
Urban	9268	29.5	8848	30.2	462	21.3
Region						
North	3960	12.6	3755	12.8	218	10.0
Central	6593	21.0	5759	19.7	761	35.1
East	7439	23.6	6951	23.7	492	22.7
Northeast	935	3.0	898	3.1	42	1.9
West	5401	17.2	5080	17.3	331	15.3
South	7136	22.7	6851	23.4	325	15.0
Total	31 464	100.0	29 294	100.0	2170	100.0

\*The sample is low due to missing cases and non-response.

ADL, Activities of Daily Living; IADL, Instrumental ADL; MPCE, monthly per capita *consumption* expenditure.

supplemental table S1) presents the regression estimates of older adults with food insecurity suffering from major depression. In model 1 (unadjusted), older adults with food insecurity had higher odds of major

depression in comparison to their food secure counterparts (UOR: 1.39; CI 1.27 to 1.51). Similarly, in the adjusted model (model 2), older adults with food insecurity had higher odds of major depression compared

**Table 2** Percentage of older adults suffering from major depression in India, 2017–2018

	Not depressed	Depressed	
Background characteristics	Percentage (n)	Percentage (n)	P value
Food security factors			
Reduced the size of meals			<0.001
No	92.6 (27 297)	7.4 (2174)	
Yes	76.4 (1523)	23.6 (470)	
Did not eat enough food of one’s choice			0.984
No	92.1 (17 425)	7.9 (1497)	
Yes	90.9 (11 395)	9.1 (1147)	
Hungry but did not eat			<0.001
No	92.6 (27 511)	7.4 (2200)	
Yes	74.7 (1309)	25.3 (444)	
Did not eat for a whole day			<0.001
No	92.3 (27 833)	7.7 (2319)	
Yes	75.2 (987)	24.8 (325)	
Lost weight due to lack of food			<0.001
No	92.5 (27 477)	7.5 (2218)	
Yes	75.9 (1343)	24.1 (426)	
Individual factors			
Age			0.207
Young old	91.7 (16 887)	8.3 (1523)	
Old old	91.9 (8734)	8.1 (767)	
Oldest old	90 (3199)	10 (354)	
Sex			<0.001
Male	92.8 (13 851)	7.2 (1080)	
Female	90.5 (14 969)	9.5 (1564)	
Education			<0.001
No education/primary not completed	90.7 (19 392)	9.3 (1989)	
Primary completed	92.2 (3246)	7.8 (274)	
Secondary completed	94.2 (4119)	5.8 (253)	
Higher and above	94.2 (2064)	5.8 (127)	
Living arrangements			<0.001
Alone	86.8 (1551)	13.2 (236)	
With spouse	91.9 (5880)	8.1 (516)	
With children	92.1 (19 777)	7.9 (1698)	
Others	89.3 (1612)	10.7 (193)	
Working status			<0.001
Working	92.3 (8933)	7.7 (747)	
Not working/retired	90.5 (12 185)	9.5 (1284)	
Never worked	92.6 (7702)	7.4 (612)	
Social participation			<0.001
No	91.5 (27 503)	8.5 (2550)	
Yes	93.4 (1317)	6.6 (94)	
Self-rated health			<0.001
Good	95.6 (15 848)	4.4 (734)	
Poor	87.2 (12 973)	12.8 (1910)	

Continued

Table 2 Continued

Background characteristics	Not depressed	Depressed	P value
	Percentage (n)	Percentage (n)	
Difficulty in ADL			<0.001
No	93.4 (22 229)	6.6 (1573)	
Yes	86 (6591)	14 (1071)	
Difficulty in IADL			<0.001
No	94.5 (15 238)	5.5 (892)	
Yes	88.6 (13582)	11.4 (1752)	
Morbidity status			<0.001
0	93.1 (13 755)	6.9 (1017)	
1	91.6 (8402)	8.4 (770)	
2+	88.6 (6663)	11.4 (857)	
<b>Household/community factors</b>			
MPCE quintile			<0.001
Poorest	91.4 (6240)	8.6 (589)	
Poorer	92.3 (6308)	7.7 (524)	
Middle	92.2 (6074)	7.8 (516)	
Richer	91.5 (5522)	8.6 (516)	
Richest	90.4 (4676)	9.6 (499)	
Religion			<0.001
Hindu	91.6 (23 702)	8.4 (2169)	
Muslim	91.1 (3232)	8.9 (316)	
Christian	92.9 (837)	7.1 (64)	
Others	91.7 (1049)	8.3 (96)	
Caste			<0.001
Scheduled Caste	90.3 (5370)	9.7 (579)	
Scheduled Tribe	95.3 (2435)	4.7 (121)	
Other Backward Class	91 (12 955)	9 (1276)	
Others	92.4 (8061)	7.7 (668)	
Place of residence			<0.001
Rural	90.6 (20 116)	9.4 (2081)	
Urban	93.9 (8705)	6.1 (563)	
Region			<0.001
North	93.3 (3695)	6.7 (265)	
Central	85.9 (5666)	14.1 (927)	
East	91.9 (6839)	8.1 (600)	
Northeast	94.5 (884)	5.5 (51)	
West	92.5 (4997)	7.5 (404)	
South	94.4 (6740)	5.6 (397)	
<b>Total</b>	91.6 (28 820)	8.4 (2644)	

The estimates are weighted.

ADL, Activities of Daily Living; IADL, Instrumental ADL; MPCE, monthly per capita consumption expenditure.



**Table 3** Logistic regression estimates for older adults suffering from major depression in India, 2017–2018

Background characteristics	Model 1	Model 2
	UOR (95% CI)	AOR (95% CI)
<b>Food security factors</b>		
Reduced the size of meals		
No	Ref.	Ref.
Yes	1.95* (1.61 to 2.37)	1.76* (1.44 to 2.15)
Did not eat enough food of one's choice		
No	Ref.	Ref.
Yes	1.01 (0.92 to 1.10)	0.92 (0.84 to 1.02)
Hungry but did not eat		
No	Ref.	Ref.
Yes	1.46* (1.16 to 1.85)	1.35* (1.06 to 1.72)
Did not eat for a whole day		
No	Ref.	Ref.
Yes	1.15 (0.90 to 1.47)	1.33* (1.03 to 1.71)
Lost weight due to lack of food		
No	Ref.	Ref.
Yes	2.17* (1.80 to 2.6)	1.57* (1.3 to 1.89)
<b>Individual factors</b>		
Age		
Young old		Ref.
Old old		0.81* (0.73 to 0.91)
Oldest old		0.74* (0.63 to 0.86)
Sex		
Male		Ref.
Female		1.15* (1.02 to 1.28)
Education		
No education/primary not completed		1.03 (0.83 to 1.28)
Primary completed		1.08 (0.85 to 1.37)
Secondary completed		0.99 (0.78 to 1.24)
Higher and above		Ref.
Living arrangements		
Alone		1.05 (0.82 to 1.33)
With spouse		0.72* (0.59 to 0.89)
With children		0.8* (0.66 to 0.96)
Others		Ref.
Working status		
Working		Ref.
Not working/retired		0.99 (0.88 to 1.11)
Never worked		0.80* (0.69 to 0.93)
Social participation		
No		0.92 (0.74 to 1.14)
Yes		Ref.
Self-rated health		
Good		Ref.
Poor		2.38* (2.15 to 2.64)

Continued

**Table 3** Continued

Background characteristics	Model 1	Model 2
	UOR (95% CI)	AOR (95% CI)
Difficulty in ADL		
No		Ref.
Yes		1.56* (1.4 to 1.74)
Difficulty in IADL		
No		Ref.
Yes		1.54* (1.38 to 1.72)
Morbidity status		
0		Ref.
1		1.23* (1.09 to 1.37)
2+		1.56* (1.38 to 1.76)
Household/community factors		
MPCE quintile		
Poorest		0.75* (0.65 to 0.88)
Poorer		0.77* (0.67 to 0.89)
Middle		0.70* (0.6 to 0.81)
Richer		0.88 (0.76 to 1.01)
Richest		Ref.
Religion		
Hindu		Ref.
Muslim		1.06 (0.92 to 1.22)
Christian		0.93 (0.74 to 1.18)
Others		1.09 (0.88 to 1.36)
Caste		
Scheduled Caste		Ref.
Scheduled Tribe		0.58* (0.47 to 0.7)
Other Backward Class		1.11 (0.98 to 1.26)
Others		0.92 (0.8 to 1.06)
Place of residence		
Rural		Ref.
Urban		0.82* (0.73 to 0.92)
Region		
North		Ref.
Central		1.96* (1.67 to 2.29)
East		0.97 (0.83 to 1.14)
Northeast		0.62* (0.49 to 0.79)
West		1.09 (0.92 to 1.29)
South		0.65* (0.55 to 0.77)

Model 2 was adjusted for all the individual and household factors whereas model 1 represents the unadjusted estimates.

\*If  $p < 0.05$ .

ADL, Activities of Daily Living; AOR, adjusted OR; IADL, Instrumental ADL; MPCE, monthly per capita consumption expenditure; Ref, reference; UOR, unadjusted OR.

with their food secure counterparts (AOR: 2.56; CI 2.28 to 2.88).

## DISCUSSION

The current study aimed to explore the prevalence of specific indicators of food insecurity and associated depression in late life through descriptive and regression analyses of a large country-representative survey data. With the increasing age, because of the development of more physical disabilities, there is a higher probability of increased food insecurity.<sup>59</sup> Consistently, the results have shown a substantial proportion of older population reducing the size of their meals due to food shortage, not eating food of own choice, not eating enough food, remaining hungry for a whole day and losing their body weight. In line with the recent literature, we also observed strong positive associations between food insecurity indicators and depression even after adjusting for several sociodemographic and health variables.<sup>60–62</sup>

However, since food insecurity and depression are multifactorial in nature, the mechanisms through which food insecurity affects depression are not adequately understood. There is a growing body of literature suggesting that food insecurity may act as an environmental stressor that can lead to late-life depressive disorders.<sup>63–64</sup> Food insecurity is considered as a major source of anxiety and life stress in the psychological pathways, leading to the shame or concern about one's position in the society.<sup>61</sup> Similarly, financial constraints may enhance feelings of worry and anxiety about the food situation and asking for food that is considered to be socially unacceptable creates feelings of stress.<sup>28</sup> On the other hand, the link between nutritional deficiencies and depressive symptoms is well documented.<sup>65</sup> A couple of studies have shown poor nutrition and dietary imbalances leading to depression.<sup>66–67</sup> Also, burgeoning studies on nutritional psychiatry attempt to address the pathway of availability and accessibility of food and the onset and the severity of depressive disorders.<sup>65–68</sup> The findings suggest that addressing food at the population level that is often an overlooked issue in the context of mental health, especially among the older population, may contribute to better mental health and psychological well-being in an ageing population.

Reduced intake of food and weight loss being indicators of food insecurity was found to be significantly associated with the risk of depression among the older sample in this study, and the associations were stronger than other indicators. This was consistent with several previous studies that had identified less nutrient intakes and to be associated with poor mental health and depressive symptoms.<sup>69–71</sup> The mechanism of this contribution can be explained by the relationship between stressors emerging as a result of reduced body weight and morbidities and negative mental health outcomes.<sup>72</sup> The inverse causation also has been discussed in past studies, suggesting that weight changes in both direction increased or decreased

intake may be caused by the appetite changes due to depression.<sup>73</sup> Seeking alternative food sources through food support plans, food banks or social networks is challenging for older adults due to social isolation, loss of independence and weakness, increasing with age,<sup>74</sup> in turn, they may feel particularly incapable when faced with food insecurity, probably raising the likelihood of depression,<sup>75</sup> suggesting the bidirectional association of food insecurity and depressive symptoms in old age. Our findings also suggest that promoting food security should be regarded as an important aspect of preventing psychological morbidity among the older population who are food insecure. Besides, the findings imply that food security interventions can have both nutritional and non-nutritional impacts, including an improved mental health status.

Other important findings of the current study include significant age and sex differences in the prevalence of major depressive disorder. The odds of suffering from depression were higher among young and female older adults compared with their male and oldest counterparts. The previous studies have also reported that depression is more prevalent among women, as the global depression ratio stands in favour of men.<sup>76</sup> Several studies elaborate that the gender disparity may stem primarily from behavioural as well as socioeconomic differences such as diet, tobacco and alcohol use and education.<sup>77</sup> Some studies reported that the higher prevalence of depression among women can be related to their biological conditions such as menstrual disorders, postmenopausal depression and anxiety and postpartum depression.<sup>76</sup> Furthermore, older people by increasing age tend to accept ill health as an impact of ageing and are less likely to be worried about their poor mental status.<sup>78</sup> On the other side, younger population becomes more aware of their psychological problems and is more sensitive towards any deficit in their mental well-being. The declining trend in depression in older ages was also reported in multiple previous studies.<sup>79–80</sup>

However, the findings of the current study need to be interpreted in light of major limitations. First, the study was conducted with a cross-sectional design, hence causality cannot be inferred between outcome variable that is depression and predictor variables. Also, the food security indicators were self-reported by older adults, which may result in their recall or reporting biases. Similarly, the huge variations in the prevalence of food insecurity measured through several indicators that vary from 4.2% for the question regarding 'did not eat for a whole day due to food shortage' to 39.9% for the question regarding 'did not have food of own choice' may result in misclassification effects, for example, people who are food secure might be marked as food insecure in specific indicator, which suggests the need for further investigation of appropriate measures of food security in Indian setting. However, due to the use of a large country representative data set with detailed measures of food insecurity, results can be generalisable to the broader

population. In addition, depression was assessed with a globally accepted scale of CIDI-SF that adds to the validity and reliability of the present study. For older adults, food insecurity is a vital psychosocial stressor that adds to variations in major depression across socioeconomic strata.<sup>81</sup>

A majority of research considered food insecurity as a static experience; however, both life transitions and cumulative experiences could also influence depression in old age.<sup>82–83</sup> Importantly, older adults may be significantly exposed to the consequence of food insecurity. For instance, some evidence indicates that food insecurity is more prone to poor diet condition among older adults than younger age groups.<sup>81–84</sup> In turn, poor diet quality is associated with depression, potentially a source of chronic systemic inflammation.<sup>85</sup> Food insecurity also intensifies the medical conditions common in older age, like diabetes, poor health status and medical morbidity that are recognised as risk factors for older age depression.<sup>86–88</sup> These aspects suggest future investigation of several pathways of food insecurity, including adverse childhood and adulthood exposures, leading to mental stressors and late-life depression.

## CONCLUSION

The results showed that self-reported food insecurity indicators, including the reduction in the size of meals, not eating food of one's choice, not eating enough food, remaining hungry for a whole day and losing the body weight, are strongly associated with major depression among older Indian adults. The findings can be of special interest to health-decision makers and researchers involved in the areas of mental health of ageing population in India and other low-income and middle-income countries with similar demographic and economic transitional stages. The findings suggest that the national food security programmes should be enhanced as an effort to improve mental health status and quality of life among older population.

**Contributors** Conceived and designed the research paper: MT and SS; analysed the data: SS; contributed agents/materials/analysis tools: MT; wrote the manuscript: KMS, MT and DD; refined the manuscript and guarantor: MT and SS.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not applicable.

**Ethics approval** The Indian Council of Medical Research (ICMR) extended the necessary guidance and ethical approval for conducting the LASI.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data may be obtained from a third party and are not publicly available. <https://www.ipsindia.ac.in/content/lasi-wave-i>.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content

includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Muhammad T <http://orcid.org/0000-0003-1486-7038>

Shobhit Srivastava <http://orcid.org/0000-0002-7138-4916>

## REFERENCES

- 1 Cafiero C, Viviani S, Nord M. Food security measurement in a global context: the food insecurity experience scale. *Measurement* 2018;116:146–52.
- 2 FAO. *Rome Draft Declaration of the World Summit on Food Security. World Food Summit, 16–18 of November 2009. World Food Summit, 2009*: 1–7.
- 3 FAO, IFAD, UNICEF, WFP W. *The state of food security and nutrition in the world 2017. building resilience for peace and food security [Policy support and Governance]*. Food and Agriculture Organization of the United Nations.
- 4 UN. *Transforming our world: the 2030 agenda for sustainable development* 2016.
- 5 Castetbon K. *Measuring food insecurity*. Sustain Nutr Chang World, 2017: 35–41.
- 6 Webb P, Coates J, Frongillo EA. Advances in Developing Country Food Insecurity Measurement Measuring Household Food Insecurity : Why It's So Important and Yet So. *Adv Dev Ctry Food Insecurity Meas* 2006;136:1404S–8.
- 7 Gardner BL, Rausser GC. *Handbook of agricultural economics*. 2B. 1st Edition. First. North Holland, 2002.
- 8 McCorriston S, Hemming DJ, Lamontagne-Godwin JD. What is the evidence of the impact of agricultural trade liberalisation on food security in developing countries? A systematic review 2013.
- 9 Aurino E, Morrow V. "Food prices were high, and the dal became watery". Mixed-method evidence on household food insecurity and children's diets in India. *World Dev* 2018;111:211–24.
- 10 Alderman H, Gentilini U, Yemtsov R. *The 1.5 billion people question: food, vouchers, or cash transfers?* Washington, DC: World Bank, 2018.
- 11 Narayanan S, Gerber N. Social safety nets for food and nutrition security in India. *Glob Food Sec* 2017;15:65–76.
- 12 Holben DH, American Dietetic Association. Position of the American dietetic association: food insecurity in the United States. *J Am Diet Assoc* 2010;110:1368–77.
- 13 Vozoris NT, Tarasuk VS. Household food insufficiency is associated with poorer health. *J Nutr* 2003;133:120–6.
- 14 Nelson K, Cunningham W, Andersen R, et al. Is food insufficiency associated with health status and health care utilization among adults with diabetes? *J Gen Intern Med* 2001;16:404–11.
- 15 Siefert K, Heflin CM, Corcoran ME, et al. Food insufficiency and the physical and mental health of low-income women. *Women Health* 2001;32:159–77.
- 16 Stuff JE, Casey PH, Szeto KL, et al. Household food insecurity is associated with adult health status. *J Nutr* 2004;134:2330–5.
- 17 Vilar-Compte M, Gaitán-Rossi P, Pérez-Escamilla R. Food insecurity measurement among older adults: implications for policy and food security governance. *Glob Food Sec* 2017;14:87–95.
- 18 Sharkey JR, Schoenberg NE. Prospective study of black-white differences in food insufficiency among homebound elders. *J Aging Health* 2005;17:507–27.
- 19 Nord M, Kantor LS. Seasonal variation in food insecurity is associated with heating and cooling costs among low-income elderly Americans. *J Nutr* 2006;136:2939–44.
- 20 Lee JS, Sinnott S, Bengtson R, et al. Unmet needs for the older Americans act nutrition program. *J Appl Gerontol* 2011;30:587–606.
- 21 Chung WT, Gallo WT, Giunta N, et al. Linking neighborhood characteristics to food insecurity in older adults: the role of perceived safety, social cohesion, and walkability. *J Urban Health* 2012;89:407–18.
- 22 Lee JS, Johnson MA, Brown A, et al. Food security of older adults requesting older Americans act nutrition program in Georgia can



- be validly measured using a short form of the U.S. household food security survey module. *J Nutr* 2011;141:1362–8.
- 23 Dean WR, Sharkey JR. Food insecurity, social capital and perceived personal disparity in a predominantly rural region of Texas: an individual-level analysis. *Soc Sci Med* 2011;72:1454–62.
  - 24 Bhargava V, Lee JS, Jain R, et al. Food insecurity is negatively associated with home health and out-of-pocket expenditures in older adults. *J Nutr* 2012;142:1888–95.
  - 25 Bengler R, Sinnett S, Johnson T, et al. Food insecurity is associated with cost-related medication non-adherence in community-dwelling, low-income older adults in Georgia. *J Nutr Elder* 2010;29:170–91.
  - 26 Holben DH, Barnett MA, Holcomb JP. Food insecurity is associated with health status of older adults participating in the commodity supplemental food program in a rural appalachian Ohio County. *J Hunger Environ Nutr* 2007;1:89–99.
  - 27 Gao X, Scott T, Falcon LM, et al. Food insecurity and cognitive function in Puerto Rican adults. *Am J Clin Nutr* 2009;89:1197–203.
  - 28 Kim K, Frongillo EA. Participation in food assistance programs modifies the relation of food insecurity with weight and depression in elders. *J Nutr* 2007;137:1005–10.
  - 29 WHO. *Mental health and development: targeting people with mental health conditions as a vulnerable group*, 2010: 1–6.
  - 30 Murata C, Kondo K, Hirai H, et al. Association between depression and socio-economic status among community-dwelling elderly in Japan: the Aichi Gerontological evaluation study (AGEs). *Health Place* 2008;14:406–14.
  - 31 Barua A, Ghosh MK, Kar N, et al. Socio-Demographic factors of geriatric depression. *Indian J Psychol Med* 2010;32:87–92.
  - 32 Copeland JR, Beekman AT, Dewey ME, et al. Depression in Europe. geographical distribution among older people. *Br J Psychiatry* 1999;174:312–21.
  - 33 Dudek D, Rachel W, Cyranka K. Depression in older people. *Encycl Biomed Gerontol* 2019;165:500–5.
  - 34 Heikkinen R-L, Kauppinen M. Depressive symptoms in late life: a 10-year follow-up. *Arch Gerontol Geriatr* 2004;38:239–50.
  - 35 Golden J, Conroy RM, Bruce I, et al. Loneliness, social support networks, mood and wellbeing in community-dwelling elderly. *Int J Geriatr Psychiatry* 2009;24:694–700.
  - 36 Steunenberg B, Beekman ATF, Deeg DJH, et al. Personality predicts recurrence of late-life depression. *J Affect Disord* 2010;123:164–72.
  - 37 Chen C-M, Mullan J, Su Y-Y, et al. The longitudinal relationship between depressive symptoms and disability for older adults: a population-based study. *J Gerontol A Biol Sci Med Sci* 2012;67:1059–67.
  - 38 Lee JS, Frongillo EA. Factors associated with food insecurity among U.S. elderly persons: importance of functional impairments. *J Gerontol B Psychol Sci Soc Sci* 2001;56:S94–9.
  - 39 German L, Kahana C, Rosenfeld V, et al. Depressive symptoms are associated with food insufficiency and nutritional deficiencies in poor community-dwelling elderly people. *J Nutr Health Aging* 2011;15:3–8.
  - 40 Klesges LM, Pahor M, Shorr RI, et al. Financial difficulty in acquiring food among elderly disabled women: results from the women's health and aging study. *Am J Public Health* 2001;91:68–75.
  - 41 International Institute for Population Sciences (IIPS), NPHCE, MoHFW. *Longitudinal ageing study in India (LASI) wave 1*. Mumbai, India, 2020.
  - 42 Kessler RC, Üstün TB. The world mental health (WMH) survey initiative version of the world Health organization (who) composite international diagnostic interview (CIDI). *Int J Methods Psychiatry Res* 2004;13:93–121.
  - 43 Muhammad T, Meher T, Sekher TV. Association of elder abuse, crime victimhood and perceived neighbourhood safety with major depression among older adults in India: a cross-sectional study using data from the LASI baseline survey (2017–2018). *BMJ Open* 2021;11:e055625.
  - 44 Sethi V, Maitra C, Avula R, et al. Internal validity and reliability of experience-based household food insecurity scales in Indian settings. *Agric Food Secur* 2017;6:21.
  - 45 Srivastava S, Chauhan S, Muhammad T, et al. Older adults' psychological and subjective well-being as a function of household decision making role: Evidence from cross-sectional survey in India. *Clin Epidemiol Glob Health* 2021;10:100676.
  - 46 Muhammad T, Srivastava S. Tooth loss and associated self-rated health and psychological and subjective wellbeing among community-dwelling older adults: a cross-sectional study in India. *BMC Public Health* 2022;22:1–11.
  - 47 Srivastava S, Muhammad T. Violence and associated health outcomes among older adults in India: a gendered perspective. *SSM Popul Health* 2020;12:100702.
  - 48 Muhammad T, Srivastava S. Why rotational living is bad for older adults? Evidence from a cross-sectional study in India. *J Popul Ageing* 2020;31.
  - 49 Keetile M, Navaneetham K, Letamo G. Prevalence and correlates of multimorbidity among adults in Botswana: a cross-sectional study. *PLoS One* 2020;15:e0239334.
  - 50 Corsi DJ, Subramanian SV. Socioeconomic gradients and distribution of diabetes, hypertension, and obesity in India. *JAMA Netw Open* 2019;2:e190411.
  - 51 International Institute for Population Sciences (IIPS) and ICF. *National family health survey (NFHS-4, 2017)*: 199–249.
  - 52 McHugh ML. The chi-square test of independence. *Biochem Medica* 2012.
  - 53 Mchugh ML. The chi-square test of independence lessons in biostatistics. *Biochem Medica*.
  - 54 Osborne J, King JE. Binary Logistic Regression. In: *Best Practices in Quantitative Methods*. SAGE Publications, Inc, 2011: 358–84.
  - 55 Lewis-Beck M, Bryman A, Futing Liao T. Variance Inflation Factors. In: *The SAGE encyclopedia of social science research methods*, 2012.
  - 56 Miles J. Tolerance and Variance Inflation Factor. In: *Wiley StatsRef: Statistics Reference Online*, 2014.
  - 57 Vogt W. Variance Inflation Factor (VIF). In: *Dictionary of Statistics & Methodology*, 2015.
  - 58 StataCorp. *Stata statistical software: release 14*. 2015, 2015.
  - 59 Rafat R, Rezazadeh A, Arzhang P, et al. The association of food insecurity with sociodemographic factors and depression in the elderly population of Qarchak City – Iran. *NFS* 2021;51:114–24.
  - 60 Bishwajit G, Kota K, Buh A, et al. Self-Reported food insecurity and depression among the older population in South Africa. *Psych* 2019;2:34–43.
  - 61 Mesbah SF, Sulaiman N, Shariff ZM, et al. Does food insecurity contribute towards depression? A cross-sectional study among the urban elderly in Malaysia. *Int J Environ Res Public Health* 2020;17:3118.
  - 62 Smith L, Il Shin J, McDermott D, et al. Association between food insecurity and depression among older adults from low- and middle-income countries. *Depress Anxiety* 2021;38:439–46.
  - 63 Silverman J, Krieger J, Kiefer M, et al. The relationship between food insecurity and depression, diabetes distress and medication adherence among low-income patients with poorly-controlled diabetes. *J Gen Intern Med* 2015;30:1476–80.
  - 64 Montgomery AL, Ram U, Kumar R, et al. Maternal mortality in India: causes and healthcare service use based on a nationally representative survey. *PLoS One* 2014;9:e83331.
  - 65 Jacka FN. Nutritional psychiatry: where to next? *EBioMedicine* 2017;17:24–9.
  - 66 Popa TA, Ladea M. Nutrition and depression at the forefront of progress. *J Med Life* 2012;5:414–9.
  - 67 Rao TSS, Asha MR, Ramesh BN, et al. Understanding nutrition, depression and mental illnesses. *Indian J Psychiatry* 2008;50:77.
  - 68 LaChance LR, Ramsey D. Antidepressant foods: an evidence-based nutrient profiling system for depression. *World J Psychiatry* 2018;8:97–104.
  - 69 Yang YJ. Socio-Demographic characteristics, nutrient intakes and mental health status of older Korean adults depending on household food security: based on the 2008–2010 Korea National health and nutrition examination survey. *Korean J Community Nutr* 2015;20:30.
  - 70 Weaver LJ, Hadley C. Moving beyond hunger and nutrition: a systematic review of the evidence linking food insecurity and mental health in developing countries. *Ecol Food Nutr* 2009;48:263–84.
  - 71 Faulconbridge LF, Wadden TA, Berkowitz RI, et al. Changes in symptoms of depression with weight loss: results of a randomized trial. *Obesity* 2009;17:1009–16.
  - 72 Vahabi M, Schindel Martin L, Martin LS. Food security: who is being excluded? A case of older people with dementia in long-term care homes. *J Nutr Health Aging* 2014;18:685–91.
  - 73 Forman-Hoffman VL, Yankey JW, Hillis SL, et al. Weight and depressive symptoms in older adults: direction of influence? *J Gerontol B Psychol Sci Soc Sci* 2007;62:43–51.
  - 74 Fitzpatrick K, Greenhalgh-Stanley N, Ver Ploeg M. The impact of food Deserts on food insufficiency and SNAP participation among the elderly. *Am J Agric Econ* 2016;98:19–40.
  - 75 Davison TE, McCabe MP, Knight T, et al. Biopsychosocial factors related to depression in aged care residents. *J Affect Disord* 2012;142:290–6.
  - 76 Albert PR. Why is depression more prevalent in women? *J Psychiatry Neurosci* 2015;40:219–21.
  - 77 Verplaetse TL, Smith PH, Pittman BP. Associations of gender, smoking, and stress with transitions in major depression diagnoses. *Yale J Biol Med* 2016;89:123–9.



- 78 Cahoon CG. Depression in older adults. *Am J Nurs* 2012;112:22–30.
- 79 Bryant C. Anxiety and depression in old age: challenges in recognition and diagnosis. *Int Psychogeriatr* 2010;22:511–3.
- 80 Schaakxs R, Comijs HC, van der Mast RC, *et al*. Risk factors for depression: differential across age? *Am J Geriatr Psychiatry* 2017;25:966–77.
- 81 Bergmans RS, Wegryn-Jones R. Examining associations of food insecurity with major depression among older adults in the wake of the great recession. *Soc Sci Med* 2020;258:113033.
- 82 Murayama Y, Yamazaki S, Yamaguchi J, *et al*. Chronic stressors, stress coping and depressive tendencies among older adults. *Geriatr Gerontol Int* 2020;20:297–303.
- 83 Vergare MJ. Depression in the context of late-life transitions. In: *Bulletin of the Menninger clinic*, 1997: 240–8.
- 84 Bergmans RS, Berger LM, Palta M, *et al*. Participation in the supplemental nutrition assistance program and maternal depressive symptoms: moderation by program perception. *Soc Sci Med* 2018;197:1–8.
- 85 Bergmans RS, Malecki KM. The association of dietary inflammatory potential with depression and mental well-being among U.S. adults. *Prev Med* 2017;99:313–9.
- 86 Cole MG, Dendukuri N. Risk factors for depression among elderly community subjects: a systematic review and meta-analysis. *Am J Psychiatry* 2003;160:1147–56.
- 87 Seligman HK, Bolger AF, Guzman D, *et al*. Exhaustion of food budgets at month's end and hospital admissions for hypoglycemia. *Health Aff* 2014;33:116–23.
- 88 Bergmans RS, Sadler RC, Wolfson JA, *et al*. Moderation of the association between individual food security and poor mental health by the local food environment among adult residents of Flint, Michigan. *Health Equity* 2019;3:264–74.

**Table-S1.** Logistic regression estimates for older adults suffering from major depression in India, 2017-18

Background characteristic	Model-1	Model-2
	UOR (95% CI)	AOR (95% CI)
<b>Food insecurity</b>		
No	Ref.	Ref.
Yes	1.39* (1.27, 1.51)	2.56* (2.28, 2.88)

\*if  $p < 0.05$ ; Ref: Reference; UOR: Unadjusted odds ratio; AOR: Adjusted odds ratio; CI: Confidence interval. Model-2 was controlled for individual and household factors; Food insecurity variable was generated using the following five questions: -

1. In the last 12 months, did you ever reduce the size of your meals or skip meals because there was not enough food at your household?
2. In the last 12 months, did you eat enough food of your choice?
3. In the last 12 months, were you hungry but didn't eat because there was not enough food at your household?
4. In the past 12 months did you ever not eat for a whole day because there was not enough food at your household?
5. Do you think that you have lost weight in the last 12 months because there was not enough food at your household?

All the variables were coded as 0 "no" and 1 "yes". Then a summation score of 0-5 was generated using a egen command in STATA. Lastly, the variable was coded as 0 if the summation score was 0 and 1 if the summation score ranges from 1-5. The variable was hence categorized as food insecurity (no and yes).