

# BMJ Open Identifying non-communicable disease multimorbidity patterns and associated factors: a latent class analysis approach

Parul Puri ,<sup>1</sup> Shri Kant Singh,<sup>1</sup> Sanghamitra Pati<sup>2</sup>

**To cite:** Puri P, Singh SK, Pati S. Identifying non-communicable disease multimorbidity patterns and associated factors: a latent class analysis approach. *BMJ Open* 2022;**12**:e053981. doi:10.1136/bmjopen-2021-053981

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-053981>).

Received 01 June 2021  
Accepted 27 May 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Department of Survey Research and Data Analytics, International Institute for Population Sciences, Mumbai, Maharashtra, India

<sup>2</sup>Department of Health Research, Indian Council of Medical Research Chandrasekharapur, Bhubaneswar, Orissa, India

## Correspondence to

Parul Puri;  
[parulpuri93@gmail.com](mailto:parulpuri93@gmail.com)

## ABSTRACT

**Objective** In the absence of adequate nationally-representative empirical evidence on multimorbidity, the existing healthcare delivery system is not adequately oriented to cater to the growing needs of the older adult population. Therefore, the present study identifies frequently occurring multimorbidity patterns among older adults in India. Further, the study examines the linkages between the identified patterns and socioeconomic, demographic, lifestyle and anthropometric correlates.

**Design** The present findings rest on a large nationally-representative sample from a cross-sectional study.

**Setting and participants** The study used data on 58 975 older adults (45 years and older) from the Longitudinal Ageing Study in India, 2017–2018.

**Primary and secondary outcome measures** The study incorporated a list of 16 non-communicable diseases to identify commonly occurring patterns using latent class analysis. The study employed multinomial logistic regression models to assess the association between identified disease patterns with unit-level socioeconomic, demographic, lifestyle and anthropometric characteristics.

**Results** The present study demonstrates that older adults in the country can be segmented into six patterns: ‘relatively healthy’, ‘hypertension’, ‘gastrointestinal disorders–hypertension–musculoskeletal disorders’, ‘musculoskeletal disorders–hypertension–asthma’, ‘metabolic disorders’ and ‘complex cardiometabolic disorders’. Additionally, socioeconomic, demographic, lifestyle and anthropometric factors are significantly associated with one or more identified disease patterns.

**Conclusions** The identified classes ‘hypertension’, ‘metabolic disorders’ and ‘complex cardiometabolic disorders’ reflect three stages of cardiometabolic morbidity with hypertension as the first and ‘complex cardiometabolic disorders’ as the last stage of disease progression. This underscores the need for effective prevention strategies for high-risk hypertension group. Also, targeted interventions are essential to reduce the burden on the high-risk population and provide equitable health services at the community level.

## BACKGROUND

Estimates generated by the World Bank suggest that life expectancy at birth has increased by 28 years in the past six decades in

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The study provides empirical evidence on non-communicable disease patterns using a large nationally-representative sample of older adults.
- ⇒ The study explored emerging non-communicable multimorbidity classes using a probability-based classification method.
- ⇒ Non-communicable diseases included in the study are based on self-reporting, leading to misclassification bias.
- ⇒ Considering the geographical and contextual variations in disease burden and profiling, these results should be explicitly apprehended for the older adults in India.
- ⇒ The findings were based on a single round of a longitudinal survey; thus, no causality was tested.

India.<sup>1</sup> Despite hypothesising this increased life expectancy as a byproduct of improved living conditions and advancement in the healthcare infrastructure and health delivery services, the past two decades have been fettered by the increasing morbidity burden, with a preponderance of non-communicable diseases (NCD) in the country.<sup>2</sup> Despite the ever-growing healthcare sector, India is still not adequately positioned to deal with the mounting NCD prevalence. The major reason for this shortfall is that NCDs result from shared pathophysiologies and risk factors. Thus, they tend to interact and coexist, commonly referred to as associative morbidities.<sup>3</sup>

The vast majority of the existing morbidity research in India still explores various realms of single chronic conditions,<sup>4–6</sup> with scarce focus on the simultaneous occurrence of multiple morbid conditions, also known as multimorbidity.<sup>2 7–9</sup> Moreover, the studies on multimorbidity in India are based on one-dimensional indices like simple count or score, which suffers from the limitation of ill-defined cut-offs.<sup>8 10</sup> In addition to this ongoing debate, these crude measures specifically focus on quantity rather than

the nature of chronic conditions and how they interact. Consequently, the evidence generated by the method above (multimorbidity score) can estimate the multimorbidity burden but is not adequate to form policies or design interventions, majorly due to its deficiencies in understanding the complexities related to multimorbidity.<sup>2 9 11</sup> In the absence of this, it is challenging for a low resource economy like India to satisfy the multifaceted service requirements of the multimorbid older adults in the country.

Recent studies suggest harnessing the advantages of latent class analysis (LCA) to deal with this.<sup>12–14</sup> These studies propose that chronic NCDs are disproportionately distributed in any community setting. Therefore various subgroups of the population have varied healthcare priorities and needs.<sup>14–16</sup> Thus, the disease patterns identified through LCA can be used to reorient the existing healthcare delivery system from a disease-centred model to a more patient-centred approach.<sup>9</sup> Therefore, the present study aims to bridge this knowledge gap and identify frequently occurring NCD multimorbidity patterns among older adults in India. Further, the study would examine the linkages between socioeconomic, demographic, lifestyle and anthropometric factors and the identified non-communicable disease patterns.

## MATERIALS AND METHODS

### Data source, ethical consideration and sample size

The present study is based on the Longitudinal Ageing Study in India (LASI), 2017–2018. LASI was conducted under the stewardship of the Ministry of Health and Family Welfare, Government of India.<sup>17</sup> The survey included information on individuals aged 45 years or older and their spouses dwelling in the same household, unrelated to age.<sup>17</sup> LASI implemented a multistage stratified probability cluster sampling design to draw nationally representative data from 35 states/union territories (except Sikkim). A detailed account of the survey design and sample size can be seen elsewhere.<sup>17</sup>

LASI received ethical approval from the Indian Council of Medical Research and Institutional Review Board held at International Institute for Population Sciences (IIPS), India. Additional written informed consent was administered from all the study participants.<sup>17</sup> The data were downloaded from the public repository of LASI held at IIPS, Mumbai. The study used merged data sets (N=65900) containing information from the individual, biomarker and household data files. The final analysis is based on 58975 individuals aged 45 years or older obtained from this merged data set.

### Patient and public involvement

It was not appropriate or possible to involve patients or the public in our research's design, conduct, reporting or dissemination plans.

## Measures

The question commissioned to measure each disease based on self-reported medical diagnoses was, 'Has any health professional ever diagnosed you with the following chronic conditions or diseases?'. The list included information on 16 diseases, namely asthma, cancer, chronic bronchitis, chronic heart disease, chronic renal failure, chronic obstructive pulmonary disease, diabetes, gastrointestinal disorder, high cholesterol, hypertension, musculoskeletal disorder, neurological and psychiatric disorders, skin disease, stroke, thyroid disease and urinary incontinence. The diseases mentioned above were recoded as no (1) or yes (2) in accordance to the statistical package used.

The socioeconomic, demographic, lifestyle and anthropometric variables evaluated based on the existing literature included age (45–49/50–54/55–59/60–64/65–69/70–74/75–79/80 years or older), sex (men/women), residence (urban/rural), religion (Hindu/Muslim/Christian/Others), social group (Scheduled Caste (SC)/Scheduled Tribes (ST)/Other Backward Class (OBC)/Others), level of education (no education/less than 5 years/5–9 years/10 years or more), wealth status (poor/non-poor), current marital status (in union/not in a union), current working status (never worked/worked in the past but not currently working/currently work), tobacco consumption (never used tobacco/quit tobacco/currently consuming), ever consumed alcohol (no/yes), physical activity (physically active/physically inactive), body mass index (BMI) (underweight/normal/overweight/obese) and waist-hip ratio (WHR) (low risk/high risk).

Religion and social group (which comprises four major caste categories) are essential, as they build the social structure in the context of India. They play a decisive role in the social acceptance, cultural and dietary aspects of any individual. Four major social groups included in the study are SC, ST, OBC and Others (this comprises all other caste groups). Based on social stratification, SC, ST and OBC are considered socially disadvantaged, while the 'others' category mostly includes the socially accepted 'upper' castes.<sup>18</sup>

Physical activity was categorised into two categories: physically inactive and active. Physically inactive individuals were those who were not engaged in any moderate or vigorous physical activity throughout the week. On the contrary, physically active individuals satisfied one of the three criteria:

1. Only vigorous activities: Those who perform at least 75 min of vigorous-intensity physical activity throughout the week.
2. Only moderate activities: Those who perform at least 150 min of moderate-intensity physical activity throughout the week.
3. Both vigorous and moderate activities: Those who are engaged in a combination of moderate-intensity and vigorous-intensity activity.

LASI included information on weight, height, waist and hip circumference. We used BMI, calculated as weight (in kgs) divided by the square of height (in m<sup>2</sup>) and WHR, calculated as waist circumference (in cm) divided by hip circumference (in cm). BMI was categorised into four, namely underweight (BMI  $\leq 18.4$  kg/m<sup>2</sup>), normal ( $18.5 \text{ kg/m}^2 \leq \text{BMI} \leq 24.9 \text{ kg/m}^2$ ), overweight ( $25.0 \text{ kg/m}^2 \leq \text{BMI} \leq 29.9 \text{ kg/m}^2$ ) and obese (BMI  $\geq 30$  kg/m<sup>2</sup>). WHR was categorised into high and low risk after stratifying for the sex of the individual. Low risk included WHR  $< 0.90$  for men and WHR  $< 0.85$  for women, while high risk included WHR  $\geq 0.90$  for men and WHR  $\geq 0.85$  for women.

### Statistical analysis

We calculated descriptive statistics, including unweighted frequency and weighted percentages. In addition to this, disease profile, that is, the prevalence of all NCDs included in the study, was also reported. To identify the recurrent disease combinations, all possible two-way combinations ( $^{16}C_2=120$ ) were explored. Further, these were supplemented with  $\chi^2$  p values to identify statistically significant associations or associative morbidities.

LCA was used to identify frequently occurring disease patterns. LCA is a statistical procedure used to identify unobserved homogeneous (latent) subgroups within a population.<sup>19 20</sup> These subgroups are identified based on the categorical responses collected from the study population. An array of latent class models was fitted before selecting the optimal number of latent classes (say, *n* classes). The existing literature defines no definitive way of the best criteria; therefore, the most widely accepted statistical criteria were used.<sup>21</sup> This included reporting multiple fit statistics, along with parsimony and theoretical interpretability.<sup>12 15 19 20</sup> Thus, the selection criterion was based on multiple model fit indices, Akaike information criterion (AIC), Bayesian information criterion (BIC), adjusted Bayesian information criterion (aBIC), consistent Akaike information criterion (cAIC).<sup>12 13</sup> Parsimony and theoretical interpretability were preferred in identifying the optimum number (*n*) of disease classes. Furthermore,  $\chi^2$  goodness of fit, G2 statistics, entropy and likelihood ratio tests were reported in addition to the smallest class count and their respective percentage were also reported.<sup>19 22</sup>

These '*n*' classes were labelled based on the item-response probabilities of the selected chronic morbidities included in the study.<sup>12 13</sup> Existing literature suggests that item-response probabilities are similar to factor loadings.<sup>23</sup> Given this, it is recommended that a standard loading of 0.3 or above should be employed to define a particular factor.<sup>24–26</sup> Thus, a cut-off of 0.3 was chosen to assign labels. A particular item (or disease) with an item-response probability of 0.3 or above depicts a strong association with the identified latent pattern or class. Thus, this item was most informative and was used to assign labels to a specific latent class. Once the optimal number of latent classes was selected, the entire study population was

segregated into '*n*' classes (identified disease classes).<sup>12 14</sup> Finally, a multivariable multinomial logistic regression (posterior analysis) was performed to identify the factors associated with the multimorbidity patterns identified.

Analysis for this study was done using 'poLCA' package in RStudio V.1.1.463 (R Studio).<sup>21</sup> The study followed the Strengthening the Reporting of Observational Studies in Epidemiology reporting guideline (online supplemental file 1). All the estimates generated in the study were presented after suitable application of sampling weights provided by LASI, 2017–2018.<sup>17</sup>

## RESULTS

### Sample description and distribution of selected chronic conditions

The present analysis is based on 58975 individuals in age 45 years and above from LASI, wave-1, 2017–2018. Table 1 describes the study population, which comprised 19% of the individuals in the age group 45–49 years. Around 54% of the individuals were women, and 71% lived in rural areas. Around 82% followed the Hindu faith and 45% belonged to the OBC group. Fifty-one per cent received no education, and 74% were currently married. Around 62% never consumed tobacco, 85% were lifetime alcohol abstainers and 29% led a physically inactive lifestyle. Furthermore, 20% were overweight, 7% were obese and 77% had a high-risk WHR.

### Disease profile and association between selected NCDs

Figure 1 illustrates the disease profile, linkages and prevalence of all possible two-way combinations between selected NCDs among older adults in India. The values in the diagonal cells (coloured in light blue) represent the prevalence of all the diseases included in the study. Findings suggest that 14 diseases had a prevalence greater than 1%, whereas 11 had a prevalence of greater than 2%. Hypertension (26.9%), gastrointestinal disorders (18.4%), musculoskeletal disorders (16.2%), diabetes (11.7%) and skin diseases (5.2%) were commonly occurring among older adults in India.

The off-diagonal values represent the prevalence rates (in %) of all-possible two-way combinations. In addition, the colours yellow and green indicate the degree of association between the two diseases in any specific combinations. The study explored 120 diseases combinations ( $^{16}C_2=120$ ), out of which 110 were statistically significant and 20 had a prevalence of more than a per cent. The most prevalent disease combinations identified in the present study were 'diabetes-hypertension' (7.8%), 'musculoskeletal disorders-hypertension' (6.4%), 'gastrointestinal disorders-hypertension' (6.1%), 'gastrointestinal disorders-musculoskeletal disorders' (3.9%), 'musculoskeletal disorders-diabetes' (2.7%) and 'chronic heart disease- hypertension' (2.5%). Hypertension was present in five statistically significant associations (links); similarly, gastrointestinal disorders and diabetes were

**Table 1** Socio-demographic, lifestyle and anthropometric characteristic of the study population (45 years or older), Longitudinal Ageing Study in India, wave-1, 2017–2018

| Correlates             | Categories                               | Unweighted frequency (N=58 975) | Weighted percentage |
|------------------------|--|---------------------------------|---------------------|
| Age (in years)         | 45–49                                    | 11 917                          | 18.86               |
|                        | 50–54                                    | 9863                            | 15.88               |
|                        | 55–59                                    | 9057                            | 15.09               |
|                        | 60–64                                    | 9195                            | 15.39               |
|                        | 65–69                                    | 7981                            | 14.45               |
|                        | 70–74                                    | 5126                            | 9.45                |
|                        | 75–79                                    | 2990                            | 5.53                |
|                        | 80+                                      | 2846                            | 5.34                |
| Sex                    | Male                                     | 27 374                          | 46.21               |
|                        | Female                                   | 31 601                          | 53.79               |
| Place of residence     | Rural                                    | 38 715                          | 70.68               |
|                        | Urban                                    | 20 260                          | 29.32               |
| Religion               | Hindu                                    | 43 238                          | 82.45               |
|                        | Muslim                                   | 6968                            | 11.07               |
|                        | Christian                                | 5984                            | 2.99                |
|                        | Others                                   | 2785                            | 3.50                |
| Social group           | Scheduled Castes                         | 9959                            | 19.64               |
|                        | Scheduled Tribes                         | 10 348                          | 8.70                |
|                        | Other Backward Class                     | 22 243                          | 45.27               |
|                        | Others                                   | 16 425                          | 26.40               |
| Level of education     | No education                             | 27 766                          | 50.93               |
|                        | Less than 5 years                        | 10 959                          | 17.79               |
|                        | 5–9 years                                | 9380                            | 13.98               |
|                        | 10 years or more                         | 10 870                          | 17.30               |
| Wealth status          | Poor                                     | 23 537                          | 42.74               |
|                        | Non-poor                                 | 35 438                          | 57.26               |
| Current marital status | In union                                 | 44 427                          | 74.39               |
|                        | Not in union                             | 14 548                          | 25.61               |
| Current working status | Never worked                             | 16 115                          | 25.74               |
|                        | Worked in past but not currently working | 15 510                          | 27.15               |
|                        | Currently working                        | 27 350                          | 47.11               |
| Tobacco consumption    | Never used tobacco                       | 46 028                          | 61.90               |
|                        | Quit tobacco                             | 1471                            | 4.92                |
|                        | Currently consuming                      | 11 476                          | 33.18               |
| Ever consumed alcohol  | No                                       | 48 354                          | 84.81               |
|                        | Yes                                      | 10 621                          | 15.19               |
| Physical activity      | Physically active                        | 41 295                          | 71.27               |
|                        | Physically inactive                      | 17 680                          | 28.73               |
| Body mass index        | Underweight                              | 10 822                          | 21.39               |
|                        | Normal                                   | 30 529                          | 51.31               |
|                        | Overweight                               | 12 651                          | 19.62               |
|                        | Obese                                    | 4298                            | 6.69                |
|                        | Do not know/Not measured                 | 675                             | 1.00                |

Continued



Table 1 Continued

| Correlates      | Categories | Unweighted frequency (N=58975) | Weighted percentage |
|-----------------|------------|--------------------------------|---------------------|
| Waist-hip ratio | Low risk   | 11 763                         | 22.72               |
|                 | High risk  | 47 212                         | 77.28               |

present with four links, making them the most interactive diseases.

### Latent classes of multimorbidity

Table 2 illustrates the LCA model fit results. AIC, BIC, cAIC and aBIC decreased up to the six-class model. Class 7, 8 and 9 models were not well identified, that is, they could not attain a maximum likelihood even after 10 000 iterations. A six-class model was selected as optimum based on the lowest BIC value (in addition, all other fit indices were in agreement) along with model's theoretical interpretability. Furthermore, a six-class model reported an acceptable level of entropy (entropy=0.62). The selected model had a minimum class count of 457, covering 0.78% of the study sample.

The results from class proportion and item-response probabilities ( $p$ ) are presented in table 3. These estimated item-response probabilities ( $p \geq 0.3$ ) were employed to assign labels to the six-class model identified. Class 1 comprised individuals with low probabilities for all 16 NCDs compared with other classes. This group comprised 65% of the study population and was labelled as *relatively healthy*. Multimorbidity prevalence in this group was 3.4%, and the average number of chronic conditions was 0.4.

Class 2 comprised 23.6% of the study population was labelled as *hypertension* and reported high probabilities

(0.76) for hypertension. Multimorbidity prevalence in this group was 61.76%, and the average number of chronic conditions was 1.8.

Class 3 comprised 6.6% of the study population was labelled as *gastrointestinal disorders–hypertension–musculoskeletal disorders* and reported high probabilities for gastrointestinal disorders (0.53), hypertension (0.35) and musculoskeletal disorders (0.30). Multimorbidity prevalence in this group was 100.0%, and the average number of chronic conditions was 2.8.

Class 4 comprised 0.8% of the study population and was labelled as *musculoskeletal disorders–hypertension–asthma* and reported high probabilities for musculoskeletal disorders (0.30), hypertension (0.33) and asthma (0.30). Multimorbidity prevalence in this group was 100.0%, and the average number of chronic conditions was 3.0.

Class 5 comprised 2.6% of the study population and was labelled as *metabolic disorders* and reported high probabilities for hypertension (0.75), high cholesterol (0.57) and diabetes (0.48). Multimorbidity prevalence in this group was 100.0%, and the average number of chronic conditions was 3.2.

Class 6 comprised 1.5 % of the study population and was labelled as *complex cardiometabolic disorders* and reported high probabilities for hypertension (0.85), along with

|      | AS   | MKS   | CA   | CB   | CRF  | COPD | DM    | GD    | CHD  | HC   | HYP   | NPD  | SD   | ST   | THY  | UI   |
|------|------|-------|------|------|------|------|-------|-------|------|------|-------|------|------|------|------|------|
| AS   | 4.69 |       |      |      |      |      |       |       |      |      |       |      |      |      |      |      |
| MKS  | 1.26 | 16.23 |      |      |      |      |       |       |      |      |       |      |      |      |      |      |
| CA   | 0.03 | 0.11  | 0.58 |      |      |      |       |       |      |      |       |      |      |      |      |      |
| CB   | 0.23 | 0.48  | 0.58 | 1.19 |      |      |       |       |      |      |       |      |      |      |      |      |
| CRF  | 0.05 | 0.18  | 0.01 | 0.02 | 0.68 |      |       |       |      |      |       |      |      |      |      |      |
| COPD | 0.30 | 0.52  | 0.02 | 0.31 | 0.04 | 1.34 |       |       |      |      |       |      |      |      |      |      |
| DM   | 0.70 | 2.73  | 0.12 | 0.38 | 0.14 | 0.39 | 11.74 |       |      |      |       |      |      |      |      |      |
| GD   | 1.02 | 3.88  | 0.15 | 0.26 | 0.23 | 0.27 | 2.22  | 18.40 |      |      |       |      |      |      |      |      |
| CHD  | 0.32 | 1.11  | 0.04 | 0.28 | 0.06 | 0.31 | 1.24  | 0.85  | 3.86 |      |       |      |      |      |      |      |
| HC   | 0.13 | 0.63  | 0.04 | 0.04 | 0.04 | 0.05 | 0.70  | 0.56  | 0.40 | 2.32 |       |      |      |      |      |      |
| HYP  | 1.68 | 6.40  | 0.26 | 0.63 | 0.31 | 0.69 | 7.83  | 6.05  | 2.49 | 1.47 | 26.91 |      |      |      |      |      |
| NPD  | 0.21 | 0.92  | 0.03 | 0.06 | 0.04 | 0.08 | 0.37  | 0.65  | 0.21 | 0.18 | 0.99  | 2.35 |      |      |      |      |
| SD   | 0.36 | 1.14  | 0.03 | 0.07 | 0.06 | 0.09 | 0.81  | 1.54  | 0.18 | 0.19 | 1.72  | 0.17 | 5.21 |      |      |      |
| ST   | 0.11 | 0.49  | 0.03 | 0.02 | 0.04 | 0.04 | 0.49  | 0.43  | 0.24 | 0.09 | 1.18  | 0.18 | 0.13 | 1.78 |      |      |
| THY  | 0.23 | 0.79  | 0.05 | 0.09 | 0.07 | 0.10 | 0.71  | 0.73  | 0.19 | 0.32 | 1.47  | 0.21 | 0.24 | 0.07 | 2.75 |      |
| UI   | 0.29 | 0.89  | 0.04 | 0.09 | 0.12 | 0.12 | 0.62  | 1.19  | 0.19 | 0.14 | 1.18  | 0.19 | 0.28 | 0.16 | 0.19 | 2.96 |

$p < 0.01$

$p < 0.05$

**Figure 1** Disease profile and association between all possible two-way combinations of selected non-communicable diseases among older adults, Longitudinal Ageing Study in India, wave-1, 2017–2018. AS, asthma; CA, cancer; CB, chronic bronchitis; CHD, chronic heart disease; COPD, chronic obstructive pulmonary disorder; CRF, chronic renal failure; DM, diabetes; GD, gastrointestinal disorders; HC, high cholesterol; HYP, hypertension; MKS, musculoskeletal disorder; NPD, neurological and psychiatric disorders; SD, skin disease; ST, stroke; THY, thyroid disorder; UI, urinary incontinence.

**Table 2** Model fit and diagnostic criteria for latent class analyses, Longitudinal Ageing Study in India, wave-1, 2017–2018

| Class | df                  | LL         | AIC      | BIC      | cAIC     | aBIC      | Smallest class count (n) | Smallest class size (%) | G2     | X <sup>2</sup> goodness of fit | Likelihood ratio test |                  |                |
|-------|---------------------|------------|----------|----------|----------|-----------|--------------------------|-------------------------|--------|--------------------------------|-----------------------|------------------|----------------|
|       |                     |            |          |          |          |           |                          |                         |        |                                | Entropy               | Likelihood ratio | Critical value |
| 2     | 58942               | -182 959.3 | 365984.7 | 366281.2 | 366314.2 | 366 176.3 | 11 895                   | 20.17                   | 6347.2 | 6 096 559                      | 0.5817                | -                | -              |
| 3     | 58925               | -182 347.1 | 364794.1 | 365243.4 | 365293.4 | 365 084.5 | 3774                     | 6.40                    | 5122.6 | 1 779 553                      | 0.6215                | 1224.4           | 27.587         |
| 4     | 58908               | -182 074.9 | 364283.9 | 364885.9 | 364952.9 | 364 672.9 | 1109                     | 1.88                    | 4578.4 | 227 023.2                      | 0.5453                | 544.4            | 27.587         |
| 5     | 58891               | -181 959.9 | 364087.7 | 364842.4 | 364926.4 | 364 575.5 | 590                      | 1.00                    | 4348.2 | 295 466.3                      | 0.5812                | 230.0            | 27.587         |
| 6     | 58874               | -181 853.5 | 363909.0 | 364816.5 | 364917.5 | 364 495.5 | 457                      | 0.78                    | 4135.5 | 126 168                        | 0.6158                | -                | -              |
| 7     | Not well identified |            |          |          |          |           |                          |                         |        |                                |                       |                  |                |
| 8     | Not well identified |            |          |          |          |           |                          |                         |        |                                |                       |                  |                |
| 9     | Not well identified |            |          |          |          |           |                          |                         |        |                                |                       |                  |                |

aBIC, Adjusted Bayesian Information Criteria; AIC, Akaike Information Criteria; BIC, Bayesian Information Criteria; cAIC, Consistent Akaike Information Criteria; df, degree of freedom; LL, log likelihood.

musculoskeletal disorders (0.51), diabetes (0.50), gastrointestinal disorders (0.43), high cholesterol (0.28) and heart disease (0.27). It is worth mentioning that item-response probabilities for hypertension, musculoskeletal disorders, diabetes and gastrointestinal disorders are higher than the prescribed cut-off of 0.3. However, the item-response probabilities for high cholesterol (0.28) and chronic heart disease (0.27) can be considered as 0.3 if rounded off to one decimal. Multimorbidity prevalence in this group was 100.0%, and the average number of chronic conditions was 5.1. A detailed description of the socio-demographic, lifestyle and anthropometric characteristics of six identified latent classes is presented in online supplemental file 2.

### Posterior analysis

Table 4 depicts multinomial logistic regression analysis findings considering *relatively healthy* class as the base outcome. The findings suggest that higher age increased the likelihood of belonging to any of the five multimorbidity classes. In particular, being woman increased the likelihood of belonging to *hypertension, gastrointestinal disorders–hypertension–musculoskeletal disorders, metabolic disorders and complex cardiometabolic disorders* class. Residing in the urban areas increased the likelihood of belonging to class: *hypertension, metabolic disorders and complex cardiometabolic disorders* compared with class *relatively healthy*. The likelihood of belonging to class: *hypertension, gastrointestinal disorders–hypertension–musculoskeletal disorders, metabolic disorders and complex cardiometabolic disorders* was higher for respondents belonging to the Muslim faith compared with the *relatively healthy* class.

Belonging to the ST decreased the likelihood of belonging to class *gastrointestinal disorders–hypertension–musculoskeletal disorders, musculoskeletal disorders–hypertension–asthma, metabolic disorders and complex cardiometabolic disorders* compared with *relatively healthy* class.

In particular, the educated respondents were associated with a higher likelihood of belonging to class *hypertension, metabolic disorders and complex cardiometabolic disorders* than respondents from a *relatively healthy* class. Respondents belonging to the wealthy class were more likely to belong to any of the five multimorbidity class than the *relatively healthy* class.

The findings suggest that currently working was associated with a lower likelihood of belonging to *hypertension, gastrointestinal disorders–hypertension–musculoskeletal disorders, musculoskeletal disorders–hypertension–asthma and metabolic disorders*.

Considering the lifestyle and anthropometric predictors, individuals who have quit smoking were more likely to belong to any of the five multimorbid classes as compared with *relatively healthy*. Consumption of alcohol increased the likelihood of belonging to the class *metabolic disorders*. Overweight increased the likelihood of belonging to class *hypertension, gastrointestinal disorders–hypertension–musculoskeletal disorders, metabolic disorders and complex cardiometabolic disorders* compared with the *relatively healthy* class.

**Table 3** Class proportions and item-response probabilities from six-latent class model of chronic morbidities, Longitudinal Ageing Study in India, wave-1, 2017–2018

| Latent class   | 1                  | 2            | 3   | 4   | 5                   | 6                                 |
|--|--------------------|--------------|---|---|---------------------|-----------------------------------|
| Assigned labels  | Relatively healthy | Hypertension | Gastrointestinal disorders–hypertension–musculoskeletal disorders | Musculoskeletal disorders–hypertension–asthma | Metabolic disorders | Complex cardiometabolic disorders |
| Class proportion   | 65.03              | 23.59        | 6.57  | 0.78  | 2.58                | 1.45                              |
| Non-communicable diseases  |                    |              |   |   |                     |                                   |
| Asthma   | 0.02               | 0.03         | 0.06  | <b>0.30</b>                                   | 0.03                | 0.12                              |
| Cancer   | 0.00               | 0.01         | 0.01  | 0.01  | 0.01                | 0.03                              |
| Chronic bronchitis   | 0.00               | 0.00         | 0.01  | 0.14  | 0.01                | 0.05                              |
| Chronic heart disease  | 0.01               | 0.06         | 0.03  | 0.11  | 0.18                | <b>0.27</b>                       |
| Chronic renal disease  | 0.00               | 0.01         | 0.02  | 0.02  | 0.01                | 0.06                              |
| Chronic obstructive pulmonary disease  | 0.00               | 0.01         | 0.01  | 0.18  | 0.01                | 0.06                              |
| Diabetes   | 0.04               | 0.28         | 0.10  | 0.04  | <b>0.48</b>         | <b>0.50</b>                       |
| Gastrointestinal disorder  | 0.12               | 0.16         | <b>0.53</b>   | 0.19  | 0.19                | <b>0.43</b>                       |
| High cholesterol   | 0.00               | 0.00         | 0.02  | 0.03  | <b>0.57</b>         | <b>0.28</b>                       |
| Hypertension   | 0.04               | <b>0.76</b>  | <b>0.35</b>   | <b>0.33</b>                                   | <b>0.75</b>         | <b>0.85</b>                       |
| Musculoskeletal disorders  | 0.08               | 0.16         | <b>0.30</b>   | <b>0.34</b>                                   | 0.19                | <b>0.51</b>                       |
| Neurological and psychiatric disorder  | 0.01               | 0.02         | 0.05  | 0.08  | 0.01                | 0.16                              |
| Skin disorder  | 0.03               | 0.03         | 0.16  | 0.07  | 0.05                | 0.13                              |
| Stroke   | 0.00               | 0.04         | 0.01  | 0.01  | 0.03                | 0.15                              |
| Thyroid disorder   | 0.01               | 0.03         | 0.05  | 0.03  | 0.14                | 0.17                              |
| Urinary incontinence   | 0.01               | 0.02         | 0.11  | 0.06  | 0.01                | 0.19                              |
| Multimorbidity (in %)  | 3.44               | 61.76        | 100.00  | 100.00  | 100.00              | 100.00                            |
| Average number of diseases   | 0.38               | 1.82         | 2.75  | 3.00  | 3.20                | 5.12                              |
| The bold values depict the diseases which are used in assigning labels to the identified latent classes. |                    |              |   |   |                     |                                   |

*healthy* class. Obesity increased the likelihood of belonging to class any of the five multimorbid classes compared with the *relatively healthy* class. Furthermore, being physically inactive increased the likelihood of belonging to class *hypertension* and *complex cardiometabolic disorders*. High-risk WHR increased the likelihood of belonging to class *hypertension*, *gastrointestinal disorders–hypertension–musculoskeletal disorders*, *metabolic disorders* and *complex cardiometabolic disorders* compared with *relatively healthy* class.

## DISCUSSION

This study used LCA to investigate the emerging multimorbidity patterns and identify its predictors among older adults in India. Based on a nationally representative sample of 58 975 individuals, the study included information on 16 NCDs.

LCA identified six multimorbidity classes, including a predominant *relatively healthy* (65.0%) class, *hypertension*

(23.6%), *gastrointestinal disorders–hypertension–musculoskeletal disorders* (6.6%), *musculoskeletal disorders–hypertension–asthma* (0.78%), *metabolic disorders* (2.58%) and *complex cardiometabolic disorders* (1.45%). Considering 329 775 000 (approximately 0.33 billion) older adult population as per World Population Prospects (2019),<sup>27</sup> we can infer that approximately 0.12 billion individuals belonged to one of the five multimorbidity classes (35.0%).

These resultant classes hinted towards a quantitative distinction, but three of these patterns were quite similar qualitatively. Considering the quantitative aspect, it is clear that *complex cardiometabolic disorders* followed by *metabolic disorders* were the most complicated disease patterns, with many NCDs coexisting simultaneously. Hypertension emerged as the most prevalent NCD with a high membership probability in five multimorbid classes. In addition, the disease patterns *hypertension*, *metabolic disorders* and *complex cardiometabolic disorders* reflect three stages of

**Table 4** Multinomial analysis of multimorbidity classes among older adults, Longitudinal Ageing Study in India, wave-1, 2017–2018

| Class                     | 2                    | 3  | 4  | 5                    | 6                                  |
|---------------------------|----------------------|--|--|----------------------|------------------------------------|
| Assigned labels           | Hypertension*        | Gastrointestinal disorders–hypertension–musculoskeletal disorders* | Musculoskeletal disorders–hypertension–asthma* | Metabolic disorders* | Complex cardiometabolic disorders* |
| <b>Correlates</b>         |                      |  |  |                      |                                    |
| Age (in years)            |                      |  |  |                      |                                    |
| 45–49 (Ref.)              | 1.00                 | 1.00   | 1.00   | 1.00                 | 1.00                               |
| 50–54                     | 1.45* (1.34 to 1.56) | 1.21* (1.08 to 1.37)   | 1.28 (0.83 to 1.97)                            | 1.70* (1.41 to 2.06) | 1.94* (1.38 to 6.72)               |
| 55–59                     | 1.78* (1.34 to 1.55) | 1.31* (1.16 to 1.48)   | 1.78* (1.18 to 2.67)                           | 2.22* (1.83 to 2.69) | 3.09* (2.22 to 4.29)               |
| 60–64                     | 2.02* (1.65 to 1.91) | 1.50* (1.33 to 1.69)   | 2.02* (1.35 to 3.01)                           | 2.79* (2.31 to 3.38) | 3.37* (2.43 to 4.68)               |
| 65–69                     | 2.52* (1.87 to 2.17) | 1.76* (1.55 to 1.99)   | 2.53* (1.69 to 3.78)                           | 2.85* (2.32 to 3.51) | 4.66* (3.34 to 6.46)               |
| 70–74                     | 2.72* (2.33 to 2.72) | 1.93* (1.67 to 2.23)   | 2.76* (1.78 to 4.28)                           | 3.29* (2.60 to 4.18) | 6.79* (4.80 to 9.61)               |
| 75–79                     | 2.76* (2.49 to 2.98) | 2.21* (1.86 to 2.62)   | 3.24* (2.00 to 5.24)                           | 3.26* (2.43 to 4.34) | 7.13* (4.84 to 11.04)              |
| 80+                       | 2.75* (2.48 to 3.07) | 2.53* (2.11 to 3.01)   | 4.14* (2.57 to 6.67)                           | 1.82* (1.24 to 2.68) | 7.04* (4.67 to 11.06)              |
| <b>Sex</b>                |                      |  |  |                      |                                    |
| Male (Ref.)               | 1.00                 | 1.00   | 1.00   | 1.00                 | 1.00                               |
| Female                    | 1.29* (1.21 to 1.36) | 1.44* (1.30 to 1.59)   | 0.96 (0.72 to 1.28)                            | 1.32* (1.12 to 1.56) | 1.50* (1.19 to 1.89)               |
| <b>Place of residence</b> |                      |  |  |                      |                                    |
| Rural (Ref.)              | 1.00                 | 1.00   | 1.00   | 1.00                 | 1.00                               |
| Urban                     | 1.30* (1.24 to 1.37) | 0.97 (0.90 to 1.05)  | 1.16 (0.93 to 1.46)                            | 1.61* (1.12 to 1.56) | 1.44* (1.22 to 1.69)               |
| <b>Religion</b>           |                      |  |  |                      |                                    |
| Hindu (Ref.)              | 1.00                 | 1.00   | 1.00   | 1.00                 | 1.00                               |
| Muslim                    | 1.29* (1.21 to 1.39) | 1.35* (1.21 to 1.50)   | 1.17 (0.85 to 1.58)                            | 2.93* (2.52 to 3.39) | 1.91* (1.54 to 2.34)               |
| Christian                 | 1.08* (1.24 to 1.37) | 0.90 (0.77 to 1.04)  | 0.56** (0.33 to 0.93)                          | 1.61* (1.32 to 1.96) | 0.98 (0.72 to 1.35)                |
| Others                    | 1.23* (1.21 to 1.38) | 1.09 (0.91 to 1.13)  | 0.56 (0.30 to 1.05)                            | 2.52* (2.07 to 3.04) | 1.08 (0.78 to 1.52)                |
| <b>Social group</b>       |                      |  |  |                      |                                    |
| Scheduled Castes (Ref.)   | 1.00                 | 1.00   | 1.00   | 1.00                 | 1.00                               |
| Scheduled Tribes          | 0.73 (0.66 to 0.78)  | 0.53* (0.46 to 0.62)   | 0.51* (0.33 to 0.78)                           | 0.65* (0.51 to 0.83) | 0.41* (0.27 to 0.59)               |
| Other Backward Classes    | 1.00 (0.94 to 1.06)  | 0.89* (0.81 to 0.98)   | 1.08 (0.81 to 1.42)                            | 1.14* (0.93 to 1.38) | 0.92* (0.72 to 0.92)               |
| Others                    | 1.05 (0.98 to 1.12)  | 1.01 (0.91 to 1.13)  | 0.99 (0.73 to 1.36)                            | 1.27 (1.04 to 1.53)  | 1.12 (0.87 to 1.43)                |
| <b>Level of education</b> |                      |  |  |                      |                                    |
| No education (Ref.)       | 1.00                 | 1.00   | 1.00   | 1.00                 | 1.00                               |
| Less than 5 years         | 1.23* (1.16 to 1.31) | 1.33* (1.21 to 1.45)   | 1.40* (1.08 to 1.81)                           | 2.61* (2.21 to 3.09) | 2.10* (1.69 to 2.59)               |
| 5–9 years                 | 1.21* (1.13 to 1.29) | 1.17* (1.05 to 1.30)   | 1.11 (0.81 to 1.81)                            | 2.62* (2.19 to 3.13) | 2.16* (1.72 to 2.73)               |
| 10 years or more          | 1.23* (1.15 to 1.32) | 0.99 (0.89 to 1.12)  | 0.72 (0.50 to 1.03)                            | 3.63* (3.06 to 4.31) | 1.94* (1.53 to 2.46)               |

Continued



**Table 4** Continued

| Class                                    | 2                     | 3  | 4  | 5                      | 6                                  |
|--|-----------------------|--|--|------------------------|------------------------------------|
| Assigned labels                          | Hypertension*         | Gastrointestinal disorders–hypertension–musculoskeletal disorders* | Musculoskeletal disorders–hypertension–asthma* | Metabolic disorders*   | Complex cardiometabolic disorders* |
| Wealth status                            |                       |  |  |                        |                                    |
| Poor (Ref.)                              | 1.00                  | 1.00   | 1.00   | 1.00                   | 1.00                               |
| Non-Poor                                 | 1.30* (1.24 to 1.36)  | 1.37* (1.27 to 1.47)   | 1.39* (1.13 to 1.71)                           | 1.69* (1.49 to 1.93)   | 2.13* (1.78 to 2.56)               |
| Current marital status                   |                       |  |  |                        |                                    |
| In union (Ref.)                          | 1.00                  | 1.00   | 1.00   | 1.00                   | 1.00                               |
| Not in union                             | 1.16 (1.10 to 1.23)   | 0.97 (0.89 to 1.06)  | 1.06 (0.83 to 1.35)                            | 1.05 (0.91 to 1.21)    | 0.96 (0.79 to 1.15)                |
| Current working status                   |                       |  |  |                        |                                    |
| Never worked (Ref.)                      | 1.00                  | 1.00   | 1.00   | 1.00                   | 1.00                               |
| Worked in past but not currently working | 1.13 (1.06 to 1.19)   | 1.04 (0.94 to 1.15)  | 1.16 (0.88 to 1.54)                            | 0.98 (0.84 to 1.14)    | 1.31 (1.07 to 1.60)                |
| Currently working                        | 0.73* (0.69 to 0.78)  | 0.72* (0.65 to 0.79)   | 0.59* (0.43 to 0.82)                           | 0.52* (0.43 to 0.61)   | 0.45 (0.34 to 0.57)                |
| Tobacco consumption                      |                       |  |  |                        |                                    |
| Never used tobacco (Ref.)                | 1.00                  | 1.00   | 1.00   | 1.00                   | 1.00                               |
| Quit tobacco                             | 1.12* (1.02 to 1.23)  | 1.75* (1.52 to 2.03)   | 3.01* (2.19 to 4.13)                           | 1.43* (1.15 to 1.78)   | 3.01* (1.95 to 3.27)               |
| Currently consuming                      | 0.89 (0.84 to 1.93)   | 1.40 (1.28 to 1.52)  | 1.17 (0.91 to 1.49)                            | 0.56 (0.46 to 0.66)    | 1.01 (0.82 to 1.24)                |
| Ever consumed alcohol                    |                       |  |  |                        |                                    |
| No (Ref.)                                | 1.00                  | 1.00   | 1.00   | 1.00                   | 1.00                               |
| Yes                                      | 1.01 (0.94 to 1.08)   | 1.03 (0.92 to 1.14)  | 1.20 (0.91 to 1.58)                            | 1.52* (1.27 to 1.81)   | 1.01 (0.78 to 1.29)                |
| Physical activity                        |                       |  |  |                        |                                    |
| Physically active (Ref.)                 | 1.00                  | 1.00   | 1.00   | 1.00                   | 1.00                               |
| Physically inactive                      | 1.13** (1.08 to 1.18) | 0.95 (0.89 to 1.03)  | 1.17 (0.94 to 1.45)                            | 1.08 (0.95 to 1.21)    | 1.46* (1.25 to 1.72)               |
| Body mass index                          |                       |  |  |                        |                                    |
| Underweight (Ref.)                       | 1.00                  | 1.00   | 1.00   | 1.00                   | 1.00                               |
| Normal                                   | 1.64* (1.54 to 1.75)  | 1.30* (1.17 to 1.43)   | 0.82* (0.63 to 1.05)                           | 2.59* (1.90 to 3.52)   | 2.10* (1.53 to 2.90)               |
| Overweight                               | 2.78* (2.53 to 4.28)  | 1.90* (1.69 to 2.14)   | 0.96 (0.39 to 3.21)                            | 6.26* (4.58 to 8.56)   | 5.28* (3.78 to 7.38)               |
| Obese                                    | 3.88* (3.53 to 4.28)  | 2.68* (2.30 to 3.13)   | 2.11* (1.39 to 3.21)                           | 10.50* (7.57 to 11.45) | 9.52* (6.62 to 13.68)              |
| Waist-hip ratio                          |                       |  |  |                        |                                    |
| Low risk (Ref.)                          | 1.00                  | 1.00   | 1.00   | 1.00                   | 1.00                               |
| High risk                                | 1.24* (1.17 to 1.31)  | 1.14** (1.04 to 1.26)  | 1.02 (0.79 to 1.31)                            | 2.38* (1.87 to 3.03)   | 1.51* (1.15 to 1.97)               |

\*p<0.001, \*\*p<0.05.

\*All estimates are computed considering relatively healthy as the base category.

cardiometabolic morbidity. It can be hypothesised that if individuals are affected with hypertension, they are more likely to accumulate other cardiometabolic morbidities, like diabetes, high cholesterol and heart diseases. From a programmatic standpoint, the study findings underscore an urgent need for effective prevention and management strategies for individuals affected with hypertension.

As the present study is the first attempt to explore prominent disease patterns among older adults in India, comparison with other studies is taxing, primarily due to the ambiguities in the operational definition of multimorbidity, differences in the age groups involved and other methodological differences. However, recent studies proposed cardiovascular and metabolic disorders as frequently occurring diseases in the country.<sup>18 28</sup> It is worth mentioning that the studies above included fewer chronic morbidities. In addition, identified disease clusters are in concordance with the systematic review based on 14 studies set in different geographical settings.<sup>29</sup>

Age originated as one of the primary predictors of all identified multimorbidity patterns among older adults. Ageing is a universal process characterised by unescapable alterations in various biological and neurological processes, resulting in progressive functional decline at cellular levels.<sup>30</sup> These alterations might also reflect an added predisposition towards chronic diseases. However, ageing is not synonymous with ill-health or multimorbidity. Consciously employing healthful dietary patterns, incorporating an active lifestyle including physical activities and managing other behavioural aspects and environmental exposures can alter the situation.<sup>31</sup>

Similarly, belonging to the affluent class was significantly associated with all multimorbidity patterns, whereas being educated increased the likelihood of belonging to *hypertension*, *metabolic disorders* and *complex cardiometabolic disorders*. At the same time, residing in urban areas increased the likelihood of belonging to *hypertension*, *metabolic disorders* and *complex cardiometabolic disorders*. This hints toward the preponderance of multimorbidity among individuals with better availability and accessibility to early screening, diagnoses and treatment services.<sup>18 32</sup> Through 'Ayushman Bharat', the government of India has established health and wellness centres across the country for timely detection, prevention and effective management of diseases.<sup>33</sup> Despite this, there are issues like improper referral networks, inefficient supply chain management and staff.<sup>18</sup> In addition, healthcare providers' poor behaviours and low-personal competence have been highlighted by recent studies.<sup>34</sup> This majorly affects the quality of government-managed public health centres. Thus, the individuals who do not have economic or non-economic resources like lack of transportation facilities, having a disability, living in rural areas and work timings face challenges in accessing affordable quality healthcare. However, this might not thoroughly explain the higher disease burden among affluent, educated and urban residents. Additional reasons could be better access to health information and education in this group,

enabling them to identify early symptoms of NCDs and better compliance with treatment regimens.<sup>35</sup>

Women were more prone to being in disease class *hypertension*, *gastrointestinal disorders–hypertension–musculoskeletal disorders*, *metabolic disorders* and *complex cardiometabolic disorders*. Our study findings are in concordance with the existing literature, which suggests that India is experiencing feminisation of multimorbidity,<sup>7 18 36</sup> primarily due to increased life expectation and prolonged exposure to social isolation.<sup>8 18</sup>

Belonging to the ST social group decreased the likelihood of belonging to class *gastrointestinal disorders–hypertension–musculoskeletal disorders*, *musculoskeletal disorders–hypertension–asthma*, *metabolic disorders* and *complex cardiometabolic disorders*. Similar findings have also been highlighted in recent studies.<sup>8 37</sup> In a way, this could indicate better health among the ST population. However, as the data used in the present study is self-reported diagnoses, the estimates generated are primarily based on individual's perception and awareness. They thus could indicate an underdiagnosed NCD burden among this population.<sup>38</sup> Tribal population is considered the most disadvantaged population subgroup in India.<sup>39</sup> They reside in remote locations, have inadequate awareness about health risks and have limited access to good quality healthcare.<sup>40 41</sup> Another issue is the low acceptability of healthcare among this social group which is highlighted as a significant cause of low diagnoses and treatment rates. These findings, therefore, hint toward unequal access to healthcare, which in turn suggests an urgent need for adequate awareness, accessibility, affordability, availability and acceptability of the healthcare services by this population.<sup>39 42 43</sup>

The lifestyle predictor: quitting tobacco increased the likelihood of belonging to the five multimorbid classes. As tobacco cessation is the key to NCD reduction, the medical practitioner recommended that multimorbid individuals quit tobacco.<sup>44</sup> Obesity increased the likelihood of belonging to class any of the five multimorbid classes. Furthermore, being physically inactive increased the likelihood of belonging to class: *hypertension and complex cardiometabolic disorders*, whereas high-risk WHR increased the likelihood of belonging to *hypertension*, *gastrointestinal disorders–hypertension–musculoskeletal disorders*, *metabolic disorders* and *complex cardiometabolic disorders*. These findings indicate the importance of lifestyle and anthropometric dynamics in intensifying exposure to associative chronic morbidities in any community, as already highlighted in existing literature for various subsections of the population.<sup>9 18</sup>

The study's major strength is that it provides empirical evidence on recurrent disease patterns using a large nationally-representative sample of older adults. However, the NCDs included in the study are based on self-reporting, which can lead to misclassification bias. No causality was tested in the study as findings are based on a single round of a longitudinal survey. In addition, the methodological limitations of LCA, a data driven technique cannot

be denied. Class assignment is based on the probability-based classification method, which is a potential source of bias. In addition, as the designated class labels are based on the author's discretion thus, subjectivity and naming fallacy cannot be ruled out. Furthermore, considering the geographical and contextual variations in disease burden and profiling, these results should be explicitly apprehended for the older adults in India.

## CONCLUSIONS

These findings can recognise associated diseases among patients affected with one or more morbidities during hospital visits. The classes 'hypertension', 'metabolic disorders' and 'complex cardiometabolic disorders' reflect three stages of cardiometabolic morbidity, with hypertension as the first and 'complex cardiometabolic disorders' as the last stage of disease progression. This underscores the need for an effective prevention strategy. These findings can assist physicians and policymakers in devising practical strategies for control, prevention and management for individuals belonging to a specific disease class not to accumulate additional diseases. Thus, advocating policies to reorganise the existing healthcare services in a way to accommodate the rising requirements of the older adult population (45 years and older), which is estimated to reach 0.66 billion by 2050.<sup>27</sup> Alternatively, targeted interventions in the form of equitable prevention strategies, are essential to reduce the burden on the high-risk hypertensive older adult population in the country.

**Acknowledgements** The authors are grateful to the Longitudinal Ageing Study in India (LASI) for assembling and publishing accurate, nationally representative data on a range of health, biomarkers and healthcare utilisation indicators for the population aged 45 years and older. The authors are also grateful to LASI's project partners, the International Institute for Population Sciences (IIPS), Harvard T. H. Chan School of Public Health and the University of Southern California (USC). The findings of this article were presented at the British Society for Population Studies (BSPS) Conference 2021, London School of Economics, UK. The parts of these article were also presented at the IUSSP's International Population Conference (IPC), 2021.

**Contributors** PP conceived the idea for the study, conducted the statistical analysis and is responsible for the initial draft of the report. PP, SKS and SP contributed to subsequent and final draft. SKS and SP were responsible for supervision. PP acts as guarantor for the final manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not applicable.

**Ethics approval** The study used de-identified data from a secondary data source, Longitudinal Ageing Study in India, 2017–2018. The survey followed all necessary guidelines and received ethical approval from the Indian Council of Medical Research (ICMR), and Institute Review Board held at the International Institute for Population Sciences, Mumbai, India. Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. The data has been archived in the public repository of the Longitudinal Ageing Study in India, 2017–18. Access to the data requires registration and is granted only for legitimate

research purposes. A guide on applying for the data set access is available at <https://www.iipsindia.ac.in/content/LASI-data>. The data can be accessed by filling out the request form available at [https://iipsindia.ac.in/sites/default/files/LASI\\_DataRequestForm\\_0.pdf](https://iipsindia.ac.in/sites/default/files/LASI_DataRequestForm_0.pdf) and sending it to the Data Center ([datacenter@iipsindia.ac.in](mailto:datacenter@iipsindia.ac.in)) held at the International Institute for Population Sciences, Mumbai, India.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iD

Parul Puri <http://orcid.org/0000-0001-6272-837X>

## REFERENCES

- 1 The World Bank. Life expectancy at birth, 2019. Available: <https://data.worldbank.org/indicator/SP.DYN.LE00.IN?locations=IN>
- 2 Puri P, Singh SK, Pati S. Temporal dynamics, patterns and correlates of single and multimorbidity in India, 1994–2018. *J Multimorbidity Comorbidity* 2021;11:263355652110627.
- 3 Pati S, Swain S, Hussain MA, et al. Prevalence, correlates, and outcomes of multimorbidity among patients attending primary care in Odisha, India. *Ann Fam Med* 2015;13:446–50.
- 4 Shekhar C, Shil A. Linkages between occupation and elevated blood pressure among men in India: a cross-sectional study. *Int J Public Health* 2020;65:835–46.
- 5 Singh S, Puri P, Subramanian SV. Identifying spatial variation in the burden of diabetes among women across 640 districts in India: a cross-sectional study. *J Diabetes Metab Disord* 2020;19:523–33.
- 6 Puri P, Shil A, Shetty A, et al. Contribution of modifiable risk factors on the burden of diabetes among women in reproductive age-group in India: a population based cross-sectional study. *J Public Health Policy* 2022;43:89–108.
- 7 Arokiasamy P, Uttamacharya JK. Multi-morbidity, functional limitations, and self-rated health among older adults in India: cross-sectional analysis of LASI pilot survey, 2010. *SAGE Open* 2015;5.
- 8 Pati S, Swain S, Metsemakers J, et al. Pattern and severity of multimorbidity among patients attending primary care settings in Odisha, India. *PLoS One* 2017;12:e0183966–19.
- 9 Puri P, Kothavale A, Singh SK, et al. Burden and determinants of multimorbidity among women in reproductive age group: a cross-sectional study based in India. *Wellcome Open Res* 2021;5:275–21.
- 10 Srivastava S, Joseph K J V, Drishti D, et al. Interaction of physical activity on the association of obesity-related measures with multimorbidity among older adults: a population-based cross-sectional study in India. *BMJ Open* 2021;11:e050245–10.
- 11 Jackson CA, Jones M, Tooth L, et al. Multimorbidity patterns are differentially associated with functional ability and decline in a longitudinal cohort of older women. *Age Ageing* 2015;44:810–6.
- 12 Larsen FB, Pedersen MH, Friis K, et al. A latent class analysis of multimorbidity and the relationship to socio-demographic factors and health-related quality of life. A national population-based study of 162,283 Danish adults. *PLoS One* 2017;12:e0169426–17.
- 13 Yap KH, Warren N, Allotey P, et al. Chronic disease profiles of subjective memory complaints: a latent class analysis of older people in a rural Malaysian community. *Ageing Ment Health* 2020;24:709–16.
- 14 Park B, Lee HA, Park H. Use of latent class analysis to identify multimorbidity patterns and associated factors in Korean adults aged 50 years and older. *PLoS One* 2019;14:e0216259–13.
- 15 Craig LS, Hotchkiss DR, Theall KP, et al. Prevalence and patterns of multimorbidity in the Jamaican population: a comparative analysis of latent variable models. *PLoS One* 2020;15:e0236034.
- 16 Olaya B, Moneta MV, Caballero FF, et al. Latent class analysis of multimorbidity patterns and associated outcomes in Spanish older adults: a prospective cohort study. *BMC Geriatr* 2017;17:1–10.

- 17 International Institute for Population Sciences (IIPS). Longitudinal ageing study in India (LASI); 2020: 1–632. [https://www.ipsindia.ac.in/sites/default/files/LASI\\_India\\_Report\\_2020\\_compressed.pdf](https://www.ipsindia.ac.in/sites/default/files/LASI_India_Report_2020_compressed.pdf) [Accessed 07-07-2021].
- 18 Puri P, Singh SK. Patterns and correlates of non-communicable disease multimorbidity among older adults in India : Evidence from Longitudinal Ageing Study in India (LASI), 2017-18. *J Public Health Policy*. Published online 2022.
- 19 Weller BE, Bowen NK, Faubert SJ. Latent class analysis: a guide to best practice. *J Black Psychol* 2020;46:287–311.
- 20 Craig LS, Cunningham-Myrie CA, Hotchkiss DR, et al. Social determinants of multimorbidity in Jamaica: application of latent class analysis in a cross-sectional study. *BMC Public Health* 2021;21:1–15.
- 21 Linzer DA, Lewis JB. polCA : An R Package for Polytomous Variable. *J Stat Softw* 2011;42:1–29.
- 22 Zhang Z, Abarda A, Contractor AA, et al. Exploring heterogeneity in clinical trials with latent class analysis. *Ann Transl Med* 2018;6:119.
- 23 Lanza S. Latent Class Analysis. In: *Statistical horizons*, 2017: 1–16. <https://statisticalhorizons.com/wp-content/uploads/LCA-Sample-Materials.pdf>
- 24 Northstone K, Ness AR, Emmett PM, et al. Adjusting for energy intake in dietary pattern investigations using principal components analysis. *Eur J Clin Nutr* 2008;62:931–8.
- 25 Yong AG, Pearce S. A Beginner's Guide to Factor Analysis: Focusing on Exploratory Factor Analysis. *Tutor Quant Methods Psychol* 2013;9:79–94.
- 26 Arifin WN, Yusoff MSB. Confirmatory factor analysis of the Universiti Sains Malaysia emotional quotient inventory among medical students in Malaysia. *SAGE Open* 2016;6:215824401665024.
- 27 United Nations. World population prospects 2019, 2019. Available: <https://population.un.org/wpp/Download/Standard/Population/>
- 28 Pati S, Mahapatra P, Kanungo S, et al. Managing multimorbidity (multiple chronic diseases) amid COVID-19 pandemic: a community based study from Odisha, India. *Front Public Health* 2020;8:1–9.
- 29 Prados-Torres A, Calderón-Larrañaga A, Hanco-Saavedra J, et al. Multimorbidity patterns: a systematic review. *J Clin Epidemiol* 2014;67:254–66.
- 30 Adav SS, Wang Y. Metabolomics signatures of aging: recent advances. *Aging Dis* 2021;12:646–16.
- 31 Jin K. Modern biological theories of aging. *Aging Dis* 2010;1:72–4.
- 32 Pati S, Sinha R, Mahapatra P. Non-communicable disease risk reduction teaching in India: a curricular landscape. *Front Public Health* 2019;7:1–7.
- 33 Ministry of Health and Family Welfare. Ayushman Bharat - Health and Wellness Centre; 2020: 1–24. [https://nhm.gov.in/New\\_Updates\\_2018/publication/AB-HWC-Brochure-March-2021-English.pdf](https://nhm.gov.in/New_Updates_2018/publication/AB-HWC-Brochure-March-2021-English.pdf) [Accessed 07-07-2021].
- 34 Kasthuri A. Challenges to healthcare in India - The five A's. *Indian J Community Med* 2018;43:141–3.
- 35 Puri P, Kothavale A, Singh SK, et al. Burden and determinants of multimorbidity among women in reproductive age group: a cross-sectional study based in India. *Wellcome Open Res* 2020;5:275.
- 36 Schäfer I, Kaduszkiewicz H, Nguyen TS, et al. Multimorbidity patterns and 5-year overall mortality: results from a claims data-based observational study. *J Comorb* 2018;8:2235042X1881658.
- 37 Mini GK, Thankappan KR. Pattern, correlates and implications of non-communicable disease multimorbidity among older adults in selected Indian states: a cross-sectional study. *BMJ Open* 2017;7:e013529.
- 38 United Nations. Situation of Indigenous peoples and rights to health. 2016.
- 39 Puri P, Pati S. Exploring the linkages between non-communicable disease multimorbidity. *Health Care Utilization and Expenditure Among Aboriginal Older Adult Population in India* 2022;67.
- 40 Haddad S, Mohindra KS, Siekmans K, et al. "Health divide" between indigenous and non-indigenous populations in Kerala, India: Population based study. *BMC Public Health* 2012;12:1.
- 41 Thresia CU, Srinivas PN, Mohindra KS. The health of Indigenous populations in South Asia: a critical review in a critical time. *Int J Heal Serv* 2020.
- 42 Davy C, Harfield S, McArthur A, et al. Access to primary health care services for Indigenous peoples: a framework synthesis. *Int J Equity Health* 2016;15:1–9.
- 43 Government of India. Report of the expert Committee on tribal health: tribal health in India: bridging the AGP and a roadmap for the future. India; 2018: 1–56. [http://nhm.gov.in/nhm\\_components/tribal\\_report/Executive\\_Summary.pdf](http://nhm.gov.in/nhm_components/tribal_report/Executive_Summary.pdf) [Accessed 07-07-2021].
- 44 Wilson MS, Metink-Kane MM. Effective tobacco control is key to apid progress in reduction of non-communicable diseases. *Lancet* 2012;23:1–7.



**Supplementary File 1.** STROBE cross-sectional checklist for the study.

|                            | Reporting Item  | Page No. |
|----------------------------|---|----------|
| <b>Title and abstract</b>  |   |          |
| Title                      | <a href="#">#1a</a> Indicate the study's design with a commonly used term in the title or the abstract  | 1        |
| Abstract                   | <a href="#">#1b</a> Provide in the abstract an informative and balanced summary of what was done and what was found   | 2        |
| <b>Introduction</b>        |   |          |
| Background / rationale     | <a href="#">#2</a> Explain the scientific background and rationale for the investigation being reported   | 4-5      |
| Objectives                 | <a href="#">#3</a> State specific objectives, including any prespecified hypotheses   | 4-5      |
| <b>Methods</b>             |   |          |
| Study design               | <a href="#">#4</a> Present key elements of study design early in the paper  | 5        |
| Setting                    | <a href="#">#5</a> Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection  | 5-6      |
| Eligibility criteria       | <a href="#">#6a</a> Give the eligibility criteria, and the sources and methods of selection of participants.  | NA       |
|                            | <a href="#">#7</a> Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable   | 5-7      |
| Data sources / measurement | <a href="#">#8</a> For each variable of interest give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group. Give information separately for for exposed and unexposed groups if applicable.             | 5-7      |
| Bias                       | <a href="#">#9</a> Describe any efforts to address potential sources of bias  | NA       |
| Study size                 | <a href="#">#10</a> Explain how the study size was arrived at   | 5        |
| Quantitative variables     | <a href="#">#11</a> Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen, and why   | 5-7      |
| Statistical methods        | <a href="#">#12a</a> Describe all statistical methods, including those used to control for confounding  | 7-8      |
| Statistical methods        | <a href="#">#12b</a> Describe any methods used to examine subgroups and interactions  | 7-8      |
| Statistical methods        | <a href="#">#12c</a> Explain how missing data were addressed  | NA       |
| Statistical methods        | <a href="#">#12d</a> If applicable, describe analytical methods taking account of sampling strategy   | NA       |
| Statistical methods        | <a href="#">#12e</a> Describe any sensitivity analyses  | NA       |
| <b>Results</b>             |   |          |
| Participants               | <a href="#">#13a</a> Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed. Give information separately for for exposed and unexposed groups if applicable. | 8        |
| Participants               | <a href="#">#13b</a> Give reasons for non-participation at each stage   | NA       |
| Participants               | <a href="#">#13c</a> Consider use of a flow diagram   | NA       |
| Descriptive data           | <a href="#">#14a</a> Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders. Give information separately for exposed and unexposed groups if applicable.  | 8        |
| Descriptive data           | <a href="#">#14b</a> Indicate number of participants with missing data for each variable of interest  | NA       |
| Outcome data               | <a href="#">#15</a> Report numbers of outcome events or summary measures. Give information separately for exposed and unexposed groups if applicable.   | 8        |
| Main results               | <a href="#">#16a</a> Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included   | 8-11     |
| Main results               | <a href="#">#16b</a> Report category boundaries when continuous variables were categorized  | 7        |

|                          |                      |  |       |
|--------------------------|----------------------|--|-------|
| Main results             | <a href="#">#16c</a> | If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period   | NA    |
| Other analyses           | <a href="#">#17</a>  | Report other analyses done—e.g., analyses of subgroups and interactions, and sensitivity analyses  | NA    |
| <b>Discussion</b>        |                      |  |       |
| Key results              | <a href="#">#18</a>  | Summarize key results with reference to study objectives   | 12    |
| Limitations              | <a href="#">#19</a>  | Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias.      | 3, 15 |
| Interpretation           | <a href="#">#20</a>  | Give a cautious overall interpretation considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence. | 12-15 |
| Generalizability         | <a href="#">#21</a>  | Discuss the generalizability (external validity) of the study results  | 12-15 |
| <b>Other Information</b> |                      |  |       |
| Funding                  | <a href="#">#22</a>  | Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based    | 16    |

**Supplementary File 2.** Socio-demographic, lifestyle and anthropometric characteristic of the six identified latent classes, Longitudinal Ageing Study in India (LASI), wave-1, 2017-18.

| Predictors                | Latent Class       |              |   |   |                     |                                   |
|---------------------------|--------------------|--------------|---|---|---------------------|-----------------------------------|
|                           | 1                  | 2            | 3   | 4   | 5                   | 6                                 |
|                           | Relatively Healthy | Hypertension | Gastrointestinal Disorders-Hypertension-Musculoskeletal Disorders | Musculoskeletal disorders-Hypertension-Asthma | Metabolic disorders | Complex cardiometabolic disorders |
| <b>Age (in years)</b>     |                    |              |   |   |                     |                                   |
| 45-49                     | 9029 (75.76)       | 1893 (15.88) | 654 (5.49)  | 53 (0.44)                                     | 211 (1.77)          | 77 (0.64)                         |
| 50-54                     | 6899 (69.95)       | 1995 (20.23) | 574 (5.82)  | 50 (0.51)                                     | 251 (2.54)          | 94 (0.95)                         |
| 55-59                     | 5972 (65.94)       | 2093 (23.11) | 558 (6.16)  | 60 (0.66)                                     | 255 (2.82)          | 119 (1.31)                        |
| 60-64                     | 5763 (62.68)       | 2305 (25.07) | 623 (6.78)  | 72 (0.78)                                     | 298 (3.24)          | 134 (1.45)                        |
| 65-69                     | 4619 (57.88)       | 2324 (9.12)  | 566 (7.09)  | 79 (0.99)                                     | 239 (2.99)          | 154 (2.49)                        |
| 70-74                     | 2854 (55.68)       | 1551 (30.26) | 388 (7.57)  | 54 (1.05)                                     | 151 (2.94)          | 128 (2.49)                        |
| 75-79                     | 1633 (54.62)       | 910 (30.43)  | 247 (8.26)  | 42 (1.40)                                     | 82 (2.74)           | 76 (2.54)                         |
| 80+                       | 1573 (55.27)       | 844 (29.66)  | 267 (9.38)  | 49 (1.72)                                     | 40 (1.41)           | 73 (2.57)                         |
| <b>Sex</b>                |                    |              |   |   |                     |                                   |
| Male                      | 18897 (69.03)      | 5627 (20.56) | 1627 (5.94)   | 233 (0.85)                                    | 637 (2.32)          | 353 (1.29)                        |
| Female                    | 19445 (61.53)      | 8288 (26.23) | 2250 (7.12)   | 226 (0.72)                                    | 890 (2.82)          | 502 (1.59)                        |
| <b>Place of Residence</b> |                    |              |   |   |                     |                                   |
| Rural                     | 26953 (69.62)      | 7858 (20.29) | 2592 (6.70)   | 304 (0.79)                                    | 604 (1.56)          | 404 (1.04)                        |
| Urban                     | 11389 (56.21)      | 6057 (29.90) | 1285 (6.34)   | 155 (0.77)                                    | 923 (4.56)          | 451 (1.59)                        |
| <b>Religion</b>           |                    |              |   |   |                     |                                   |
| Hindu                     | 28644 (66.25)      | 9890 (22.87) | 2870 (6.63)   | 363 (0.84)                                    | 880 (2.04)          | 591 (1.37)                        |
| Muslim                    | 3976 (57.07)       | 1907 (27.37) | 546 (7.84)  | 59 (0.85)                                     | 327 (4.69)          | 153 (2.20)                        |
| Christian                 | 4145 (69.27)       | 1316 (21.99) | 269 (4.49)  | 25 (0.42)                                     | 162 (2.71)          | 67 (1.12)                         |
| Others                    | 1577 (56.62)       | 802 (28.80)  | 192 (6.89)  | 12 (0.43)                                     | 158 (5.67)          | 44 (1.58)                         |
| <b>Social group</b>       |                    |              |   |   |                     |                                   |
| Scheduled Castes          | 6744 (67.72)       | 2175 (21.84) | 711 (7.13)  | 78 (0.78)                                     | 145 (1.46)          | 106 (1.06)                        |
| Scheduled Tribes          | 7832 (75.68)       | 1821 (17.60) | 418 (4.03)  | 46 (0.44)                                     | 163 (1.58)          | 68 (0.65)                         |
| Other Backward Classes    | 14392 (64.70)      | 5329 (23.96) | 1477 (6.64)   | 204 (0.92)                                    | 540 (2.43)          | 301 (1.35)                        |
| Others                    | 9374 (57.07)       | 4590 (27.95) | 1271 (7.74)   | 131 (0.80)                                    | 679 (4.13)          | 380 (2.31)                        |
| <b>Level of Education</b> |                    |              |   |   |                     |                                   |
| No education              | 19080 (68.72)      | 6057 (21.81) | 1825 (6.57)   | 222 (0.80)                                    | 309 (1.11)          | 273 (0.98)                        |
| Less than 5 years         | 6882 (62.80)       | 2645 (24.14) | 815 (7.44)  | 108 (0.99)                                    | 317 (2.89)          | 192 (1.75)                        |
| 5-9 years                 | 5956 (63.49)       | 2254 (24.03) | 616 (6.57)  | 72 (0.77)                                     | 311 (3.32)          | 171 (1.82)                        |
| 10 years or more          | 6424 (59.09)       | 2959 (27.22) | 621 (5.71)  | 57 (0.52)                                     | 590 (5.43)          | 219 (2.01)                        |
| <b>Wealth Tertile</b>     |                    |              |   |   |                     |                                   |
| Poor                      | 16734 (71.09)      | 4746 (20.16) | 1328 (5.64)   | 170 (0.72)                                    | 357 (1.52)          | 202 (0.85)                        |
| Non-poor                  | 21608 (60.97)      | 9169 (25.87) | 2549 (7.19)   | 289 (0.82)                                    | 1170 (3.30)         | 653 (1.84)                        |

|  |               |               |              |            |             |            |
|--|---------------|---------------|--------------|------------|-------------|------------|
|  |               |               |              |            |             |            |
| <b>Current Marital Status</b>            |               |               |              |            |             |            |
| In union                                 | 29735 (66.93) | 9736 (21.91)  | 2855 (6.43)  | 321 (0.72) | 1175 (2.64) | 605 (1.36) |
| Not in union                             | 8607 (59.16)  | 4179 (28.73)  | 1022 (7.03)  | 138 (0.95) | 352 (2.42)  | 250 (1.72) |
| <b>Current Working Status</b>            |               |               |              |            |             |            |
| Never worked                             | 9141 (56.72)  | 4674 (29.00)  | 1233 (7.65)  | 123 (0.76) | 625 (3.88)  | 319 (1.98) |
| Worked in past but not currently working | 8727 (56.27)  | 4551 (29.34)  | 1185 (7.64)  | 188 (1.21) | 482 (3.11)  | 377 (2.43) |
| Currently Working                        | 20474 (74.86) | 4690 (17.15)  | 1459 (5.33)  | 148 (0.54) | 420 (1.54)  | 159 (0.58) |
| <b>Tobacco Consumption</b>               |               |               |              |            |             |            |
| Never used Tobacco                       | 174 (1.23)    | 9698 (68.61)  | 2255 (15.95) | 242 (1.71) | 1190 (8.42) | 575 (4.07) |
| Consumed in past but not currently       | 1886 (57.54)  | 817 (24.92)   | 280 (8.54)   | 63 (1.92)  | 129 (3.94)  | 103 (3.14) |
| Currently Consuming                      | 13282 (71.56) | 3400 (18.32)  | 1342 (7.23)  | 154 (0.83) | 208 (1.12)  | 177 (0.95) |
| <b>Ever consumed alcohol</b>             |               |               |              |            |             |            |
| No                                       | 30750 (63.59) | 11970         | 3240 (6.70)  | 361 (0.75) | 1296 (2.68) | 737 (1.52) |
| Yes                                      | 7592 (71.48)  | 1945          | 637 (5.99)   | 98 (0.92)  | 231 (2.17)  | 118 (1.11) |
| <b>Physical Activity</b>                 |               |               |              |            |             |            |
| Physically Active                        | 27730 (67.15) | 9071          | 2722 (6.59)  | 285 (0.69) | 1017 (2.46) | 470 (1.13) |
| Physically Inactive                      | 10612 (60.02) | 4844          | 1155 (6.53)  | 174 (0.98) | 510 (2.88)  | 385 (2.18) |
| <b>Body Mass Index</b>                   |               |               |              |            |             |            |
| Underweight                              | 8425 (77.85)  | 1533          | 634 (5.85)   | 122 (1.13) | 46 (0.42)   | 62 (0.57)  |
| Normal                                   | 20985 (68.74) | 6576          | 1909 (6.25)  | 215 (0.70) | 526 (1.72)  | 318 (1.04) |
| Overweight                               | 6775 (53.55)  | 3977          | 925 (7.31)   | 78 (0.62)  | 610 (4.82)  | 286 (2.26) |
| Obese                                    | 1799 (41.86)  | 1620          | 362 (8.42)   | 40 (0.93)  | 325 (7.56)  | 152 (3.54) |
| Don't Know                               | 358 (53.04)   | 209           | 47 (6.96)    | 4 (0.59)   | 20 (2.96)   | 37 (5.48)  |
| <b>Waist-Hip Ratio</b>                   |               |               |              |            |             |            |
| Low Risk                                 | 8959 (76.16)  | 1891          | 661 (5.62)   | 99 (0.84)  | 72 (0.61)   | 81 (0.69)  |
| High Risk                                | 29383 (62.24) | 12024         | 3216 (6.81)  | 360 (0.76) | 1455 (3.08) | 774 (1.63) |
| <b>Total</b>                             | 38342 (65.01) | 13915 (23.59) | 3877 (6.57)  | 459 (0.78) | 1527 (2.58) | 855 (1.45) |