

BMJ Open Prevalence and factors associated with self-reported injuries in Nepal: a secondary analysis of the nationally representative cross-sectional STEPS Survey, 2019

Meghnath Dhimal ¹, Anil Poudyal ¹, Bihungum Bista,¹ Sitasnu Dahal,¹ Puspaj Pant,² Pradip Gyanwali¹

To cite: Dhimal M, Poudyal A, Bista B, *et al*. Prevalence and factors associated with self-reported injuries in Nepal: a secondary analysis of the nationally representative cross-sectional STEPS Survey, 2019. *BMJ Open* 2022;**12**:e060561. doi:10.1136/bmjopen-2021-060561

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-060561>).

MD and AP contributed equally.

MD and AP are joint first authors.

Received 27 December 2021
Accepted 26 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Meghnath Dhimal;
meghdhimal@gmail.com and
Anil Poudyal;
poudyalanil123@gmail.com

ABSTRACT

Objective This study aims to determine the prevalence and factors associated with injuries in the adult population of Nepal.

Design and participants Secondary analysis of the data from the cross-sectional WHO STEPwise Approach to NCD Risk Factor Surveillance (STEPS) Survey Nepal, 2019. A multistage cluster sample of 5593 adults aged 15–69 years who have been the usual residents of the household for at least 6 months. A binary logistic regression model was employed to identify the determinants of injuries.

Setting Data were derived from the STEPS Survey Nepal, 2019.

Primary and secondary outcome measures The primary outcome was injured person defined as one who had road traffic injuries (RTIs), had other serious unintentional/accidental injury, or had been seriously injured in a violent incident within the past 12 months. The secondary outcome measure was factors associated with injuries.

Results Over 11% of the 4996 study participants reported any injuries during the past 12 months. About 3.75% of the participants experienced a RTI, 4.71% had experienced unintentional injuries other than RTI, while 5.33% had been injured in violent incidents. Individuals belonging to the middle wealth quintile (crude OR (COR)=2.95, 95% CI 1.27 to 6.84) were associated with increased odds of RTIs. By occupation, homemaker (COR=0.45, 95% CI 0.24 to 0.84) was protective against these injuries. Likewise, currently married individuals (COR=3.74, 95% CI 1.37 to 10.17), ever married individuals (COR=3.49, 95% CI 1.08 to 11.25) and individuals not in employment (COR=2.13, 95% CI 1.16 to 3.91) were associated with an increased likelihood of sustaining an intentional injury. Injuries were higher among rural participants.

Conclusions This study provides the baseline population-based estimates of the prevalence of injuries in Nepal. It describes the mechanisms and risk factors of these injuries. It is hoped that this evidence will serve as a stimulus for future studies to elucidate comprehensive national information about injuries.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ To our knowledge, this is the first study to assess the magnitude and risk factors of injuries based on the most recent nationally representative sample of the Nepalese population.
- ⇒ We used multistage stratified cluster sampling and probability weights to avoid potential biases.
- ⇒ Self-reported information collected from past events is subject to recall bias.
- ⇒ Due to the cross-sectional nature of the study, causal relationships could not be identified.

BACKGROUND

Injuries are one of the leading causes of premature death and disabilities globally.¹ According to world health organization (WHO) estimates, around 4.4 million deaths occur globally from injuries, which is nearly 8% of total deaths from all causes.² Road traffic injuries (RTIs) are responsible for 1.3 million deaths, unintentional injuries cause 1.8 million deaths, and self-harm and violent injuries are responsible for 1.3 million deaths.³ Additionally, injuries contribute to 69.3 million years lived with disability (YLDs). The top three leading causes of death worldwide due to injuries including road traffic crashes, suicide and homicide are all predicted to increase in rank compared with other causes of death, placing them among the top 20 leading causes of death in the world by 2030.⁴ Developing countries carry the largest burden of injuries where more than 90% of deaths and 94% of disability-adjusted life years (DALYs) resulting from injuries occur in low-income and middle income countries (LMICs).⁵ The burden of injury is highest in the younger age group. This age group is believed to be the most productive, and injury will, in



turn, have an impact on the growth and financial development of a nation.⁶

Nepal is an LMIC and has a substantial burden from injuries in terms of a high number of hospital visits, hospitalisation and disabilities. Injuries are a major cause of death. The Integrated Health Management Information System record shows that injuries were the reason for 1.4 million outpatient morbidities in Nepal in the year 2020/2021.⁷ In Nepal, economic development, urbanisation, unsafe exposure to road traffic, motorisation, ageing, environmental and lifestyle changes over the past decades have led to injury being reported as the seventh leading cause of premature death.⁸ Literature indicates a range of risk factors associated with injuries including youth,⁹ low socioeconomic status,¹⁰ those covered by health insurance,¹¹ disabilities related to vision, hearing, motor and cognitive function,¹² male sex,¹¹ college-level education,¹³ presence of comorbidities, alcohol use,¹³ unsafe environment including inadequate traffic laws, poor road engineering and high-risk road users.¹⁴

Reporting of injury data in Nepal is very limited; the majority of the data on injuries are generated from hospital and police records. However, population-based epidemiological data needed to inform policy and programme development are not readily available in Nepal. According to recent estimates, in Nepal, 4.11% of all deaths were attributed to transport injuries, 3.54% were attributed to unintentional injuries, and 1.55% were attributed to self-harm and interpersonal violence.¹⁵ Previous studies conducted by the Nepal Health Research Council and reports from other studies show that RTIs are a common cause of trauma-related injuries in Nepal.^{16 17} Nepal traffic police statistics suggest that the number of road traffic-related death and injury is increasing annually.^{18–20} A community-based survey suggested that injuries occurring at home and around, such as falls and burns, poisoning, and work-related injuries also have a significant impact on Nepal's injury burden.²¹ A recent study recorded an incident rate of 22.40 per 1000 (3 months incidence) for injuries in Makwanpur district. Nearly 36% of all injuries occurred at home and 29% of injuries were work-related.²² The Nepal Demographic and Health Survey (NDHS) 2016 revealed that 9% of women had experienced physical violence from their husbands in the past 12 months.²³ Although data are not available for recent years, the NDHS 2006 found that injury was the cause of death for 2.8% postnatal and 10.7% of child (12–59 months) deaths.²⁴ While many injuries are preventable, there is a lack of attention paid to injury prevention, but it is gradually appearing in Nepal's national health and development agenda.²⁵ The Nepal Road Safety Action Plan 2013–2020,²⁶ the Nepal Health Sector Strategy (NHSS) 2015–2020,²⁷ the NHSS Implementation Plan 2016–2021²⁸ and the Fifteenth Plan (2019–2024)²⁹ have included activities related to the prevention of injuries where road safety and postcrash response are highlighted but provisions for other types of injuries are neglected. A recent review of national documents has indicated that

over 60 different existing laws and policies are relevant to this particular public health issue.³⁰ Limited progress in the implementation of these plans and policies has challenged any efforts related to prevention of injuries in Nepal. Knowing the factors that are associated with injuries is important to encourage actions regarding health promotion and prevention in different contexts. Therefore, this paper reports the prevalence and factors associated with injuries among the adult population of Nepal.

METHODS

This study used secondary data derived from population-based NCD risk factor surveys using a standardised method from the WHO, called the STEPwise Approach to NCD Risk Factor Surveillance (STEPS) Survey Nepal 2019.³¹ The STEPS 2019 Survey collected data using a WHO-recommended optional module on unintentional injuries and violence which is presented in this paper.³²

Study design and sampling

We used nationally representative population-based cross-sectional study data from the STEPS Survey 2019. In brief, the STEPS Survey applied multistage cluster sampling using the Central Bureau of Statistics data to obtain a random sample of 5593 adults aged 15–69 years who have been the usual residents of the household for at least 6 months and have stayed in the household the night before the survey. One participant per household was selected for the survey. The detailed methodology has been presented elsewhere.^{33–35}

Data collection tools and techniques

The survey was conducted using the standardised WHO NCD STEPS Questionnaire V.3.2. Data collection was done using ODK software by trained enumerators using the eSTEPS questionnaire which was loaded on android tablets. As part of the household survey, STEP 1 measures sociodemographic and behavioural information on tobacco and alcohol use, diet, physical activity, history of chronic diseases, family history of chronic conditions, health screening, and healthcare costs. STEP 2 measures physical measurements such as height, weight, blood pressure and waist circumference. STEP 3 measures biochemical tests such as fasting blood glucose, total cholesterol, triglycerides, high-density lipoprotein and low-density lipoprotein cholesterol. Unintentional injuries and violence were covered in STEP 1 as one of the optional modules, of which we used all the questions prescribed in the stepwise approach to surveillance. The data collected from this survey has been deposited in a public repository which is available for reuse on reasonable request. Data can be accessed via the South-East Asia regional microdata repository at <https://nada.searo.who.int/index.php/catalog> with the reference ID NPL_2019_STEPS_v01.³⁶

Primary outcome and measures

The primary outcome was injured person defined as one who had either been injured in a road traffic crash or

collision (RTIs), had other serious unintentional/accidental injury, or had been seriously injured in a violent incident within the past 12 months. Unintentional/accidental injuries included injuries such as a fall, burn, poisoning, near-drowning and electrocution. Violent incidents included intentional use of physical force to cause injury. A serious or severe injury was defined as an injury that required medical attention. These variables included the following questions: ‘did you have any injuries in this road traffic crash which required medical attention?’, ‘in the past 12 months, have you been accidentally injured, other than the RTIs which required medical attention?’ and ‘in the last 12 months, how many times have you been in a violent incident where you were injured and required medical attention?’.

The explanatory variables used in this study include sociodemographic and socioeconomic variables (age, sex, education, marital status, occupation, place of residence, wealth quintile); sex (male, female), marital status (never married, currently married, ever married); age in three categories (15–29, 30–44 and 45–69 years); place of residence stratified into two categories (urban and rural); educational status (none/less than primary, primary, secondary and more than secondary); occupation (categorised as employed, student, homemaker, unemployed and others) and wealth quintile (classified into the following categories: lowest, second, middle, fourth, highest). We also calculated the prevalence of protective and behavioural factors for RTIs, particularly, seat belt use (yes, never, seat belt not present) and reports of driving under the influence of alcohol (no, yes, don’t know). The dependent variable for this study was reporting of any injury within the preceding 12 months.

Statistical analysis

Statistical analysis was performed using STATA V.13 (StataCorp, USA) with appropriate methods for complex sampling design of the survey. Background characteristics for all participants were described, and descriptive summaries for all variables reported. Bivariate analyses were used to examine associations between the explanatory variables and injuries using χ^2 tests with a 95% CI. Likewise, the crude OR (COR), along with its CIs, was estimated using a binary logistic regression model to identify the determinants of injuries. All tests were two-sided and value of $p < 0.05$ was considered statistically significant.

Patient and public involvement

Neither the patients nor the public were involved in the design, conduct, reporting or dissemination plans of this research.

RESULTS

Table 1 depicts the sociodemographic characteristics of research participants and injured persons. A majority (64.21%) of the research participants were women. More men (12.14%) were injured than women (10.97%).

By age, injuries were highest (12.15%) among participants aged 15–29 years and lowest (10.85%) among the 45–69 years age group. In terms of education, injuries were highest (13.61%) among those participants who completed secondary-level education and lowest (9.87%) among those who had primary education. Injuries were lower among the participants residing in metropolitan/submetropolitan cities (9.67%) compared with those who reside in municipalities (10.89%) and rural municipalities (11.74%). By wealth quintile, injuries ranged from 10.20% among participants in the fourth wealth quintile to 12.47% among participants in the second wealth quintile. Injuries were higher (14.35%) among unemployed participants. Regarding marital status, injuries were higher (13.96%) among the ever-married and lower (11.14%) among the currently married participants.

Table 2 depicts the place of occurrence and type of unintentional injuries. Place of injury occurrence was available for 154 participants (27.06%) who reported 205 unintentional injury incidents other than road injuries. Among them, 32.68% of injuries occurred at home followed by 23.41% in the farms, 19.51% at workplaces and 19.02% on roads. Among those injured at home, 83.58% were fall injuries, followed by cut wounds (11.94%), animal bites (2.99%) and burns (1.49%). Likewise, 85% of all unintentional injuries occurring at workplaces were falls followed by near-drowning (7.50%), burns (5%) and cut wounds (2.50%). Among the 39 injuries that occurred on roads/streets/highways, majority were falls (87.18%) followed by animal bites (7.69%) and cut wounds (5.13%). Another common place for injury was farmland where 47.92% of falls and 43.75% of cuts occurred. Other injuries that occurred were animal bites and burns. All the unintentional injuries that occurred in schools and sports or leisure areas were falls.

Table 3 highlights the prevalence of protective and risk factors for road safety. Among adults who used seat belts, only 9.15% reported injury. Likewise, 9.80% of injuries were reported among alcohol users while driving.

Figure 1 shows the prevalence rate of injuries by its type. The most common types of injuries were violent incidents (5.33%), followed by unintentional injuries (4.71%) and RTIs (3.75%).

Online supplemental table 1 shows the bivariate analysis of independent variables with RTI, unintentional injury, intentional injury and overall injury. In bivariate analysis, RTI was found to be associated with wealth quintile and occupation. The middle wealth quintile had almost three times higher odds (COR) of having RTI (COR=2.95, 95% CI 1.27 to 6.84) compared with the lowest wealth quintile. Homemakers had almost half the odds of having RTIs compared with the employed (COR 0.45, 95% CI 0.24 to 0.84). Those who were unemployed had more than two times higher odds of intentional injury compared with those who were employed. Likewise, those currently married (COR=3.74, 95% CI 1.37 to 10.17) and ever married (COR=3.49, 95% CI 1.08 to 11.25) had more than three times higher odds of intentional injury

**Table 1** Sociodemographic characteristics of research participants and injured persons

Category	Injured n (%)	Not injured n (%)	Total Number (%)	P value
Sex (n=4996)				
Male	217 (12.14)	1571 (87.86)	1788 (100)	0.896
Female	352 (10.97)	2856 (89.03)	3208 (100)	
Age groups, years (n=4996)				
15–29	158 (12.15)	1142 (87.85)	1300 (100)	0.295
30–44	207 (11.40)	1608 (88.60)	1815 (100)	
45–69	204 (10.85)	1677 (89.15)	1881 (100)	
Education level (n=4995)*				
None/less than primary	285 (11.62)	2168 (88.38)	2453 (100)	0.898
Primary	94 (9.87)	858 (90.13)	952 (100)	
Secondary	106 (10.89)	867 (89.11)	973 (100)	
More than secondary	84 (13.61)	533 (86.39)	617 (100)	
Place of residence (n=4996)				
Metropolitan/submetropolitan	76 (9.67)	549 (90.33)	625 (100)	0.827
Municipality	292 (10.89)	2147 (89.11)	2439 (100)	
Rural municipality	201 (11.74)	1731 (88.26)	1932 (100)	
Wealth quintile (n=4996)				
Lowest	157 (10.97)	1274 (89.03)	1431 (100)	0.701
Second	116 (12.47)	814 (87.53)	930 (100)	
Middle	106 (12.30)	756 (87.70)	862 (100)	
Fourth	82 (10.20)	722 (89.80)	804 (100)	
Highest	108 (11.15)	861 (88.85)	969 (100)	
Occupation (n=4990)*				
Employed	195 (12.67)	1344 (87.33)	1539 (100)	0.420
Student	46 (13.11)	305 (86.89)	351 (100)	
Homemaker	295 (10.41)	2538 (89.59)	2833 (100)	
Unemployed	30 (14.35)	179 (85.65)	209 (100)	
Others	3 (5.17)	55 (94.83)	58 (100)	
Marital status (n=4995)				
Never married	58 (12.24)	416 (87.76)	474 (100)	0.629
Currently married	474 (11.14)	3782 (88.86)	4256 (100)	
Ever married	37 (13.96)	228 (86.04)	265 (100)	
Total (15–69)	569 (11.39)	4427 (88.61)	4996 (100)	

*6 cases were missing occupation and 1 each missing for education and marital status.

compared with those who were never married. None of the independent variables were found to significantly associate with unintentional injury and overall injury.

DISCUSSION

This population-based study estimated the national-level prevalence of injuries and its associated factors among adults aged 15–69 years. The overall prevalence of injury was found to be 11.39%. More men (12.14%) were injured compared with women (10.97%). Maximum injuries was

seen among the young age group (12.15%) followed by the middle age group (11.40%) and participants with a level of education above secondary (13.61%). However, none of these differences were statistically significant. Similarly, the prevalence rates of RTI, unintentional injury and violent incident were 3.75%, 4.71% and 5.33% respectively. Bivariate analysis showed that RTI was significantly associated with wealth quintile and occupation. Similarly, occupation and marital status were found to be strong predictors of intentional injury.

Table 2 Place of occurrence of injury for other unintentional injuries, n (%)

Place of injury (other than RTI)	Falls n (%)	Burns n (%)	Cuts n (%)	Near-drowning n (%)	Animal bites N (%)	Total N (%)
Home (n=67)	56 (83.58)	1 (1.49)	8 (11.94)	0	2 (2.99)	67 (100)
Farmland (n=48)	23 (47.92)	1 (2.08)	21 (43.75)	0	3 (6.25)	48 (100)
Workplace (n=40)	34 (85.00)	2 (5.00)	1 (2.50)	3 (7.50)	0	40 (100)
Road/street/highway (n=39)	34 (87.18)	0	2 (5.13)	0	3 (7.69)	39 (100)
Sports/leisure area (n=7)	7 (100)	0	0	0	0	7 (100)
School (n=4)	4 (100)	0	0	0	0	4 (100)

RTI, road traffic injury.

The prevalence of injury (11.39%) was consistent with the nationwide studies conducted in Nepal (13.1%)²¹ and Kenya (15%),³⁷ however, we found a higher prevalence compared with a study conducted in eastern Nepal (3.1%).³⁸ This calls for the implementation of laws and regulations aimed at reducing injuries which encompass environmental modifications, addressing barriers to implementing injury-control policies, continuing research for providing evidence for decision making and strengthening injury prevention policies.^{30 39}

Similarly, our findings are in line with other studies conducted in Pakistan,⁴⁰ Kenya,³⁷ Sudan⁴¹ and Tanzania⁴² that have found a higher risk of injuries among men than in women. The gender disparity can be attributed to masculine norms, expectations and identity, occupation: work on various sites, travelling by different modes,⁴³ drinking and driving,⁴⁴ and exposure to risky behaviour.⁴⁵ This calls for strengthened policies for safety promotion, especially in areas of occupational injuries and road safety. Findings from the burden of injuries in Nepal, 1990–2017 revealed that young and middle-aged adults experience a greater burden of injuries than other age groups, which is convergent to our results.¹⁵ Furthermore, a systematic analysis for the Global Burden of Disease, 2019 showed that three injury causes: road injuries, self-harm and interpersonal violence were among the top causes of DALYs in adolescents aged 10–24 years which is also in agreement with our findings.⁴⁶ A possible explanation could be that

the young age group people have risk-taking lifestyles and behaviours. The burden of injury in this age group might have a greater impact on the social and economic development of the country. Place of residence is another major factor for injury. We found that participants residing in metropolitan/submetropolitan cities were more injured as compared with those who resided in municipalities and rural municipalities. The result is similar to that of a study done in Nepal⁴⁷ in 2015, however, it contradicts the findings from USA,⁴⁸ Kenya³⁷ and Tanzania.⁴² Possible explanations might be unplanned urbanisation, high population, and the greater number of vehicles in metropolitan and submetropolitan cities of Nepal.

Bivariate analysis showed that the odds of having RTIs were almost three times higher (COR=2.95, 95% CI 1.27 to 6.84) in middle wealth quintile participants than in lower wealth quintile participants. This is an interesting finding because most of the previous studies have found that people from poorer economic backgrounds have higher rates of injury and death in comparison to wealthier people.^{4 49 50} A possible explanation could be that participants in the middle wealth quintile could afford two-wheeler vehicles and access to other means that may increase the risk of injury. Also, middle wealth quintile participants are active and engaged in occupations that may increase their chances of encountering occupational injury as compared with the participants in the lowest quintile. Likewise, participants who were

Table 3 Prevalence of protective and behavioural risk factors for road safety, n (%)

Protective and risk factors	Road traffic injuries		
	No n (%)	Yes n (%)	Total n (%)
Seat belt use			
Yes	139 (90.85)	14 (9.15)	153 (100)
Never	1367 (96.88)	44 (3.12)	1411 (100)
Seat belt not present	1799 (96.46)	66 (3.54)	1865 (100)
Drove vehicle under the influence of alcohol			
No	1417 (96.72)	48 (3.28)	1465 (100)
Yes	92 (90.20)	10 (9.80)	102 (100)
Don't know	3777 (96.80)	125 (3.20)	3902 (100)




Figure 1 Prevalence rate of different types of injuries. Y-axis represent percentage

homemakers were less likely to be involved in road traffic accidents compared with participants who were employed (COR=0.45, 95% CI 0.24 to 0.84).

Unintentional injuries are predictable and preventable. Cost-effective and equity-oriented interventions such as setting minimum conditions and product standards, imposing safe behaviour and practices through legislation, regulation and enforcement would be promising in mitigating injuries.⁵¹ Also, the study revealed that participants who were currently married (COR=3.74, 95% CI 1.37 to 10.17) and ever married (COR=3.49, 95% CI 1.08 to 11.25) had more than three times higher odds of having unintentional injuries compared with those who were never married. Ever married people might suffer from varying levels of psychological stress and those who are currently married have to adjust to several changes such as changes in child care, housework and work demands. These underlying conditions might compromise safety practices to prevent injuries. Although legislation for key risk factors such as overspeeding, drinking and driving, use of seat belts, motorcycle helmet use exists,⁵² comprehensive injury surveillance and concerted multisectoral approaches are needed for implementing existing provisions.

The data included in our study are from a large replicative sample of the general population and quality control was rigorous, increasing the generalisability. However, our study had several limitations. First, the injuries were self-reported, which might have led to recall bias leading to underestimation or overestimation of injuries and affected the actual prevalence. Second, the study was cross-sectional in nature and, therefore, it was unable to tell the direction of association. Third, the study did not take into account financial and social effects of injuries, DALYs, and YLD and mortality. Furthermore, in line with similar studies conducted in Kenya,³⁷ Sierra Leone⁵³ and Bangladesh,⁵⁴ some of the system-level components (such as conditions of the roads, management and control of traffic, traffic regulations, etc) could not be incorporated as this was a secondary analysis of the STEPS Survey, Nepal, 2019. Despite its limitations, this is the

first step towards gathering information that could be useful for injury-prevention interventions and policies at the national, provincial and local levels. Further studies are recommended to incorporate all the comprehensive national information of injuries.

CONCLUSION

This study showed that injuries are a major national public health problem and provides a baseline estimate of injury prevalence in one of the world's disaster-prone countries. Men, those in the younger age group and participants with higher education were seen to be more vulnerable to injuries—requiring special targeted injury-prevention initiatives for these groups. Further research is needed to investigate the context-specific behavioural, environmental and system-level components, and the injury circumstances that shape them. The study also calls for collaborative efforts across various disciplines so that cost-effective and equity-oriented interventions can be employed for safety improvements and behavioural promotions.

Author affiliations

¹Research section, Nepal Health Research Council, Ramshah path, Kathmandu, Bagmati province, Nepal

²Department of social sciences, Nexus Institute of Research and Innovation (NIRI), Kathmandu, Bagmati province, Nepal

Twitter Meghnath Dhimal @MeghnathDhimal, Anil Poudyal anil poudyal @PoudyalAnil, Bihungum Bista @bistabihungum and Puspa Raj Pant Puspa Raj Pant, PhD @Puspa_RPant

Acknowledgements The authors thank Dr Umesh Raj Aryal, Dr Dinesh Bhandari, consultants Nepal Health Research Council (NHRC) and Ms Hemlata Pant for reviewing the manuscript.

Contributors MD, AP and BB were responsible for the concept and design, analysis and interpretation of data, and writing the manuscript. AP wrote the first draft of the manuscript. PG, PRP and SD revised and edited the manuscript. All authors provided intellectual content during manuscript drafting and revision, accept accountability for their contributions and have approved the final manuscript. AP is the guarantors and undertake the full responsibility for the contents of the article submitted for publication. MD and AP contributed equally.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval The study fully abided by ethical guidelines, with an ethical approval from an independent Ethical Review Board (ERB) of the Nepal Health Research Council (registration number, 293/2018). Right to voluntary participation and freedom to withdraw was fully assured by taking written informed consent from the study participants.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. Data are available upon reasonable request. The data collected from this survey has been deposited in a public repository which is available for reuse upon reasonable request. Data can be accessed via the South-east Asia regional microdata repository at <https://nada.searo.who.int/index.php/catalog> with the reference ID NPL_2019_STEPS_v01

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been

peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Meghnath Dhimal <http://orcid.org/0000-0001-7176-7821>

Anil Poudyal <http://orcid.org/0000-0003-2687-2993>

REFERENCES

- James SL, Castle CD, Dingels ZV, *et al*. Global injury morbidity and mortality from 1990 to 2017: results from the global burden of disease study 2017. *Inj Prev* 2020;26:196–114.
- World Health Organization. Injuries and violence [Internet], 2021. Available: <https://www.who.int/news-room/fact-sheets/detail/injuries-and-violence>
- RothGA, AbateD, AbateKH, *et al*. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the global burden of disease study 2017. *Lancet* 2018;392:1736–88.
- World Health Organization. Violence and Injuries: the Facts. 2014 [Internet], 2014. Available: http://apps.who.int/iris/bitstream/10665/44288/1/9789241599375_eng.pdf%0A3
- Gosselin RA, Spiegel DA, Coughlin R, *et al*. Injuries: the neglected burden in developing countries. *Bull World Health Organ* 2009;87:246–246a.
- Chandran A, Hyder AA, Peek-Asa C. The global burden of unintentional injuries and an agenda for progress. *Epidemiol Rev* 2010;32:110–20.
- Department of Health Services. Annual report 2077/78 (2020/21) [Internet]. Kathmandu. Available: https://dohs.gov.np/wp-content/uploads/2022/07/DoHS-Annual-Report-FY-2077-78-date-5-July-2022-2022_FINAL.pdf
- Nepal Health Research Council (NHRC);Ministry of Health and Population; Monitoring Evaluation and Operational Research. Nepal Burden of Disease 2017: A Country Report based on the Global Burden of Disease 2017 Study [Internet], 2017. Available: http://nhrc.gov.np/wp-content/uploads/2019/04/NBoD-2017_NHRC-MoHP.pdf
- Thomas J, Kavanagh J, Tucker H. Accidental injury risk-taking behaviour and the social circumstances in which young people (aged 12–24) live: a systematic review 2007.
- Whitlock G, Norton R, Clark T, *et al*. Motor vehicle driver injury and socioeconomic status: a cohort study with prospective and retrospective driver injuries. *J Epidemiol Community Health* 2003;57:512–6.
- Moncater TJR, Nakamura K, Siongo KL, *et al*. Prevalence and determinants of self-reported injuries among community-dwelling older adults in the Philippines: a 10-year pooled analysis. *Int J Environ Res Public Health* 2020;17:4372.
- Muir SW, Gopaul K, Montero Odasso MM. The role of cognitive impairment in fall risk among older adults: a systematic review and meta-analysis. *Age Ageing* 2012;41:299–308.
- Diamond MB, Dalal S, Adebamowo C, *et al*. Prevalence and risk factor for injury in sub-Saharan Africa: a multicountry study. *Inj Prev* 2018;24:272–8.
- SEARO WHO, Forum RH. *Who south-east Asia region special issue on blood Pressure-take control*. New Delhi: WHO SEARO, 2013.
- Pant PR, Banstola A, Bhatta S, *et al*. Burden of injuries in Nepal, 1990–2017: findings from the global burden of disease study 2017. *Inj Prev* 2020;26:i57–66.
- Mishra SR, Neupane D, Bhandari PM. Burgeoning burden of non-communicable diseases in Nepal: a scoping review. *Global Health* 2015;11:1–10.
- Shrestha R, Shrestha SK, Kayastha SR, *et al*. A comparative study on epidemiology, spectrum and outcome analysis of physical trauma cases presenting to emergency department of Dhulikhel Hospital, Kathmandu university hospital and its outreach centers in rural area. *Kathmandu Univ Med J* 2013;11:241–6.
- World bank. Delivering Road Safety in Nepal [Internet], 2020. Available: <https://openknowledge.worldbank.org/bitstream/handle/10986/33340/ROAD+SAFETY+IN+NEPAL.pdf;jsessionid=E0799429DC03F4F5F203FB15EEE4E7CB?sequence=1>
- Choulagai B, Ling H, Sharma P. Epidemiology of road traffic accidents in Nepal: data review and qualitative analysis. *SM J Public Heal Epidemiol* 2015;1:1014.
- Karkee R, Lee AH. Epidemiology of road traffic injuries in Nepal, 2001–2013: systematic review and secondary data analysis. *BMJ Open* 2016;6:e010757.
- Gupta S, Gupta SK, Devkota S, *et al*. Fall injuries in Nepal: a countrywide population-based survey. *Ann Glob Health* 2015;81:487–94.
- Pant PR, Deave T, Banstola A, *et al*. Home-related and work-related injuries in Makwanpur district, Nepal: a household survey. *Inj Prev* 2021;27:450–5.
- Ministry of Health; New ERA; and ICF. Nepal Demographic and Health Survey 2016 [Internet], 2016. Available: <https://www.dhsprogram.com/pubs/pdf/fr336/fr336.pdf>
- Ministry of Health; New ERA; and ICF. Nepal Demographic Health Survey 2006 [Internet], 2006. Available: <http://www.measuredhs.com/pubs/pdf/FR191/FR191.pdf>
- Shah MT, Bhattarai S, Lamichhane N, *et al*. Assessment of the availability of technology for trauma care in Nepal. *Injury* 2015;46:1712–9.
- Government of Nepal;Ministry of Physical Planning & Transport Management. Nepal Road Safety Action Plan (2013 – 2020) [Internet]. Kathmandu, Nepal. Available: http://www.nhss.gov.np/NHSS_Archives/health_policy/NHSS_english_book_2015.pdf
- Government of Nepal;Ministry of Health and Population. NEPAL HEALTH SECTOR STRATEGY 2015 - 2020 [Internet]., 2015. Available: http://www.nhssp.org.np/NHSSP_Archives/health_policy/NHSS_english_book_2015.pdf
- Government of Nepal;Ministry of Health and Population. Nepal Health Sector Strategy Implementation Plan 2016–2021 [Internet]. Available: <http://climate.mohp.gov.np/news/31-acts/163-nepal-health-sector-strategy-implementation-plan-2016-2021>
- Government of Nepal; National Planning Commission. The Fifteenth Plan (Fiscal Year 2019/20 – 2023/24) [Internet]. Available: https://npc.gov.np/images/category/15th_plan_English_Version.pdf
- Pant PR, Mytton J, Dharel MR, *et al*. The prevention of - and first response to - injuries in Nepal: a review of policies and legislation. *Health Res Policy Syst* 2021;19:1–20.
- World Health Organization. STEPwise Approach to NCD Risk Factor Surveillance (STEPS) [Internet]. Available: <https://www.who.int/teams/noncommunicable-diseases/surveillance/systems-tools/steps>
- World Health Organization (WHO). Noncommunicable diseases and their risk factors- STEPS Optional Modules [Internet]. Available: <https://www.who.int/ncds/surveillance/steps/riskfactor/modules/en/>
- Dhimal M, Bista B, Bhattarai S, *et al*. report on Non Communicable Diseases Risk Factors: STEPS Survey Nepal 2019 [Internet], 2020. Available: https://www.who.int/docs/default-source/nepal-documents/ncds/ncd-steps-survey-2019-compressed.pdf?sfvrsn=807bc4c6_2
- Bista B, Dhimal M, Bhattarai S, *et al*. Prevalence of non-communicable diseases risk factors and their determinants: results from steps survey 2019, Nepal. *PLoS One* 2021;16:e0253605.
- Poudyal A, Bista B, Gyanwali P, *et al*. Prevalence and factors associated with joint pain in Nepal: findings from a countrywide cross-sectional steps survey. *BMJ Open* 2021;11:e051536.
- Nepal Health Research Council. Data from:Nepal - STEPS 2019. WHO South-East Asia Regional Microdata Repository [Internet], 2019. Available: <https://nada.searo.who.int/index.php/catalog/76> [Accessed 20 Oct 2021].
- Gathecha GK, Ngaruiya C, Mwai W, *et al*. Prevalence and predictors of injuries in Kenya: findings from the National steps survey. *BMC Public Health* 2018;18:1–9.
- Khanal VK, Upreti R, Oli U, *et al*. Prevalence of injury and its associated factors in a rural area of eastern Nepal. *J Chitwan Med Coll* 2016;6:7–13.
- Scott I. The future of injury prevention in Asia and the Pacific. *Inj Control Saf Promot* 2004;11:201–6.
- Hyder AA, Razzak JA. The challenges of injuries and trauma in Pakistan: an opportunity for concerted action. *Public Health* 2013;127:699–703.
- El Tayeb S, Abdalla S, Mørkve O, *et al*. Injuries in Khartoum state, the Sudan: a household survey of incidence and risk factors. *Int J Inj Contr Saf Promot* 2014;21:144–53.



- 42 Moshiro C, Heuch I, Åström AN, *et al.* Injury morbidity in an urban and a rural area in Tanzania: an epidemiological survey. *BMC Public Health* 2005;5:1–10.
- 43 Ghaffar A, Hyder AA, Mastoor MI, *et al.* Injuries in Pakistan: directions for future health policy. *Health Policy Plan* 1999;14:11–17.
- 44 Behnood A, Roshandeh AM, Mannering FL. Latent class analysis of the effects of age, gender, and alcohol consumption on driver-injury severities. *Anal Methods Accid Res* 2014;3-4:56–91.
- 45 Udry JR. Why are males injured more than females? *Inj Prev* 1998;4:94–5.
- 46 Vos T, Lim SS, Abbafati C, *et al.* Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the global burden of disease study 2019. *Lancet* 2020;396:1204–22.
- 47 Gupta S, Wong EG, Nepal S, *et al.* Injury prevalence and causality in developing nations: results from a countrywide population-based survey in Nepal. *Surgery* 2015;157:843–9.
- 48 Tiesman H, Zwerling C, Peek-Asa C, *et al.* Non-fatal injuries among urban and rural residents: the National health interview survey, 1997-2001. *Inj Prev* 2007;13:115–9.
- 49 Giashuddin SM, Rahman A, Rahman F, *et al.* Socioeconomic inequality in child injury in Bangladesh - implication for developing countries. *Int J Equity Health* 2009;8:7–8.
- 50 Cubbin C, Smith GS. Socioeconomic inequalities in injury: critical issues in design and analysis. *Annu Rev Public Health* 2002;23:349–75.
- 51 Zambon F, Injuries LB. And inequities: guidance for addressing inequities in unintentional injuries. *World Health Organization. Regional Office for Europe* 2014.
- 52 Government of Nepal. Motor Vehicles and Transport Management Rules, 2054 [Internet]. Available: <https://www.lawcommission.gov.np/en/wp-content/uploads/2018/09/motor-vehicles-and-transport-management-rules-2054-1997.pdf>
- 53 Stewart K-AA, Groen RS, Kamara TB, *et al.* Traumatic injuries in developing countries: report from a nationwide cross-sectional survey of Sierra Leone. *JAMA Surg* 2013;148:463–9.
- 54 UI Baset MK, Rahman A, Alonge O, *et al.* Pattern of road traffic injuries in rural Bangladesh: burden estimates and risk factors. *Int J Environ Res Public Health* 2017;14:1354.