

BMJ Open Effects of respiratory physiotherapy interventions on pulmonary mechanics of newborns: a protocol for a systematic review

Adriele de Moraes Nunes ¹,
Ana Tereza do Nascimento Sales Figueiredo Fernandes,²
Adynna Tévina de Castro Silva,³ Mayara Fabiana Pereira Costa,¹
Karolinne Souza Monteiro,¹ Silvana Alves Pereira ^{1,3}

To cite: Nunes AdM, Fernandes ATdNSF, de Castro Silva AT, *et al.* Effects of respiratory physiotherapy interventions on pulmonary mechanics of newborns: a protocol for a systematic review. *BMJ Open* 2022;**12**:e062910. doi:10.1136/bmjopen-2022-062910

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-062910>).

Received 15 March 2022
Accepted 03 August 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Postgraduate program in Rehabilitation Sciences, Universidade Federal do Rio Grande do Norte, Santa Cruz, RN, Brazil

²Departamento de Fisioterapia, Universidade Estadual da Paraíba, Campina Grande, PB, Brazil

³Maternidade Escola Januario Cicco - Ebserh, Universidade Federal do Rio Grande do Norte, Natal, RN, Brazil

Correspondence to

Dr Silvana Alves Pereira;
silvana.alves@ufrn.br

ABSTRACT

Introduction Although respiratory physiotherapy techniques may reduce respiratory load in newborns, manual contact with the ribcage may interfere with pulmonary mechanics. Therefore, this systematic review aims to evaluate the effects of conventional and non-conventional respiratory physiotherapies on pulmonary mechanics of newborns.

Methods and analysis We will search PubMed, LILACS, SciELO, ScienceDirect, Cochrane Central and Web of Science databases. Searches will be conducted from September 2022. We will include randomised clinical trials reporting thoracoabdominal synchrony, lung volumes and capacities, respiratory discomfort and pain in newborns aged between 1 hour and 28 days and admitted to neonatal intensive care units. We will exclude studies not fully available or incomplete and studies conducted with newborns presenting structural alterations. Two independent researchers will perform the study selection, data extraction and quality assessment. After consensus, one reviewer will proceed with the process. We will include studies published in English or Portuguese, without publication date restriction. An overview of the included studies and extracted information will be reported and the quality of studies will be assessed. A meta-analysis will be conducted if data regarding between-group comparisons are available.

Ethics and dissemination Ethics approval is not required for this systematic review. Results will be presented in journals and national and international conferences, and findings will be shared on social media using accessible language.

PROSPERO registration number CRD42021266729.

INTRODUCTION

Newborns present unique physiological and anatomical characteristics, such as fewer alveoli, more cartilaginous and horizontal ribs and greater airway resistance and ribcage compliance. Therefore, this population is more susceptible to muscle fatigue, requiring

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This systematic review protocol was prepared according to international recommendations.
- ⇒ Study inclusion and quality will be assessed independently by two reviewers.
- ⇒ Including studies published only in English and Portuguese may limit the results.
- ⇒ The study findings will provide comprehensive information for future randomised controlled trials about the influence of respiratory physiotherapy on pulmonary mechanics of newborns.

ventilatory support, hospitalisation and respiratory physiotherapy.^{1–3}

Respiratory physiotherapy uses techniques to improve respiratory function, facilitate gas exchange, prevent and treat pulmonary complications and improve ventilation–perfusion, mucociliary clearance and secretion removal.³ These techniques can be conventional or non-conventional, depending on the initial contact with the ribcage.⁴ Conventional techniques are usually older than non-conventional techniques, and the contact with the ribcage does not always consider pulmonary mechanics. In contrast, non-conventional techniques are usually employed according to anatomical and physiological differences of each age group.^{4–6}

Although conventional and non-conventional physiotherapy are commonly used in neonatal intensive care units to improve secretion mobilisation and thoracoabdominal repercussion,^{5–12} the effects on pulmonary mechanics of newborns are still unclear. Therefore, this systematic review aims to evaluate the effects of conventional and non-conventional respiratory physiotherapies, respiratory pattern, pulmonary

mechanics and pain in newborns and identify protocols for respiratory physiotherapy used in this population.

METHODS AND ANALYSIS

Registration

This protocol followed the Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P)¹³ and was registered in the International Prospective Register of Systematic Reviews (PROSPERO) in August 2021.

Eligibility criteria

Types of studies

We will include randomised controlled trials published in full. Studies not fully available or incomplete will be excluded.

Types of participants

We will include studies performed with newborns of both sexes, aged between 1 hour and 28 days, and admitted to neonatal intensive care units. We will exclude studies conducted with animals or newborns presenting structural alterations (eg, chest wall deformities, abdominal pathologies and cardiac diseases).

Types of interventions

We will include studies comparing conventional (vibration or vibrocompression, percussion and postural drainage) with non-conventional physiotherapy techniques (assisted autogenic drainage, increased expiratory flow, prolonged slow expiration and thoracoabdominal rebalancing).^{4,6} All techniques are described as follows.

Vibrocompression and vibropercussion: bronchial hygiene manoeuvres with thixotropic effects caused by the propagation of mechanic waves on the chest wall. These waves allow secretion mobilisation and facilitate its elimination.⁴

Postural drainage: bronchial hygiene manoeuvre that uses gravity to drain pulmonary secretion to the centre of the bronchial tree.⁴

Assisted autogenic drainage: the flow generated during the manoeuvre is altered and reaches the small airways and pulmonary periphery, interfering with the depuration process.⁴

Increased expiratory flow: bronchial hygiene manoeuvre consisting of changes in thoracoabdominal movements promoted by the hands of the therapist with the newborn positioned supine. The technique is applied during the expiratory phase to increase the expiratory flow of the newborn.⁴

Prolonged slow expiration: bronchial hygiene manoeuvre that consists of prolonging spontaneous breathing by manually pressing the chest and abdomen during expiration. The manual pressure must be applied continuously and synchronously in both regions to stimulate the expiratory flow and mobilise secretions.^{14 15}

Thoracoabdominal rebalancing: the technique consists of manual support on different points of the thoracoabdominal region (eg, floating ribs and sternum) to assist ribcage movement and mobilization of the thorax.¹⁰

Types of outcome measures

Primary outcomes

1. Lung compliance and airway resistance: variables assessed in the mechanical ventilator using the following formulas: $C_{dyn} = VT/PIPEEP$; $C_{stat} = VT/P_{plateau} - PEEP$; $C_{raw} = PI - P_{plateau}/\text{inspiratory flow}$ C_{dyn} , dynamic compliance; VT, tidal volume; PI, inspiratory pressure; PEEP, Positive End Expiratory Pressure; C_{stat} , static compliance C_{raw} , airway resistance; $P_{plateau}$, Plateau Pressure.
2. Lung volumes and capacities: pneumotachograph coupled to a flow transducer.
3. Thoracoabdominal synchrony: observation of the evaluator, Silverman Andersen Respiratory Severity Score and methods for assessing thoracoabdominal movement (eg, plethysmography and biophotogrammetry).

Secondary outcomes

1. Respiratory distress: the instruments used to assess respiratory distress will not be limited, but dichotomous answers (eg, yes or no) will be considered during data extraction.
2. Pain: the instruments used to assess pain will not be limited, but dichotomous answers (eg, yes or no) will be considered during data extraction.
3. Heart rate: monitored using multiparametric monitors.
4. Respiratory rate: respiratory rate (newborns under mechanical ventilation) and respiratory incursions within 1 min assessed visually or using multiparametric monitors.
5. Peripheral oxygen saturation: monitored using multiparametric monitors.

Information sources

Search strategy

Searches will be conducted from September 2022 using PubMed, LILACS, SciELO, ScienceDirect, Cochrane Central and Web of Science databases. The reference list of all included studies and relevant systematic reviews identified in the search will be screened for additional eligible studies. Finally, experts in the field will be asked for further eligible studies.

A detailed description of the search strategies is provided in the online supplemental material. The main terms, synonyms and relevant examples were identified and included in the search using the Boolean operator OR. Articles in English and Portuguese will be included with no publication date restriction. The search will be conducted in titles, abstracts and main body of the texts. To perform an up-to-date review, the search will be redone before submission and potential, new released and relevant articles will be incorporated.

Study selection

The search results will be imported into the reference list management tool Mendeley (<https://www.mendeley.com>). Any duplicates will be identified and removed using Mendeley. Then, the reference list will be exported to the Rayyan QCRI systematic review web-based application (<https://rayyan.qcri.org>). We will record the selection process in sufficient detail to complete a PRISMA flow diagram.

Two review authors (AdMN and ATdCS) will independently screen the titles and abstracts of the remaining search results and code them as 'retrieve' (eligible or potentially eligible/unclear) or 'do not retrieve'. We will retrieve the full-text study reports of all potentially eligible studies, and two review authors (AdMN and ATdNSFF) will independently screen for inclusion and record the reasons for exclusion of ineligible studies. We will resolve any disagreement through discussion or, if required, we will consult a third review author (SAP). We will identify and exclude duplicates and collate multiple reports of the same study so that each study, rather than each report, is the unit of interest in the review.

Data extraction

We will extract data from all included studies using a prepiloted form. Two authors (AdMN and MFPC) will extract data, and two authors (KSM and SAP) will verify data accuracy and integrity. We will extract the following characteristics from included studies:

1. Participants: age, sex and geographical location.
2. Methods: study design, location, sample size and year.
3. Interventions: intervention, comparison, duration of intervention and method of delivery.
4. Outcomes: primary and secondary outcomes.
5. Limitations.

Quality assessment

Two reviewers (AdMN and ATdCS) will independently assess the risk of bias using the following domains of the PEDro scale: eligibility criteria, random allocation, concealed allocation, baseline comparability, blinding of participants, blinding of therapists, blinding of assessors, adequate follow-up, intention-to-treat analysis, between-group comparisons and measures of variability. The total scores of the scale range from 0 to 10, and risk of bias will be interpreted as high (0–3), moderate (5–7) or low (8–10).¹⁶ Results will be presented in a table containing the name of authors and scores obtained on each item of the PEDro Scale.

Data analysis

The characteristics of interventions will be described and presented as mean and SD. A meta-analysis will be conducted using the Review Manager software (Copenhagen: The Nordic Cochrane Collaboration) if data regarding between-group comparisons are available.

The heterogeneity of studies will be assessed by quantifying the proportion of total variability of the estimates

(I^2). I^2 ranges from 0% to 100%, in which 0% indicates lack of variability, >25% low variability, >50% moderate variability and >75% high variability. A significance level of $p < 0.10$ will be considered statistically significant for I^2 . Effect sizes (Z value) will also be assessed, and $p < 0.05$ will be considered statistically significant.¹⁷

Certainty of evidence

Two authors (ATdNSFF and KSM) will independently assess the reliability of evidence using the Grading of Recommendations Assessment, Development and Evaluation.¹⁸ Results will be presented in tables and classified into high, moderate, low or very low.

Study timeline

This review will be conducted in 8–12 months, contemplating all steps described above.

Patient and public involvement

None.

ETHICS AND DISSEMINATION

Ethics approval is not required for this systematic review. The findings from this review will be submitted for publication in a scientific journal and presentation at national and international conferences. Findings will also be shared on social media using accessible language.

Twitter Silvana Alves Pereira @PTPereiraSA

Contributors All authors contributed substantially to the study design, development of inclusion criteria and search strategies and read and approved the final version of the protocol. AdMN developed, designed and registered the protocol in the PROSPERO database. SAP, ATdNSFF, KSM, ATdCS and MFPC provided critical insights and reviewed the protocol.

Funding The study was funded by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brazil (Finance Code 001).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Not applicable to our type of study.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Adrielle de Moraes Nunes <http://orcid.org/0000-0002-2802-4093>Silvana Alves Pereira <http://orcid.org/0000-0002-6226-2837>

REFERENCES

- 1 Newborn health care, Guide for health professionals, Ministry of Health, Health Care Secretariat. *Departamento de Ações Programáticas Estratégicas*. 3. 2, 2014.
- 2 Fiorenzano DM, Leal GN, Sawamura KSS, et al. Respiratory distress syndrome: influence of management on the hemodynamic status of ≤ 32 -week preterm infants in the first 24 hours of life. *Rev Bras Ter Intensiva* 2019;31:312.
- 3 Marques C, Lúcia A, NICOLAU, LAHÓZ. Fisioterapia respiratória em terapia intensiva pediátrica E neonatal: uma revisão baseada em evidências Pediatria (São Paulo); 2007: 216–21.
- 4 Dantas Gomes Évelim Leal de Freitas. Evidência científica das técnicas atuais E convencionais de fisioterapia respiratória em pediatria. *Fisioterapia Brasil* 2016;17:88–97.
- 5 Santos AK, Silveira J, Neves VC, Karoline A, et al. Atelectasis and lung changes in preterm neonates in the neonatal period: a blind radiological report and clinical findings. *Rev Bras Ter Intensiva* 2019;31:347–53.
- 6 Guerra JI de A, Nagem DAP, Moran CA, et al. Thoracoabdominal mobility evaluation by photogrammetry in newborns after expiratory flow increase technique. *Fisioterapia em Movimento* 2017;30:789–95.
- 7 Santos SN, RIBEIRO. Thoracoabdominal interaction and its relationship with biological risk factors in premature newborns. *Fisioterapia e Pesquisa* 2020;27:155–60.
- 8 Brennan C, Ulm L, Julian S, et al. Thoracoabdominal asynchrony is not associated with oxyhemoglobin saturation in recovering premature infants. *Neonatology* 2017;111:297–302.
- 9 OLIVEIRA DE, Costa T, et al. Bronchic hygiene techniques in newborns and infants in the intensive care unit: systematic review of clinical trials. *Revista Pesquisa em Fisioterapia* 2018;8:420–9.
- 10 Rousseno KR, Scalco JC, Rosa GJ da, et al. Reequilíbrio tóraco-abdominal em recém-nascidos prematuros: efeitos em parâmetros cardiorespiratórios, no comportamento, Na DOR E no desconforto respiratório. *Acta Fisiátrica* 2013;20:118–23.
- 11 de Abreu L, Valenti VE, de Oliveira AG, et al. Chest associated to motor physiotherapy improves cardiovascular variables in newborns with respiratory distress syndrome. *Int Arch Med* 2011;4:37–6.
- 12 Skilhan C, Rodrigues L, ALMEIDA DE. A atuação do profissional fisioterapeuta em unidades de terapia intensiva neonatal. *Cinergis* 2016;17.
- 13 SHAMSEER, Larissa. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *Bmj* 2015;349.
- 14 Samuel A, Mishra R, Dasgupta A. Effect of prolonged slow expiratory technique as an adjunct to pulmonary rehabilitation in resolving pulmonary congestion in neonates with congenital pneumonia. *J Clin Neonatol* 2020;9:82.
- 15 Samuel A, Mishra R, et al. Prolonged slow expiratory technique: a lung clearance technique for neonates admitted in neonatal intensive care units. *J Clin Neonatol* 2018;7:282.
- 16 Shiwa SR, Costa LOP, Moser AD de L, et al. PEDro: a base de dados de evidências em fisioterapia. *Fisioterapia em Movimento* 2011;24:523–33.
- 17 Higgins JPT, Thomas J, Chandler J, eds. *Cochrane Handbook for Systematic Reviews of Interventions version 6.3*. Cochrane, 2022. www.training.cochrane.org/handbook
- 18 Lewin S, Booth A, Glenton C, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings: introduction to the series. *Implement Sci* 2018;13:2.

SUPPLEMENTARY FILE

SEARCH STRATEGY

<i>QUESTION OR PROBLEM (PICO)</i>	<i>DESCRIPTION</i>	<i>MERSH TERMS</i>	<i>ENTRY TERMS E KEYWORDS</i>
P - POPULATION	Newborns	Newborn	Newborn, Newborns, Newborn Infant, Newborn Infants, Neonate, Neonates
I - INTERVENTION	Not conventional respiratory physiotherapy	Physical Therapy Modalities, Physical Therapy Specialty, Respiratory Therapy	Physical Therapy Modality, Physical Therapy Modalities, Physiotherapy (Techniques), Physiotherapies (Techniques), Physical Therapy Techniques, Physical Therapy Technique, Group Physiotherapy, Group Physiotherapies, Physical Therapy, Physical Therapies, Physical Therapy Specialty, Physiotherapy Specialty, Respiratory Therapy, Respiratory Therapies, Chest Physical Therapy, Conventional Chest Physical Therapy, Chest Physiotherapy, Physiotherapy Respiratory, Physiotherapy, Autogenic Drainage, Expiratory Flow Increase Technique, Prolonged Slow Expiratory Maneuver, Technique of Re-educating Thoracic and Abdominal, Chest Percussion, Shaking, Tapping
C - CONTROL	Conventional respiratory physiotherapy		

O - OUTCOME	Respiratory Mechanics, Lung Volumes and Capacities, Secretion, Pain, Heart Rate, Respiratory Rate	Respiratory Mechanics, Lung Volume Measurements, Sputum, Pain, Heart Rate, Respiratory Rate	Respiratory Mechanic, Respiratory Mechanics, Breathing Mechanics, Breathing Mechanic, Lung Volume Measurement, Lung Volumes, Lung Volume, Lung Capacities, Lung Capacity, Sputum, Sputums, Induced Sputum, Induced Sputums, Pain, Heart Rate, Heart Rates, Cardiac Rate, Cardiac Rates, Pulse Rate, Pulse Rates, Heartbeat, Heartbeats, Cardiac Chronotropy, Cardiac Chronotropism, Heart Rate Control, Respiratory Rate, Respiratory Rates, Respiration Rate, Respiration Rates Lung Mechanic, Lung Mechanic, Pulmonary Mechanic, Pulmonary Mechanics, Thoracoabdominal Mobility
--------------------	---	---	--

BASE STRATEGY:

(Newborn OR Newborns OR Newborn Infant OR Newborn Infants OR Neonate OR Neonates) AND (Physical Therapy Modality OR Physical Therapy Modalities OR Physiotherapy (Techniques) OR Physiotherapies (Techniques) OR Physical Therapy Techniques OR Physical Therapy Technique OR Group Physiotherapy OR Group Physiotherapies OR Physical Therapy OR Physical Therapies OR Physical Therapy Specialty OR Physiotherapy Specialty OR Respiratory Therapy OR Respiratory Therapies OR Chest Physical Therapy OR Conventional Chest Physical Therapy OR Chest Physiotherapy OR Physiotherapy Respiratory OR Physiotherapy OR Autogenic Drainage OR Expiratory Flow Increase Technique OR Prolonged Slow Expiratory Maneuver OR Technique of Re-educating Thoracic and Abdominal OR Chest Percussion OR Shaking OR Tapping) AND (Respiratory Mechanic OR Respiratory Mechanics OR Breathing Mechanics OR Breathing Mechanic OR Lung Volume Measurement OR Lung Volumes OR Lung Volume OR Lung Capacities OR Lung Capacity OR Sputum OR Sputums OR Induced Sputum OR Induced Sputums OR Pain OR Heart Rate OR Heart Rates OR Cardiac Rate OR Cardiac Rates OR Pulse Rate OR Pulse Rates OR Heartbeat

OR Heartbeats OR Cardiac Chronotropy OR Cardiac Chronotropism OR Heart Rate Control OR Respiratory Rate OR Respiratory Rates OR Respiration Rate OR Respiration Rates OR Lung Mechanic OR Lung Mechanic OR Pulmonary Mechanic OR Pulmonary Mechanics OR Thoracoabdominal Mobility)

STRATEGIES IN DATABASES

PUBMED

(Newborn OR Newborns OR Newborn Infant OR Newborn Infants OR Neonate OR Neonates) AND (Physical Therapy Modality OR Physical Therapy Modalities OR Physiotherapy (Techniques) OR Physiotherapies (Techniques) OR Physical Therapy Techniques OR Physical Therapy Technique OR Group Physiotherapy OR Group Physiotherapies OR Physical Therapy OR Physical Therapies OR Physical Therapy Specialty OR Physiotherapy Specialty OR Respiratory Therapy OR Respiratory Therapies OR Chest Physical Therapy OR Conventional Chest Physical Therapy OR Chest Physiotherapy OR Physiotherapy Respiratory OR Physiotherapy OR Autogenic Drainage OR Expiratory Flow Increase Technique OR Prolonged Slow Expiratory Maneuver OR Technique of Re-educating Thoracic and Abdominal OR Chest Percussion OR Shaking OR Tapping) AND (Respiratory Mechanic OR Respiratory Mechanics OR Breathing Mechanics OR Breathing Mechanic OR Lung Volume Measurement OR Lung Volumes OR Lung Volume OR Lung Capacities OR Lung Capacity OR Sputum OR Sputums OR Induced Sputum OR Induced Sputums OR Pain OR Heart Rate OR Heart Rates OR Cardiac Rate OR Cardiac Rates OR Pulse Rate OR Pulse Rates OR Heartbeat OR Heartbeats OR Cardiac Chronotropy OR Cardiac Chronotropism OR Heart Rate Control OR Respiratory Rate OR Respiratory Rates OR Respiration Rate OR Respiration Rates OR Lung Mechanic OR Lung Mechanic OR Pulmonary Mechanic OR Pulmonary Mechanics OR Thoracoabdominal Mobility)

Total:

SCIELO

Newborn AND Physical Therapy Modalities OR Respiratory Therapy AND Respiratory Mechanics OR Lung Volume Measurements OR Heart Rate OR Respiratory Rate

Total:

COCHRANE CENTRAL

(Newborn OR Newborns OR Newborn Infant OR Newborn Infants OR Neonate OR Neonates) AND (Physical Therapy Modality OR Physical Therapy Modalities OR Physiotherapy (Techniques) OR Physiotherapies (Techniques) OR Physical Therapy

Techniques OR Physical Therapy Technique OR Group Physiotherapy OR Group Physiotherapies OR Physical Therapy OR Physical Therapies OR Physical Therapy Specialty OR Physiotherapy Specialty OR Respiratory Therapy OR Respiratory Therapies OR Chest Physical Therapy OR Conventional Chest Physical Therapy OR Chest Physiotherapy OR Physiotherapy Respiratory OR Physiotherapy OR Autogenic Drainage OR Expiratory Flow Increase Technique OR Prolonged Slow Expiratory Maneuver OR Technique of Re-educating Thoracic and Abdominal OR Chest Percussion OR Shaking OR Tapping) AND (Respiratory Mechanic OR Respiratory Mechanics OR Breathing Mechanics OR Breathing Mechanic OR Lung Volume Measurement OR Lung Volumes OR Lung Volume OR Lung Capacities OR Lung Capacity OR Sputum OR Sputums OR Induced Sputum OR Induced Sputums OR Pain OR Heart Rate OR Heart Rates OR Cardiac Rate OR Cardiac Rates OR Pulse Rate OR Pulse Rates OR Heartbeat OR Heartbeats OR Cardiac Chronotropy OR Cardiac Chronotropism OR Heart Rate Control OR Respiratory Rate OR Respiratory Rates OR Respiration Rate OR Respiration Rates OR Lung Mechanic OR Lung Mechanic OR Pulmonary Mechanic OR Pulmonary Mechanics OR Thoracoabdominal Mobility)

Total:

LILACS

Newborn AND Physical Therapy Modalities OR Physical Therapy Specialty AND Respiratory Mechanics

Total:

SCIENCE DIRECT

"Newborn Infant" AND "Physiotherapies" OR "Physical Therapy Modalities" AND "Breathing Mechanics" OR "Breathing Mechanic" OR "Lung Volume Measurement"

Total:

WEB OF SCIENCE

"Newborn Infant" AND "Physiotherapies" OR "Physical Therapy Modalities" AND "Breathing Mechanics" OR "Breathing Mechanic" OR "Lung Volume Measurement"

Total: