

# BMJ Open Efficiency of ER:YAG laser therapy in combination with behaviour management technique in reducing anxiety among paediatric dental patients – a study protocol for a randomised clinical trial

Ani Belcheva , Maria Shindova 

**To cite:** Belcheva A, Shindova M. Efficiency of ER:YAG laser therapy in combination with behaviour management technique in reducing anxiety among paediatric dental patients – a study protocol for a randomised clinical trial. *BMJ Open* 2022;**12**:e054523. doi:10.1136/bmjopen-2021-054523

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-054523>).

Received 15 June 2021  
Accepted 10 August 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Department of Paediatric Dentistry, Faculty of Dental Medicine, Medical University–Plovdiv, Plovdiv, Bulgaria

## Correspondence to

Dr Maria Shindova;  
[mariya.shindova@gmail.com](mailto:mariya.shindova@gmail.com)

## ABSTRACT

**Introduction** When providing dental care to children with a high level of dental anxiety, the range of approaches is divided into two sections: use of behaviour management techniques (BMTs) and application of alternative methods for caries removal. In an attempt to reduce dental anxiety, they can be mixed and matched in accordance with the dentists' choice. Owing to its promoted advantages, erbium-doped yttrium aluminium garnet (Er:YAG) laser turns into an ideal alternative technique for hard dental tissue therapy in anxious paediatric patients. The aim of the study is to assess the efficacy of a modified version of the BMT 'Latent inhibition' in combination with Er:YAG laser for achieving a reduction of dental anxiety in paediatric dental patients.

**Methods and analysis** This is a protocol for a randomised controlled clinical trial. The participants will be children aged 6–9 years, requiring conservative treatment of occlusal carious lesion on a second primary molar. Patients will be randomly assigned to the experimental or control group via a computer-generated sequence. In both groups, 'Latent inhibition' will be used as an anxiety-management technique. In the experimental group, caries treatment will be performed with Er:YAG laser, whereas that in the control group it will be performed with conventional rotary instruments. Outcome measures will be dental anxiety felt before and after the treatment, reported by the patient on a modified version of the Faces Scale by LeBaron et al., and the dynamics of heart rate, registered during the treatment session, which will be measured with a mobile pulse oximeter. Data will be analysed by independent sample t-test and paired t-test ( $p < 0.05$ ).

**Ethics and dissemination** The study protocol has been approved by the Committee for Scientific Research Ethics, Medical University–Plovdiv, Bulgaria (reference number P-2839, protocol of approval number 3/30.04.2015) and registered on a publicly accessible database. This research received institutional funding from the Medical University–Plovdiv, Bulgaria. The results will be presented through peer-reviewed publications and conference presentations.

**Trial registration number** NCT04924452.

## Strengths and limitations of this study

- ⇒ The study focuses on the implementation of a known behaviour management technique in the alternative caries treatment method.
- ⇒ A key strength of this study is that all participants meeting eligibility criteria will receive active treatment.
- ⇒ Both subjective and objective tools are used to assess dental anxiety in this study.
- ⇒ A limitation of this study is that it is not a split-mouth design whose advantage is the reduction of the outcome variability estimation.

## INTRODUCTION

### Background and rationale

When providing dental care to children with a high level of dental anxiety, most paediatric dentists find the conventional rotary treatment method inefficient and uncomfortable. According to the principles of behavioural dentistry, as part of paediatric postgraduate education, the so-called '4S' principle must be adapted and modified to the individual clinical situation to provide adequate dental care to anxious paediatric patients.<sup>1</sup> The range of approaches can be divided into two sections: behaviour management techniques (BMTs), on one hand, and alternative methods for caries removal, on the other hand. In an attempt to reduce dental anxiety, they can be mixed and matched in accordance with the dentists' choice.

As it has been found for more than 20 years that lasers are effective for caries excavation, laser paediatric dentistry has been rapidly developed. It offers total innovation and changes the conventional restorative treatment in pedodontics.<sup>2</sup> Owing to the

promoted advantages such as minimal intervention and prevention, safety due to the low penetration depth of the laser beam, selective removal of caries lesion, lack of thermal damage, no pain perception and use of local anaesthesia, a significant decrease of patient discomfort and dental anxiety, and increase of subjective acceptance and tolerance of laser therapy in children, erbium-doped yttrium aluminium garnet (Er:YAG) laser turns into an ideal laser for hard dental tissue therapy in anxious paediatric patients.<sup>2-4</sup>

Based on the concepts of minimal invasive dentistry, the use of BMTs during the treatment of anxious children to reduce their anxiety is required.<sup>5</sup> Several specific BMTs are not part of the regular curricula of dental students and have been used by paediatric dentists only.<sup>4,6,7</sup> Such a psychological technique is 'Latent inhibition,' also known as gradual exposure.<sup>8,9</sup> It involves a series of several positive non-painful check-ups and preventive procedures, before any invasive or painful dental manipulations. Step by step, the child is exposed to potential anxiety-provoking procedures or instruments, resulting in an acquaintance with the dental setting and personnel, as well as being accustomed to dental treatment. Despite the specific indications, required preparation and higher time consumption, the use of this technique is very rewarding as the paediatric patient eventually becomes comfortable with the dental procedure and creates a feeling of ability to cope within the child.<sup>6,7,10</sup>

Over the recent years, dentists advance in using alternative methods for caries removal as part of their everyday practice. Therefore, the investigation of this synergetic effect of laser caries removal and the different BMTs is crucial for the present and future development of paediatric dentistry and will improve the quality of dental care.

## Objectives

The aim of the study is to assess the efficacy of a modified version of the BMT 'Latent inhibition' in combination with Er:YAG laser for achieving a reduction of dental anxiety in paediatric dental patients. The main objectives are to compare dental anxiety felt during the laser and conventional dental treatment. The outcomes will be dental anxiety assessment by self-reported anxiety scale during treatment in both groups as well as the measurement of heart rate dynamics during the procedures.

## Trial design


The research is designed as a randomised parallel-group controlled clinical study. Table 1 presents the recruiting, allocation, interventions, monitoring and analysis of the research in accordance with the Standard Protocol Items: Recommendations for Interventional Trials recommendations.<sup>11</sup> In accordance with the 'Latent inhibition' technique, patients will have two visits to the dental office—a preventive procedure, the first one, and treatment of caries lesion, the second one. Two groups will be compared. In the experimental group, the enamel conditioning of the occlusal surfaces of the permanent molars before sealant application as well as the standardised caries treatment will be performed with Er:YAG laser, whereas in the control group, the conventional rotary instruments—high-speed and low-speed dental handpieces—will be used for the caries treatment.

## METHODS AND ANALYSES

### Study setting

The study setting of this research includes the Department of Paediatric Dentistry and the Laser Centre of the

**Table 1** Trial design

Timepoint*	Study period				
	Enrolment	Allocation	Post allocation		
	-t <sub>1</sub>	0	t <sub>1</sub>	t <sub>2</sub>	t <sub>3</sub>
Enrolment					
Eligibility screening	x				
Informed consent	x				
Allocation		x			
Interventions					
Experimental group (BMT+laser treatment of dental caries)				x	
Control group (BMT+caries treatment with conventional rotary instruments)				x	
Assessments					
Self-reported dental anxiety			x		x
Heart rate					
The table summarises the enrolment, allocation, interventions and assessments in the trial.					
*Postallocation time frame: t <sub>1</sub> -before the start of the treatment; t <sub>2</sub> - during laser or conventional treatment; t <sub>3</sub> - end of the treatment, before leaving the dental chair.					
BMT, behaviour management technique.					

Faculty of Dental Medicine, Medical University–Plovdiv, Bulgaria.

## Eligibility

### Inclusion criteria

- ▶ Participants in the study are children aged 6–9 years, compliant with the cognitive development of the child.
- ▶ Children requiring conservative treatment of occlusal carious lesions on a second primary molar, without spontaneous unprovoked pain, percussion or palpation pain or other symptoms, indicating pulp involvement or periodontal pathology. Lesions are classified as a distinct cavity with visible dentin without prior restoration or sealants by the International Caries Detection and Assessment System with code 05.<sup>12–14</sup> Included are caries lesions only on vital teeth.
- ▶ Children with one or more permanent molars giving indications for pit and fissure sealing.
- ▶ Patients without previous experience with laser treatment of carious lesions.
- ▶ Children who are not considered medically compromised or medically complex patients.
- ▶ Verbal assent from the child willing to comply with all study procedures and protocol.
- ▶ Obtained written informed consent by the patient's parent/guardian for participation in the study (see online supplemental file S1, 'Patient consent form', and online supplemental file S2, 'Information leaflet').

### Exclusion criteria

1. Patients who were undergoing therapy with neurological, sedative, analgesic and/or anti-inflammatory drugs 7 days prior to treatment that might affect the heart rate.
2. Children who were first-time dental patients.
3. Children with systemic diseases or physiological development delays.
4. Children with mental or cognitive problems.
5. Active infectious diseases such as influenza, scarlet fever, etc.
6. Excluded are molars which are affected by disturbances in the development of dental structures (hypoplasia, hypomineralisation and fluorosis).

### Interventions

Patients will be divided into two groups (41 per group): experimental and control groups. All treatments will be carried out by the same operator (MS), without anaesthesia. A baseline dental self-reported anxiety will be recorded using a Faces Anxiety Scale as well as the dynamics of heart rate, measured with a mobile pulse oximeter.

#### Er:YAG laser therapy protocol (experimental group)

Er:YAG laser (LiteTouch, Light Instruments), emission wavelength 2940 nm will be used for enamel conditioning of the occlusal surfaces of the permanent molars before

sealant application as well as the standardised caries treatment. Chosen protocol parameters are modified based on previously conducted studies<sup>2–4 15 16</sup>:

- ▶ HERE IT IS NECESSARY TO REMOVE THE LIST BULLET, IT STARTS FROM THE LINE BELOW - A LOW-SPEED .....Preventive procedure—sealant application.
- ▶ A low-speed rubber cup and pumice paste (Clean-Polish, Kerr) will be used for 30s for cleaning and polishing of the occlusal surface of the chosen permanent molar.
- ▶ Tooth surface will be washed for debris and organic residue removal and dried with air spray.
- ▶ Isolation with rubber dam.
- ▶ Laser conditioning of the occlusal enamel surface. The parameter settings used will be tip-to-tissue distance, 1.5 mm from the tooth surface; tip diameter, 600 µm; laser energy, 70 mJ; pulse frequency, 10 Hz; water spray level, 8; average power, 0.7 W; and energy density, 67 J/cm<sup>2</sup>.
- ▶ The tooth surface will be etched with 35% phosphoric acid gel (Etching Gel, DMP) for 30s and rinsed for the same time.
- ▶ The tooth surface will be dried with air spray for 15s.
- ▶ Fissure sealant application (Pit&Fissure Sealant, DMP).
- ▶ Light cured for 20s.
  - HERE IT IS NECESSARY TO REMOVE THE LIST BULLET, Caries removal—parameters: enamel removal—energy, 100–200 mJ; density 9.84–13.03 J/cm<sup>2</sup>; pulse frequency, 20 Hz; tip diameter, 800 µm; water spray level, 8; tip-to-tissue distance, 0.5÷1.0 mm from the tooth surface; dentin removal—energy, 100 mJ; density, 9.84 J/cm<sup>2</sup>, pulse frequency, 20 Hz; tip diameter, 800 µm; water spray level, 8; tip-to-tissue distance, 0.5÷1.0 mm from the tooth surface; restoration with compomer; time for caries removal procedure—max 8 min.<sup>17–20</sup>

#### Conventional therapy protocol (control group)

- ▶ HERE IT IS NECESSARY TO REMOVE THE LIST BULLET, IT STARTS FROM THE LINE BELOW - A LOW-SPEED .....Preventive procedure—sealant application.
- ▶ A low-speed rubber cup and pumice paste (Clean-Polish, Kerr) will be used for 30s for cleaning and polishing of the occlusal surface of the chosen permanent molar.
- ▶ The tooth surface will be washed for debris and organic residue removal and dried with air spray.
- ▶ Isolation with rubber dam.
- ▶ The tooth surface will be etched with 35% phosphoric acid gel (Etching Gel, DMP) for 30s and rinsed for the same time.
- ▶ The tooth surface will be dried with air spray for 15s.
- ▶ Fissure sealant application (Pit&Fissure Sealant, DMP).
- ▶ Light cured for 20s.

- HERE IT IS NECESSARY TO REMOVE THE LIST BULLET. Caries removal—conventional rotary instruments will be used: high-speed and low-speed dental handpieces. For the bur preparation, 1.2 mm diameter diamond round bur Drendel & Zweiling No. 801.314 and Komet Steel round bur 016, Komet Dental Gebr were used. A new bur was used for each preparation. Restoration was performed with compomer. Time for caries removal procedure was a maximum of 4 min.

## Clinical protocol

### First visit

1. Parents/guardians are informed about the protocol of the study and the laser technique. They sign the informed consent form (see online supplemental files S1 and S2). Verbal assent from the child is obtained.
2. Oral examination and sealant application are performed according to the assigned intervention.
3. Patient's self-report of dental anxiety before leaving the dental chair.

### Second visit

1. Patients will be asked to report their dental anxiety, pointing to the face or choose the number which most closely depicted its state of anxiety using a modified version of the self-report Faces Scale by LeBaron<sup>21</sup> (see online supplemental file S3).
2. The pulse oximeter is connected to the patient's index finger. The start of heart rate monitoring and recording will be 5 min prior to treatment. Time frame: at least 5 min after the dental treatment, before leaving the dental chair.
3. Caries treatment is performed according to the assigned intervention.
4. Patient's self-report of dental anxiety before leaving the dental chair.

## Outcomes

### Primary outcome measures

The primary outcome will be the dental anxiety before and after the treatment session, reported by the patient on a modified version of the self-report Faces Scale by LeBaron et al. The scale comprises a row of five faces ranging from 'relaxed' to 'very worried' in combination with a visual analogue scale of 0–10. Each child was asked to point to the face or choose the number which most closely depicted its state of anxiety.

### Secondary outcome measures

The secondary outcome will be the dynamics of heart rate, registered during the treatment session measured with a mobile pulse oximeter (CMS50F, CONTEC), placed on the index finger of the left hand.<sup>22</sup> Throughout the whole procedure of each dental visit, data were recorded and analysed by a specially developed digital processing and graphic visualisation software SPO2 Review V.1.2 rel.

## Participant's timeline

Each eligible patient undergoes two visits. The first appointment includes screening, consenting and assenting, recording of dental anxiety and sealant application according to the assigned interventions for each group. The second appointment at the 1-week recall includes a recording of dental anxiety and treatment of a carious lesion according to the assigned interventions for each group. The manipulations will be performed by one operator.

## Sample size calculation

The sample size calculation is performed based on data from a pilot study with 20 subjects. To estimate sample size for the primary outcome—self-reported anxiety felt, according to the Faces Scale by LeBaron—a t-test for paired groups has been used (G\* Power software V.3.1,6 since we have two groups). The effect size was determined using the formula

$$ES = \frac{\text{Control} - \text{Treated}}{SD_{\text{pooled}}} = \frac{2.33 - 0.33}{3.25} = 0.62,$$

where SD is the pooled SD, an average of the SD of the experimental and control groups. The sample size is calculated to assure a test power greater than 95% and a significant level of  $\alpha=0.05$ . We estimated a sample size of 41 patients per group to detect significant differences. Thus, the final sample size for this study will be 82 patients.

## Recruitment

The patients at the Department of Paediatric Dentistry of the Faculty of Dental Medicine, Medical University–Plovdiv, Bulgaria, who meet the inclusion criteria, will be screened for eligibility. Once identified, patients will be informed about this research project and will receive information about the possibility of potential study participation. Patient recruitment starts obtaining the full quota of participants within a 1-year time frame. It begins in September 2021 with an estimated enrolment capacity of five patients per month.

## Participating centres

The patients are randomly selected from the visitors in the Department of Paediatric Dentistry of the Faculty of Dental Medicine, Medical University–Plovdiv, Bulgaria, and are treated in the Laser Centre of the same university.

## Assignment of the intervention

### Sequence generation

The patients will be randomly allocated to either the control group or the experimental group (41 patients in each group) according to the enrolment number in the trial. The randomisation will be created using a computerised random generator.

## Allocation concealment mechanism and implementation

A randomisation list will be created by a random generator before the start of the treatment and kept in a locked drawer. Assignments will be kept in separate, closed opaque, sequentially numbered envelopes, enabling



the sequence to be concealed until the intervention is assigned.

### Blinding

The randomisation will be independent, that is, the patients and parents/guardians will remain blinded to group status. The operator will get acquainted with the procedure to be performed prior to the first session. The operator is selected to be the only one performing the manipulation to prevent bias. The statistician will be blinded to treatment assignment as data will be masked before the analysis without giving the statistician the key.

### Data collection, confidentiality, storage and monitoring of the study documents

Collection, coding, storage and evaluation of personal data within the project will be carried out in accordance with The General Data Protection Regulation (EU) 2016/679. A prerequisite for data collection will be the voluntary written informed consent of the patient's parent or guardian. Confidentiality will be guaranteed by a coded ID number; access will be granted exclusively to the study investigators. The information from the paper forms will be exported to a database file and stored on a password-protected computer. Only the investigators and statistician will have access to the final data set. All data collected will be stored in sealed containers in areas of the Department of Paediatric Dentistry, Faculty of Dental Medicine, Medical University–Plovdiv, Bulgaria, with limited access.

## STATISTICAL METHODS

The obtained data will be recorded, tabulated, processed, and analysed using SPSS (Statistical Package for Social Science software) V.21.0 (IBM, USA). In all tests, the significance level of 5% probability or the corresponding *P*-value will be adopted. Descriptive statistics will be calculated. Discrete variables will be summarised by frequencies or proportions. Continuous variables will be presented as means and SD. We will compare anxiety mean scores according to the Faces Scale by LeBaron as well as heart rate mean score. Comparisons among groups will be performed by using the Independent sample t-test and paired t-test.

### Patient and public involvement

The development of the research question and outcome measures will be based on the review of available evidence in this research area. Patients will not be involved in the development of the study protocol. However, their questions and concerns will be addressed during patient recruitment and study implementation. During the conduction of the study, patients will not be informed about the results of the ongoing trial since there is no planned interim analysis. The results will be disseminated to the study participants through email and routine follow-up dental check-ups.

### Ethics and dissemination

The clinical study will be conducted in accordance with the conditions and principles of the Declaration of Helsinki, the existing EU Clinical Trial Directive (EC) No. 2001/20/EC, the recommendations of the Ethical Committee at the Medical University–Plovdiv, Bulgaria, and the international ethical and scientific quality standard for designing, recording and reporting trials that involve the participation of human subjects - *Good Clinical Practices (GCP)*.

### Research ethics approval

The study was approved by the Committee for Scientific Research Ethics, Medical University–Plovdiv, Bulgaria (reference number P-2839, protocol of approval number 3/30.04.2015) and registered on a publicly accessible database (ClinicalTrials.gov, Registration number:NCT04924452). Ethical approval for the study protocol was provided by and written informed consent for all subjects' parents/guardians was approved by the ethics committee of the Medical University–Plovdiv, Bulgaria.

### Consent

The operators will obtain written consent from patients' parents/guardians willing to participate in the trial. Additional information will be provided for all parents for the study. Completed informed consent will be collected at the Department of Paediatric Dentistry, Medical University–Plovdiv by the study investigators. A copy of the signed consent form will be handed over to the participating child's parent/guardian. After providing age-appropriate information about the study, verbal assent will be obtained as an affirmative agreement for participation from children.

### Confidentiality

Information on the participants collected during the study will be kept strictly confidential and will not be disclosed to third parties. Confidentiality will be guaranteed by a coded ID number; access will be granted exclusively to the study investigators.

### Conflict of interests

The investigators have no conflicts of interest to declare. They agree with the protocol and the informed consent of the study, and there is no financial interest to report.

### Access to data

All data collected will be stored in sealed containers in areas of the Department of Paediatric Dentistry, Faculty of Dental Medicine, Medical University–Plovdiv, Bulgaria, with limited access. The information from the paper forms will be exported to a database file and stored on a password-protected computer. Only the investigators and statistician will have access to the final data set.

### Dissemination policy

The results of the trial will be presented through peer-reviewed publications and conference presentations. In

addition, our results will be disseminated to clinicians, as well as key stakeholders, including scientific directors of postgraduate programmes 'Master of Science in Lasers in Dentistry', academic courses in Pedodontics and Preventive dentistry. The principal investigator (MS) and the scientific expert (AB) will write the first draft of the manuscript without the use of professional writers.

### Trial status

The trial is not yet recruiting patients. The process will start in September 2021 and will continue until September 2022.

**Acknowledgements** The authors show their gratitude to Associate Professor Georgi Tomov, PhD, for the fruitful discussion. Assistance with the Laser Centre of the Faculty of Dental Medicine, Medical University–Plovdiv, Bulgaria, was greatly appreciated. The authors are also thankful to Professor Nonka Mateva, PhD, for the statistical consultancy expertise in the planning of this clinical trial.

**Contributors** AB and MS conceived the ideas; MS was the primary operator, AB and MS were outcomes assessors and data collectors; and developed the design and protocol for this study. Writing was led by AB and MS. All authors made substantial contributions, participated in the analysis and reporting of the results, refined the study protocol and approved the final manuscript.

**Funding** This research received institutional funding from the Medical University–Plovdiv, Bulgaria.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, conduct, reporting or dissemination plans of this research.

**Patient consent for publication** Consent obtained from parent(s)/guardian(s).

**Ethics approval** This study involves human participants and was approved by The study was approved by the Committee for Scientific Research Ethics, Medical University–Plovdiv, Bulgaria (reference number P-2839, protocol of approval number 3/30.04.2015) and registered on a publicly accessible database (ClinicalTrials.gov). Ethical approval for the study protocol was granted by and written informed consent from all subjects' parents/guardians was approved by the Ethics Committee of the Medical University–Plovdiv, Bulgaria. Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All data relevant to the study are included in the article or uploaded as supplementary information.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### ORCID iDs

Ani Belcheva <http://orcid.org/0000-0002-9625-8684>

Maria Shindova <http://orcid.org/0000-0003-2996-3700>

### REFERENCES

- Walsh LJ. Anxiety prevention: implementing the 4s principle in conservative dentistry. *Auxiliary* 2007;17:24–6.
- Pagano S, Lombardo G, Orso M, *et al.* Lasers to prevent dental caries: a systematic review. *BMJ Open* 2020;10:e038638.
- Galui S, Pal S, Mahata S, *et al.* Laser and its use in pediatric dentistry: a review of literature and a recent update. *Int J Pediatr Rehabil* 2019;4:1–5.
- American Academy of Pediatric Dentistry. *Policy on the use of lasers for pediatric dental patients. The reference manual of pediatric dentistry*. Chicago, Ill: American Academy of Pediatric Dentistry, 2020: 116–8.
- Frencken JE, Peters MC, Manton DJ, *et al.* Minimal intervention dentistry for managing dental caries - a review: report of a FDI task group. *Int Dent J* 2012;62:223–43.
- Buldur B. Behavior management in pediatric dentistry: an overview and interpretation. *Pesqui. bras. odontopediatria clin. integr.* 2019;19:e4649:1–2.
- Khandelwal M, Shetty RM, Rath S. Effectiveness of distraction techniques in managing pediatric dental patients. *Int J Clin Pediatr Dent* 2019;12:18–24.
- Gunasekaran G, Ramakrishnan M. Evaluation of factors affecting the behaviour of uncooperative pedodontic patients. *PJAE* 2020;17:2027–38.
- Subramaniam P, Haq M, Gupta M, *et al.* Assessment of trait and state anxiety in 3–6-year old children during sequential phases of dental treatment. *Contemp Pediatr Dent* 2020;1:22–32.
- Tajadura-Jiménez A, Grehl S, Tsakiris M. The other in me: interpersonal multisensory stimulation changes the mental representation of the self. *PLoS One* 2012;7:e40682.
- Chan A-W, Tetzlaff JM, Altman DG, *et al.* SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med* 2013;158:200–7.
- Ismail AI, Sohn W, Tellez M, *et al.* The International caries detection and assessment system (ICDAS): an integrated system for measuring dental caries. *Community Dent Oral Epidemiol* 2007;35:170–8.
- International Caries Detection and Assessment System (ICDAS) Coordinating Committee. Rationale and evidence for the International caries detection and assessment system (ICDAS II), 2011. Available: <http://www.icdas.org>
- Pitts N, "Pitts N. "ICDAS"--an international system for caries detection and assessment being developed to facilitate caries epidemiology, research and appropriate clinical management. *Community Dent Health* 2004;21:193–8.
- Yilmaz H, Keles S. The effect of the ER: YAG laser on the clinical success of hydrophilic fissure sealant: a randomized clinical trial. *Eur Oral Res* 2020;54:148–53.
- Rattanacharoenthum A, Na-Lamphun P, Kantrong N. Altered adhesion of dental sealant to tooth enamel microscopically modified by Er:YAG laser irradiation: An *in vitro* study. *Laser Ther* 2019;28:19–25.
- Sarmadi R, Andersson EV, Lingström P, *et al.* A Randomized Controlled Trial Comparing Er:YAG Laser and Rotary Bur in the Excavation of Caries - Patients' Experiences and the Quality of Composite Restoration. *Open Dent J* 2018;12:443–54.
- Li T, Zhang X, Shi H, *et al.* Er:YAG laser application in caries removal and cavity preparation in children: a meta-analysis. *Lasers Med Sci* 2019;34:273–280.
- Johar S, Goswami M, Kumar G, *et al.* Caries removal by Er,Cr:YSGG laser and Air-rotor handpiece comparison in primary teeth treatment: an *in vivo* study. *Laser Ther* 2019;28:116–22.
- Montedori A, Abrahá I, Orso M, *et al.* Lasers for caries removal in deciduous and permanent teeth. *Cochrane Database Syst Rev* 2016;9:CD010229.
- LeBaron S, Zeltzer L. Assessment of acute pain and anxiety in children and adolescents by self-reports, observer reports, and a behavior checklist. *J Consult Clin Psychol* 1984;52:729–38.
- Song J-S, Chung H-C, Sohn S, *et al.* Effects of psychological behaviour management programme on dental fear and anxiety in children: a randomised controlled clinical trial. *Eur J Paediatr Dent* 2020;21:287–91.



## Consent form

For a patient's consent to publication of images and/or information about them in BMJ publications.

Name of patient: \_\_\_\_\_

Patient's parent/guardian: \_\_\_\_\_

Material – text (information about your child's oral health, level of dental anxiety and dynamics of heart rate): \_\_\_\_\_

Provisional title of article in which Material will be included: Efficacy of Combined Er:YAG Laser Therapy and Behaviour Management Technique in Reducing Anxiety among Paediatric Dental Patients

---

### CONSENT

I \_\_\_\_\_ [ENTER YOUR FULL NAME] give my consent for the Material about my child to appear in a BMJ publication.

**I confirm that I:** (please tick boxes to confirm)

- ☐ **have seen the text or other material about my child**
- ☐ **have read the article to be submitted to BMJ**
- ☐ **am legally entitled to give this consent.**

I understand the following:

- (1) The Material will be published without my child's name attached, however I understand that complete anonymity cannot be guaranteed. It is possible that somebody somewhere may recognise my child.
- (2) The Material may show or include details of my child's medical condition or injury and any prognosis, treatment or surgery that I have/the patient has, had or may have in the future.
- (3) The article may be published in a journal which is distributed worldwide. BMJ's publications go mainly to doctors and other healthcare professionals but are also seen by many others including academics, students and journalists.
- (4) The article, including the Material, may be the subject of a press release, and may be linked to from social media and/or used in other promotional activities. Once published, the article will be placed on a BMJ website and may also be available on other websites.
- (5) The text of the article will be edited for style, grammar and consistency before publication.
- (6) I and my child will not receive any financial benefit from publication of the article.

Patient consent form 050419



- (7) The article may also be used in full or in part in other publications and products published by BMJ and/or by other publishers. This includes publication in English and in translation, in print, in digital formats, and in any other formats that may be used by BMJ or other publishers now and in the future. The article may appear in local editions of journals or other publications, published in the UK and overseas.
- (8) I can revoke my consent at any time before publication, but once the article has been committed to publication ("gone to press") it will not be possible to revoke the consent.
- (9) This consent form will be retained securely and in confidence by BMJ in accordance with the law, for no longer than necessary. Personal data provided in this form will be used and retained in accordance with BMJ's Privacy Policy available at <https://www.bmj.com/company/your-privacy/>.

Signed: \_\_\_\_\_ Print name: \_\_\_\_\_

Address: \_\_\_\_\_ Email address: \_\_\_\_\_

\_\_\_\_\_ Telephone no: \_\_\_\_\_

\* signing on behalf of the patient who is under the age of 18

\_\_\_\_\_ Date: \_\_\_\_\_

#### Corresponding author

Signed: \_\_\_\_\_

Author's name: Maria Shindova

Position: Senior Assistant Professor

Address: 3 Hristo Botev Bulv., Plovdiv, Bulgaria

Institution: Department of Pediatric

Dentistry, Faculty of Dental Medicine,

Medical University of Plovdiv

Email address:  
mariya.shindova@gmail.com

Telephone no: + 359 898 390 935

Date: \_\_\_\_\_



## INFORMATION LEAFLET

### DEPARTMENT OF PAEDIATRIC DENTISTRY LASER CENTER FACULTY OF DENTAL MEDICINE, MEDICAL UNIVERSITY OF PLOVDIV, BULGARIA

#### EFFICIENCY OF ER:YAG LASER THERAPY IN COMBINATION WITH BEHAVIOUR MANAGEMENT TECHNIQUE IN REDUCING ANXIETY AMONG PAEDIATRIC DENTAL PATIENTS

**DESCRIPTION:** You and your child are invited to participate in a research study on the efficiency of Er:YAG laser therapy in combination with behavior management technique in reducing anxiety among pediatric dental patients.

**PROCEDURES:** With your permission, we would like to collect information about your children's dental anxiety before, during and after dental treatment of a caries lesion. This study does not involve any experiments, just preventive procedures and dental treatment, collection, and study of the required information.

**RISKS AND BENEFITS:** There are no anticipated risks associated with this study. You will not receive any direct benefit from participation.

**TIME INVOLVEMENT:** Your child's participation in this study will not require more time from you other than for the first visit including an explanation of the study, oral examination and a preventive procedure (sealant application). The second appointment at 7-day recall will include dental treatment.

**PAYMENTS:** You will not be paid to participate in this study. You will not pay for the treatment of your child in this study.

**PARTICIPANT'S RIGHTS:** If you have read this form and have decided your child to participate in this research, please understand your participation is voluntary and you have the right to withdraw your consent or discontinue participation at any time without penalty or loss of benefits to which you are otherwise entitled.

The results of this research study may be presented at scientific or professional meetings or published in scientific journals. However, your identity will not be disclosed.

Thank you for your time and attention!

Name of parent/guardian .....

Signature of parent /guardian .....

Name of patient .....

Telephone number .....

Name of the dentist .....

Signature of the dentist .....

For additional information regarding the trial, you can contact us at the given address, emails, or phone numbers.

**Researchers:**

Maria Shindova, DDS, MSc, PhD  
Chief Assistant Professor  
Department of Pediatric Dentistry  
Faculty of Dental Medicine  
Medical University – Plovdiv  
3 Hristo Botev Blvd  
4000 Plovdiv, Bulgaria  
Mobile: + 359 898 390 935  
mariya.shindova@gmail.com  
Mariya.Shindova@mu-plovdiv.bg

Ani Belcheva, DDS, MSc, PhD  
Professor  
Department of Pediatric Dentistry  
Faculty of Dental Medicine  
Medical University - Plovdiv  
3 Hristo Botev Blvd  
4000 Plovdiv, Bulgaria  
Mobile: + 359 889 528 932  
abeltcheva@yahoo.com

