


BMJ Open Acupuncture versus massage for pain in patients living with advanced cancer: a protocol for the IMPACT randomised clinical trial

Sally A D Romero,¹ Nicholas Emard,² Raymond E Baser,³ Katherine Panageas,² Jodi MacLeod,⁴ Desiree Walker,⁴ Margaret Barton-Burke,⁵ Kevin Liou,⁴ Gary Deng,⁴ John Farrar,⁶ Han Xiao,⁷ Jun J Mao ⁴, Andrew Epstein⁷

To cite: Romero SAD, Emard N, Baser RE, *et al*. Acupuncture versus massage for pain in patients living with advanced cancer: a protocol for the IMPACT randomised clinical trial. *BMJ Open* 2022;**12**:e058281. doi:10.1136/bmjopen-2021-058281

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-058281>).

Received 14 October 2021
Accepted 06 June 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Jun J Mao; maoj@mskcc.org

ABSTRACT

Introduction Pain, comorbid fatigue and sleep disturbances are common and distressing symptoms for patients with advanced cancer, negatively impacting their quality of life. Clinical guidelines recommend non-pharmacological interventions, including acupuncture and massage, for pain management in adult patients with cancer in adjunct to conventional care. However, high-quality evidence about the comparative effectiveness and long-term durability of these therapies for symptom management is limited.

Methods and analysis We describe the design of a two-arm, parallel group, multicentre randomised controlled trial that investigates the use of acupuncture versus massage for musculoskeletal pain among 300 patients with diverse types of advanced cancer. The primary aim is to evaluate the long-term effectiveness (26 weeks from randomisation) of acupuncture vs massage for pain (primary outcome) and comorbid symptoms (fatigue, sleep disturbance and quality of life). The secondary aim is to identify patient-level demographic characteristics (eg, sex, race, age), clinical factors (eg, insomnia, pain severity) and psychological attributes that are associated with a greater reduction in pain for either acupuncture or massage. Patients will receive weekly acupuncture or massage treatments for 10 weeks, followed by monthly booster sessions up to 26 weeks. The primary endpoint will be the change in worst pain intensity score from baseline to 26 weeks. We will collect validated patient-reported outcomes at multiple time points over 26 weeks.

Ethics and dissemination The Institutional Review Board at Memorial Sloan Kettering Cancer Center in New York approved this protocol. Results will be disseminated via peer-reviewed scientific journals and conference presentations. Our findings will help patients and healthcare providers make informed decisions about incorporating non-pharmacological treatments to manage pain for patients with advanced cancer.

Trial registration number NCT04095234.

INTRODUCTION

Cancer is a leading cause of morbidity and mortality, second only to heart disease.¹ Because of recent innovations in cancer

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This study represents the largest randomised controlled trial to date comparing the effectiveness of acupuncture versus massage for pain management among patients with advanced cancer.
- ⇒ By recruiting a diverse population in terms of race/ethnicity and cancer types, this study will offer insight into the sociodemographic, clinical factors and physiological attributes that can inform and help predict factors to personalise treatment.
- ⇒ All participants will be followed up to 26 weeks.
- ⇒ The study design does not include a control group comparing the standard of care for pain management as prescribed by the clinical team.
- ⇒ The study design does not allow crossover between the acupuncture and massage groups.

therapeutics, the definition for advanced cancer is challenging because some patients with metastatic cancer can now be 'cured' or at least enter long-term remission leaving them to often live with symptomatic sequelae. Compared with the general population, patients with advanced cancer are at a greater risk for chronic physical and psychological symptoms.^{2–5} Among patients with advanced cancer, symptoms of pain, fatigue and insomnia are the most commonly reported, often clustered together, and are generally not well managed.^{4–10} Previous studies have shown prevalence rates of pain as high as 66% among patients with advanced cancer.^{11 12}

Historically, pain management in cancer has predominantly relied on drug therapies; however, increasing clinical evidence suggesting the potential harm over time of long-term opioid therapy for chronic cancer pain, not to mention the current opioid abuse epidemic sweeping the USA, underscore a need for additional treatments.^{13 14} As more individuals with advanced cancer

live longer, patient-centred pain management integrating non-pharmacological interventions based on research evidence has strong potential to improve the quality of pain management for this population. Hence, clinical guidelines and leading medical organisations recommend non-pharmacological interventions, including acupuncture and massage, in conjunction with drug therapies for pain management.^{14–19}

Acupuncture, a therapy of traditional Chinese medicine, involves penetrating the skin with thin, solid, metallic needles that are manipulated by hand or electrical stimulation.²⁰ With respect to the efficacy of acupuncture for chronic pain in cancer populations, a systematic review and meta-analysis found that when acupuncture is incorporated into conventional cancer care, it is more effective than conventional drug management alone for cancer pain.²¹ A recent comparative effectiveness randomised controlled trial (RCT) found that electroacupuncture and auricular acupuncture were significantly more efficacious for pain reduction than usual care among diverse cancer survivors (N=360).²² Further, there is some evidence suggesting that acupuncture may improve sleep disturbances, fatigue and anxiety in patients with cancer experiencing pain.^{23 24}

Massage, which involves the manual manipulation of muscles and other soft tissue areas of the body, is one of the earliest known forms of pain relief. Since massage therapy techniques promote joint flexibility, relieve muscular tension and improve range of motion, massage therapy has mechanistic plausibility for addressing musculoskeletal pain in patient populations.^{13 25} In a recent meta-analysis conducted by the Evidence for Massage Therapy Working Group, massage therapy was effective at treating pain compared with other controls (such as reading, usual care or active attention) in cancer populations.²⁶ In addition to pain management, massage therapy may improve fatigue, sleep and anxiety in cancer populations.^{26–30}

Despite acupuncture and massage therapy both being widely available and commonly used as non-pharmacological treatments for pain,^{13 31} there is currently a gap in the evidence regarding the comparative effectiveness of these options as well as the long-term durability of their treatment effects among patients living with advanced cancer. We planned an RCT to evaluate the long-term comparative effectiveness of acupuncture vs massage for pain in patients living with advanced cancer. Our primary aim is to compare the long-term effectiveness (26 weeks from randomisation) of acupuncture versus massage for pain (primary outcome) and comorbid symptoms (fatigue, sleep disturbance and quality of life (QoL)) in patients living with advanced cancer. Our secondary aim is to identify patient-level demographic characteristics (eg, sex, race, age), clinical factors (eg, insomnia, pain severity) and psychological attributes (ie, outcome expectation) that are associated with a greater reduction in pain for either acupuncture or massage.

METHODS AND ANALYSIS

Study design

The Integrative Medicine for Pain in Patients with Advanced Cancer Trial (IMPACT) is a two-arm, parallel group RCT to compare the effectiveness of acupuncture and massage for pain and comorbid symptoms in a heterogeneous sample of 300 patients living with advanced cancer who have been experiencing moderate to severe pain (figure 1). Eligible patients will be randomly assigned in a 1:1 ratio to acupuncture or massage. Patients will receive weekly acupuncture or massage treatments for 10 weeks followed by monthly booster sessions up to week 26. All patients will continue to receive their standard medical care and pain management as prescribed by their physicians. The primary endpoint will be the change in worst pain intensity score (as assessed by the Brief Pain Inventory (BPI)) from baseline to 26 weeks. We will also collect validated patient-reported outcome measures of pain and comorbid symptoms at seven timepoints over 26 weeks (table 1).

Participants

We will recruit study participants in the USA through Memorial Sloan Kettering Cancer Center (MSK), a National Cancer Institute-designated comprehensive cancer centre, with a main campus in Manhattan and numerous regional sites in New York (Westchester County and Long Island) and New Jersey (Bergen, Monmouth and Basking Ridge). We will also recruit patients from the Baptist Health Miami Cancer Institute (MCI), which is an affiliate of MSK's strategic alliance. For MSK-affiliated patients, we will use a population-based method by mailing out letters to potentially eligible patients identified through a data query of MSK's electronic health records. We will also use stakeholders and partnering clinicians to publicise the study and provide referrals. The target accrual goal is 300 participants. Enrolment began in October 2019 and study participant assessments are scheduled to be completed by July 2022.

Inclusion and exclusion criteria

Patients will be eligible for the study if they are English or Spanish-speaking, over 18 years old, and able to walk with only occasional assistance (Karnofsky functional score of ≥ 60). They must also have a diagnosis of the following: stage III or IV lung cancer; any stage pancreatic cancer; unresectable cholangiocarcinoma; unresectable liver cancer; unresectable ampullary or periampullary cancer or other stage IV gastrointestinal cancer; stage III or IV ovarian or fallopian tube cancers or other stage IV gynecologic cancer; stage IV breast cancer; stage IV genitourinary cancer; stage III or IV sarcoma; stage IV melanoma; stage III or IV head/neck cancer; stage IV endocrine cancer; or haematological malignancies (lymphoma, myeloma and leukaemia). Patients will need to have an expected prognosis of 6 months or greater from their treating physician or the study clinician.




Figure 1 Study schema for the Integrative Medicine for Pain in Patients with Advanced Cancer Trial. PROMIS, Patient-Reported Outcomes Measurement Information System.

To be eligible, patients must also report ongoing musculoskeletal pain, defined as regional (eg, joints, extremities, back, neck) or more generalised (ie, fibromyalgia) pain, as their primary source of pain. The pain must be present for at least 1 month and occur for at least 15 days of the preceding 30 days. In addition, patients must report that their pain is four or greater on a numerical rating scale of 0–10. Non-musculoskeletal pain syndromes

(eg, headache, facial pain, chest pain or visceral abdominal pain) may be present if musculoskeletal pain is the primary source of pain. Patients will be excluded from the study if they have a blood platelet count of less than 15 000 platelets per microliter.

Table 1 Schedule of data collection

Outcome	Active intervention			Follow-up			
	Week 0	Week 4	Week 10	Week 14	Week 18	Week 22	Week 26
Primary outcome—pain							
Brief Pain Inventory	X	X	X	X	X	X	X
Secondary outcomes—fatigue, sleep, anxiety and quality of life							
Brief Fatigue Inventory	X		X		X		X
Insomnia Severity Index	X		X		X		X
Hospital Anxiety and Depression Scale	X		X		X		X
PROMIS-10 Global Health	X		X		X		X
Patients' Global Impression of Change			X		X		X
Covariates							
Demographics (eg, age, sex, race/ethnicity)	X						X
Clinical characteristics (eg, tumour type, stage, cancer therapy)	X						X
Pain medication diary	X	X	X	X			X
Predictive variables							
Mao expectancy of treatment effects	X		X				
PROMIS-10, Patient-Reported Outcomes Measurement Information System.							

Procedure

All potential participants will undergo an initial screening with a research coordinator in person or over the telephone. At this initial contact, the research coordinator will explain the study goals and procedures and screen participants for eligibility. Next, a study healthcare provider will meet with screened and interested patients to confirm eligibility. Once deemed eligible, patients will complete the informed consent and undergo randomisation. Patients will complete assessments online using Research Electronic Data Capture (REDCap), a data management software system, at seven time points: weeks 0, 4, 10, 14, 18, 22 and 26. To encourage adherence to the study procedures, participants will receive reminders to complete study assessments. Additionally, all participants will be compensated with a US\$40 gift card for completion of the week 10 visit and a US\$60 gift card for the completion of the week 26 visit, for a total of US\$100.

Randomisation

We will randomise 300 participants to acupuncture or massage using MSK's Clinical Research Database (CRDB), a secure computer system that ensures full allocation concealment. Randomisation will be performed by the method of random permuted block stratified by any current opioid use (yes vs no) and by accrual site (MSK main campus, MSK regional sites vs MCI). Given the nature of the interventions, patients and providers will not be blinded to treatment assignments. The principal investigator (PI), study statisticians and outcome assessment clinical research coordinator will be blinded to treatment assignments.

Primary outcome

The short-form BPI is one of the most widely used instruments to measure pain and has been demonstrated to be a reliable, valid and responsive measure (Cronbach's alpha 0.77–0.91).³² The BPI contains four pain severity items and seven pain interference items, all rated on a scale from 0 to 10 (higher ratings indicate worse pain intensity/interference). A pain interference subscale can be computed by taking the average rating of the seven pain interference items. A pain severity subscale score can similarly be computed for the four pain severity items; however, the Worst Pain severity item and the Average Pain severity item are often examined separately from the pain intensity subscale in clinical research because they tend to be more sensitive indicators of changes in patients' perceived pain. The primary outcome of this study will be the patient's rating of their Worst Pain in the past week with response choices of 0 'no pain' to 10 'pain as bad as you can imagine.' The Average Pain rating in the past week and the pain interference subscale will be used as secondary pain outcomes.

Secondary outcomes

The Patients' Global Impression of Change (PGIC) is a one-item survey used to define a clinically important

change in pain from the patient's perspective.^{33 34} The PGIC can be used as an anchor to derive anchor-based minimally important differences for pain measures like the BPI. Participants will be asked 'How would you describe your pain since the first clinical visit? I am: very much worse, much worse, a little worse, the same, a little improved, much improved, very much improved.' Subjects reporting 'much improved' or 'very much improved' will be considered responders.

The Brief Fatigue Inventory is a nine-item instrument designed to assess fatigue severity and has been shown to be reliable and valid in multiple languages and diverse populations.^{35 36} Three items ask patients to rate the severity of their fatigue at its 'worst,' 'usual,' and 'now' during normal waking hours, with 0 being 'no fatigue' and 10 being 'fatigue as bad as you can imagine.' Six items ask patients to rate the amount that fatigue has interfered with different aspects of their life during the past 24 hours, with 0 being 'does not interfere' and 10 being 'completely interferes.'³⁵ A composite fatigue severity score can be found by averaging the nine-item scores.

The Insomnia Severity Index is a reliable and valid seven-item scale used to assess subjective insomnia severity.^{37 38} The items are scored on a five-point Likert response scale (eg, 0=no problem; 4=very severe problem), yielding a total score ranging from 0 to 28 with higher scores representing more severe insomnia symptoms. Established cutoffs are: <8, no clinically significant insomnia; 8–14, subthreshold insomnia; 15–21, clinical insomnia (moderate severity); >21, clinical insomnia (severe).³⁷ A reduction of eight points is considered to be clinically meaningful improvement among those with insomnia.³⁹

The Hospital Anxiety and Depression Scale is a 14-item scale with seven items measuring depression and seven items measuring anxiety that has been shown to be both reliable and valid.^{40 41} Each item is answered by the patient on a four-point (0–3) response category so possible scores range from 0 to 21 for anxiety and depression, with higher scores indicating higher symptomatology. Established cutoffs are: 0–7 not significant; 8–10 subclinical, and 11–21 clinically significant depression/anxiety.⁴²

The Patient-Reported Outcomes Measurement Information System Scale V.1.2—Global Health is a brief instrument composed of 10 items that demonstrates adequate reliability and validity^{43 44} as a measure of health-related QOL in general and clinical populations.^{45 46} The measure yields two scores for physical health and mental health with higher scores indicative of better QOL.

Assessment of outcome expectancy as a predictive variable for treatment response

Outcome expectancy has long been considered an important predictor of treatment outcomes and has gained increasing recognition in massage and acupuncture clinical trials.^{47 48} The Mao Expectancy of Treatment Effects,⁴⁹ originally developed as the Acupuncture Expectancy Scale,⁵⁰ is a four-item instrument to measure outcome expectancy for various interventions (eg,

acupuncture, herbs, cognitive behavioural therapies)^{49 51} and has demonstrated reliability and validity.⁵⁰ The score ranges from 4 to 20, with a higher score indicating greater expectancy. We will use this measure to explore whether expectancy predicts treatment outcomes and may impact the observed differences between groups.

Covariates

We will collect specific demographic (eg, age, sex, race/ethnicity) and other relevant historical medical data (eg, cancer treatment). We will also track participants' self-reported use of analgesic medications (eg, acetaminophen, non-steroidal anti-inflammatory drugs, opioids and adjuvants for neuropathic pain) by having participants complete weekly pain medication diaries to calculate weekly average analgesic medication usage throughout the study time period.⁵² As pain often results in increased healthcare utilisation, we will track emergency department visits and hospitalisations via the patient's electronic health record. Additionally, we will collect participants' reasons for either stopping treatment or dropping out of the clinical trial, such as treatment adverse events, disease complications or scheduling issues with work.

Interventions

Licensed and oncology-experienced acupuncturists and massage therapists will deliver all treatments. All acupuncturists and massage therapists will be given a manual with the specific treatment protocols for acupuncture and massage (see online supplemental appendices 1 and 2) and will be trained by the PI and/or lead acupuncturist and massage therapist. For quality assurance, the lead therapists will audit at least two charts for each therapist per week for adherence to treatment protocol and documentation standards, and all therapists will be recertified twice yearly. We have extensive experience in conducting integrative medicine symptom trials including ensuring the quality of interventions.^{22 23 29 53–55}

For the acupuncture intervention, we will use a treatment protocol developed and tested by our group that has demonstrated improvements in pain, fatigue and sleep among patients with cancer.^{22 23 54–56} After sterilising the skin, the acupuncturist will place between 10 and 20 needles at a minimum of four local points around the body area with the most pain and at individual points depending on the participant's comorbid symptoms. The acupuncture needles will be inserted to appropriate depths depending on the location on the body and body type of the participant.⁵⁷ The acupuncturist will manipulate the needles to achieve the 'De Qi' sensation for the participants. 'De Qi' is a local sensation of soreness, numbness, or distension that accompanies the insertion and manipulation of needles during acupuncture.⁵⁸ The needles at the four local points for pain will be electrically stimulated at 2 Hz by connecting to a TENS unit. The acupuncturist will leave the needles in place for 20 min with brief manipulation at the beginning and end of the treatment.

For the massage intervention, we will use a treatment protocol developed and tested by our group that has shown improvements in pain and fatigue among patients with cancer undergoing chemotherapy.²⁹ Consistent with oncology massage practice, therapists will administer compressions with light to moderate pressure and will use any of the following oncology massage techniques: compression; muscle stripping; active/passive range of motion, post-isometric stretching; effleurage (gliding); myofascial release; positional release; and trigger/tender point release.^{59 60} Therapists will start with a 5 min protocol including guided diaphragmatic breathing exercise, rib mobilisations and occipital release to increase parasympathetic tone. Next, depending on the participant's primary area of pain, the therapist will focus 20 min of massage on that specific body area followed by effleurage toward the heart. The massage therapist will focus on the following identified areas of pain: head/jaw; cervical spine; thoracic spine; shoulder; upper extremity; lumbar; sacral; pelvic; hip and lower extremity.

Before each massage or acupuncture treatment, the massage therapist or acupuncturist will review the participant's current health status and modify his/her techniques if needed. In the case of acupuncture, shallow needling with minimal stimulation will be used, and needles will only be placed in the extremities. For participants with electronically charged medical devices, no stimulation will be used. In the case of massage, light touch will be used, and areas of bruising will be avoided. The massage therapist or acupuncturist will document any treatment modifications and the medical reason for the modification, which will allow us to systematically capture participants who received a modified treatment. Patients will be monitored for side effects at each visit. Adverse events related to the administration of either acupuncture or massage will be collected each week before and after each treatment by the acupuncturist/massage therapist or clinical research coordinator. The Common Terminology Criteria for Adverse Events V.5 will be used for toxicity evaluation.

Analytical approach

We will perform the analysis for each aim following the intention-to-treat (ITT) principle (ie, participants will be analysed according to the treatment group to which they will be randomly allocated regardless of drop-out or treatment adherence status). For all specific aims, our main analytic tool will be linear mixed-effects models (LMMs) because our primary outcome (worst pain severity) and secondary outcomes are repeated continuous outcomes over time.⁶¹ The general template of each LMM will model the outcome as a function of treatment arm and assessment time, controlling for the randomisation stratification variables (baseline opioid use and accrual site), and including a subject-specific random intercept and slope.

For aim 1, we will plot the outcome measure trajectories by randomisation arm over time and summarise each

outcome measure at each assessment time by treatment arm using descriptive statistics. Tests of ITT differences between randomisation arms with respect to changes in outcomes will be based on coefficients from specific time-by-arm interactions added to the general LMM template described above. Our primary effectiveness comparison will focus on changes in BPI Worst Pain from baseline to 26 weeks between acupuncture versus massage. Aim 1 secondary outcomes (eg, fatigue, insomnia, QOL) will be analysed using the same methods described above. We will also perform responder analyses by considering those who experienced 30% or greater reduction in BPI Worst Pain at end of treatment (week 10) as responders.^{34 62 63} We will compare the proportion of responders in acupuncture and massage at the end of the intervention period using descriptive cross-tabulations and logistic regression adjusting for the randomisation strata.

For aim 2, we will conduct exploratory, hypothesis-generating heterogeneity of treatment effect (HTE) analyses to identify patient-level factors associated with treatment response to either acupuncture or massage by incorporating relevant variables (eg, sex, race/ethnicity, expectation, opioid use) and variable-by-intervention interaction terms in linear regression models predicting week 26 worst pain controlling for baseline worst pain and stratification factors. Each variable of interest will be assessed for HTE in a separate model. For these exploratory regression analyses, we will guard against inflated type I error due to multiple testing by adjusting the variable-by-intervention interaction p values for the false discovery rate.^{64 65} Our current focus on evaluating and reporting HTE will be based on the approach proposed by Kent *et al.*⁶⁶ However, we will also apply promising emerging Bayesian^{67 68} and machine learning^{69 70} methods, which can identify HTE and subgroups based on multiple variables simultaneously and are potentially more powerful than traditional univariate methods. Since our inclusion criteria allows for patients with various cancer types, we will also perform exploratory subgroup analysis to see if there is any difference in treatment effect (both primary and secondary outcomes) among patients with solid tumour cancers vs blood cancers. Because our trial will enrol patients with advanced cancer, interventions may need to be modified for patient safety issues such as for those with low platelets or bruising in the area where there is pain. We will conduct exploratory analyses to examine if there are any differences in outcomes for those patients who received non-modified treatments versus those who had modified treatments. We will also conduct exploratory analyses to see whether individuals with low platelet counts experienced more adverse events compared with patients with normal platelet counts.

To address missing data, we will perform sensitivity analyses (eg, assess impact on results of adjusting for patient disease progression or death) and apply data analysis strategies that are as robust as possible to data losses. We will first explore whether missingness is associated with observed variables (eg, randomisation arm and

the baseline outcome measures) by comparing participants with complete and incomplete data. Of note, the LMMs described above validly include participants with incomplete data under the missing at random assumption. However, our exploration of the data may deem the missing at random assumption to be inappropriate; hence, we will perform sensitivity analyses to evaluate the robustness of our LMM results by refitting the models after imputing the missing week 26 outcomes using multiple imputation.

Power analysis and sample size

For our sample size/power considerations, we calculated the smallest standardised effect size (ie, Cohen's d) we will be able to detect with 80% power, given our sample size of 300 and other assumptions. Using the 'power.mmrn' function from the R package 'longpower,' we applied the formulas in Lu *et al*,⁷¹ to derive the smallest detectable effect size for the coefficient of the time-by-arm interaction term in our LMM, given our study design and assumptions, which we transformed to represent the standardised mean difference (ie, Cohen's d) between the two arms at 26 weeks postrandomisation. Based on our prior experience^{72 73} and given that patients living with advanced cancer may have unanticipated health issues (eg, hospitalisations, death), we conservatively anticipate lost to follow-up to be 20% by 26 weeks. Assuming this 20% lost to follow-up, correlation between baseline and 26-week BPI Worst Pain of 0.5, and two-sided alpha of 0.05, and with 150 participants in each of the two active intervention arms, we will have 80% power to detect an effect size of 0.35 (standardised mean difference, Cohen's d) at 26 weeks post-randomisation between acupuncture vs massage. Based on our own preliminary data in patients with stage IV cancer who experienced moderate to severe pain (N=284), the mean BPI Worst Pain score was 6.3 with SD of 1.7. A difference of 1 on the BPI Worst Pain score (considered a clinically meaningful difference in pain) based on SD of 1.7 equals an effect size (Cohen's d) of 0.59. In this study, we have 99% power to detect this clinically meaningful mean difference of 1 point (Cohen's d of 0.59) on the BPI Worst Pain score. Our trial is more than sufficiently powered to detect a clinically meaningful difference between acupuncture and massage at 26 weeks.

Patient and public involvement

Recognising the value of incorporating feedback from patients and their families, we organised a formal patient/stakeholder advisory board composed of 10 members (ie, patients, caregivers and stakeholders from advocacy and cancer organisations) to contribute to the study design, optimal delivery of interventions, recruitment and retention strategies, and implementation and dissemination efforts. By collaborating with patient/stakeholder partners, the patient perspective is included and helps to ensure that the research conducted is relevant and not unduly burdensome for patients. Our

patient/stakeholder advisory board members helped generate the research questions, choose the comparison groups, develop patient-centred inclusion and exclusion criteria, determine the timing of the primary endpoint, refine the research protocol, choose the most appropriate outcomes, decide on specific measurement tools and create patient-friendly recruitment materials. Throughout the project, our patient/stakeholder partners will have specific roles in recruitment activities and will help to ensure that our trial is accessible to participants from diverse communities. Additionally, patient/stakeholder partners' involvement will contribute to effectively translating and disseminating the study findings to patient, family, stakeholder and research audiences to effect real-world change.

DISCUSSION

Pain and comorbid fatigue and sleep disturbance are among the most common and distressing symptoms for patients living with advanced cancer.⁴⁻⁹ These co-occurring symptoms also negatively impact patients' QoL and functional performance.^{10 74 75} Unlike drug therapies that mostly focus on treating one symptom, acupuncture and massage can address multiple symptoms during treatment, which makes them potentially beneficial not only for pain but also for its related comorbid symptoms (eg, fatigue and sleep disturbance) among patients with advanced cancer. Acupuncture and massage are both widely available and commonly used nonpharmacological treatments for pain and other comorbid symptoms in cancer populations. Therefore, this RCT study will provide high quality evidence of the comparative effectiveness and durability of acupuncture versus massage that can be readily incorporated into clinical care to improve patient-centred decision making for pain management.

ETHICS AND DISSEMINATION

The institutional review board at MSK Cancer Center (MSK) approved the study protocol; most recent version of the protocol approved on 19 May 2021. For this trial, we will adhere to the guidelines from the Consolidated Standards of Reporting Trials⁷⁶ and Standards for Reporting Interventions in Clinical Trials of Acupuncture.⁷⁷ This trial is funded by the Patient-Centered Outcomes Research Institute (SMPAI-2018C2-12883).

The results of this study will be presented at national and international meetings, and a manuscript will be submitted for publication in a peer-reviewed journal. This research will inform which therapy (acupuncture or massage) is more effective for reducing pain and comorbid fatigue and insomnia in patients living with advanced cancer. Such information will lead to evidence-based and patient-centred decision making to incorporate these approaches for optimal pain management for the growing population of individuals living with advanced cancer. By collaborating with patient/stakeholder

partners, patient/stakeholder partners help to interpret both expected and unexpected study findings in a way that is culturally sensitive and relevant to patients' lived experiences. Patient/stakeholder partners' active involvement will contribute to effectively translating and disseminating the study findings to patient, family, stakeholder and research audiences to effect real-world change by providing education and awareness of the benefits of integrative, non-pharmacological options for pain management in people with advanced cancer.

Author affiliations

¹Department of Obstetrics, Gynecology, and Reproductive Sciences, University of California, San Diego, CA, USA

²Department of Psychiatry and Behavioral Sciences, Memorial Sloan Kettering Cancer Center, New York, NY, USA

³Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY, USA

⁴Integrative Medicine Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

⁵Office of Nursing Research, Memorial Sloan Kettering Cancer Center, New York, NY, USA

⁶Department of Biostatistics and Epidemiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

⁷Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

Acknowledgements The authors would like to thank the patients, healthcare providers, clinical staff, research staff and patient/stakeholder partners at all study sites for their contributions to this study.

Contributors SADR, REB, KP, JMac, DW, MB-B, JMao, AE: conceptualisation. REB, KP, JMao: data curation. REB, KP, JF: formal analysis. JMao: funding acquisition. SADR, NE, JMao: investigation. SADR, NE, JMao: project administration. JMao: resources. SADR, NE, KL, GD, HX, JMao: supervision. SADR, NE, JMao: writing-original draft. All authors: writing-review and editing.

Funding Research reported in this article was funded through a Patient-Centered Outcomes Research Institute (PCORI) award (SMPAI-2018C2-12883). The statements presented in this work are solely the responsibility of the authors and do not necessarily represent the views of the Patient-Centered Outcomes Research Institute (PCORI), its Board of Governors or Methodology Committee. It was also supported in part through the National Institutes of Health/National Cancer Institute Cancer Center Support Grant (P30 CA008748) and the Translational and Integrative Medicine Research Fund (award/grant number not applicable) at Memorial Sloan Kettering Cancer Center.

Disclaimer The funders do not have any role in study design; collection, management, analysis, and interpretation of data; writing of the report; and decision to submit for publication.

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially,

and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Jun J Mao <http://orcid.org/0000-0001-9229-0380>

REFERENCES

- American Cancer Society. *Cancer facts and statistics*. American Cancer Society, 2017.
- Delgado-Guay MO, Ferrer J, Ochoa J, *et al*. Characteristics and outcomes of advanced cancer patients who received palliative care at a public hospital compared with those at a comprehensive cancer center. *J Palliat Med* 2018;21:678–85.
- Cleeland CS, Zhao F, Chang VT, *et al*. The symptom burden of cancer: evidence for a core set of cancer-related and treatment-related symptoms from the eastern cooperative Oncology Group symptom outcomes and practice patterns study. *Cancer* 2013;119:4333–40.
- Johnsen AT, Petersen MA, Pedersen L, *et al*. Symptoms and problems in a nationally representative sample of advanced cancer patients. *Palliat Med* 2009;23:491–501.
- Kirkova J, Rybicki L, Walsh D, *et al*. The relationship between symptom prevalence and severity and cancer primary site in 796 patients with advanced cancer. *Am J Hosp Palliat Care* 2011;28:350–5.
- Dong ST, Butow PN, Costa DSJ, *et al*. Symptom clusters in patients with advanced cancer: a systematic review of observational studies. *J Pain Symptom Manage* 2014;48:411–50.
- Dong ST, Butow PN, Tong A, *et al*. Patients' experiences and perspectives of multiple concurrent symptoms in advanced cancer: a semi-structured interview study. *Support Care Cancer* 2016;24:1373–86.
- Walsh D, Donnelly S, Rybicki L. The symptoms of advanced cancer: relationship to age, gender, and performance status in 1,000 patients. *Support Care Cancer* 2000;8:175–9.
- Walsh D, Rybicki L. Symptom clustering in advanced cancer. *Support Care Cancer* 2006;14:831–6.
- Dodd MJ, Miaskowski C, Paul SM. Symptom clusters and their effect on the functional status of patients with cancer. *Oncol Nurs Forum* 2001;28:465–70.
- van den Beuken-van Everdingen MHJ, Hochstenbach LMJ, Joosten EAJ, *et al*. Update on prevalence of pain in patients with cancer: systematic review and meta-analysis. *J Pain Symptom Manage* 2016;51:e1079:1070–90.
- Porta-Sales J, Nabal-Vicuna M, Vallano A, *et al*. Have we improved pain control in cancer patients? A multicenter study of ambulatory and hospitalized cancer patients. *J Palliat Med* 2015;18:923–32.
- Tick H, Nielsen A, Pelletier KR, *et al*. Evidence-based nonpharmacologic strategies for comprehensive pain care: the Consortium pain Task force white paper. *Explore* 2018;14:177–211.
- Paice JA, Portenoy R, Lacchetti C, *et al*. Management of chronic pain in survivors of adult cancers: American Society of clinical oncology clinical practice guideline. *J Clin Oncol* 2016;34:3325–45.
- Swarm RA, Abernethy AP, Angheliescu DL, *et al*. Adult cancer pain. *J Natl Compr Canc Netw* 2013;11:992–1022.
- Network NCC. NCCN clinical practice guidelines in oncology: adult cancer pain version 1, 2018. Available: https://www.nccn.org/professionals/physician_gls/pdf/pain.pdf [Accessed 22 Jun 2018].
- Dowell D, Haegerich TM, Chou R. CDC guideline for prescribing opioids for chronic pain-United States, 2016. *JAMA* 2016;315:1624–45.
- Commission TJ. R3 report issue 11: pain assessment and management standards for hospitals, 2018. Available: https://www.jointcommission.org/r3_issue_11/ [Accessed 25 Jun 2018].
- Qaseem A, Wilt TJ, McLean RM. Clinical guidelines committee of the American College of P. noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2017;166:514–30.
- Mao JJ, Kapur R. Acupuncture in primary care. *Prim Care* 2010;37:105–17.
- Chiu HY, Hsieh YJ, Tsai PS. Systematic review and meta-analysis of acupuncture to reduce cancer-related pain. *Eur J Cancer Care* 2017;26:12457. doi:10.1111/ecc.12457
- Mao JJ, Liou KT, Baser RE, *et al*. Effectiveness of electroacupuncture or auricular acupuncture vs usual care for chronic musculoskeletal pain among cancer survivors: the peace randomized clinical trial. *JAMA Oncol* 2021;7:720–7.
- Mao JJ, Farrar JT, Bruner D, *et al*. Electroacupuncture for fatigue, sleep, and psychological distress in breast cancer patients with aromatase inhibitor-related arthralgia: a randomized trial. *Cancer* 2014;120:3744–51.
- Hershman DL, Unger JM, Greenlee H, *et al*. Effect of acupuncture vs sham acupuncture or waitlist control on joint pain related to aromatase inhibitors among women with early-stage breast cancer: a randomized clinical trial. *JAMA* 2018;320:167–76.
- Buckenmaier C, Cambron J, Werner R, *et al*. Massage therapy for pain-call to action. *Pain Med* 2016;17:1211–4.
- Boyd C, Crawford C, Paat CF, *et al*. The impact of massage therapy on function in pain Populations-a systematic review and meta-analysis of randomized controlled trials: part II, cancer pain populations. *Pain Med* 2016;17:1553–68.
- Kutner JS, Smith MC, Corbin L, *et al*. Massage therapy versus simple touch to improve pain and mood in patients with advanced cancer: a randomized trial. *Ann Intern Med* 2008;149:369–79.
- Lopez G, Liu W, Milbury K, *et al*. The effects of oncology massage on symptom self-report for cancer patients and their caregivers. *Support Care Cancer* 2017;25:3645–50.
- Mao JJ, Wagner KE, Seluzicki CM, *et al*. Integrating oncology massage into chemoinfusion suites: a program evaluation. *J Oncol Pract* 2017;13:e207–16.
- Cassileth BR, Vickers AJ. Massage therapy for symptom control: outcome study at a major cancer center. *J Pain Symptom Manage* 2004;28:244–9.
- Yun H, Sun L, Mao JJ. Growth of integrative medicine at leading cancer centers between 2009 and 2016: a systematic analysis of NCI-Designated comprehensive cancer center websites. *J Natl Cancer Inst Monogr* 2017;2017:4. doi:10.1093/jncimonographs/igx004
- Cleeland CS, Ryan KM. Pain assessment: global use of the brief pain inventory. *Ann Acad Med Singap* 1994;23:129–38.
- Hurst H, Bolton J. Assessing the clinical significance of change scores recorded on subjective outcome measures. *J Manipulative Physiol Ther* 2004;27:26–35.
- Farrar JT, Young JP, LaMoreaux L, *et al*. Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale. *Pain* 2001;94:149–58.
- Mendoza TR, Wang XS, Cleeland CS, *et al*. The rapid assessment of fatigue severity in cancer patients: use of the brief fatigue inventory. *Cancer* 1999;85:1186–96.
- Lin C-C, Chang A-P, Chen M-L, *et al*. Validation of the Taiwanese version of the brief fatigue inventory. *J Pain Symptom Manage* 2006;32:52–9.
- Bastien CH, Vallières A, Morin CM. Validation of the insomnia severity index as an outcome measure for insomnia research. *Sleep Med* 2001;2:297–307.
- Savard M-H, Savard J, Simard S, *et al*. Empirical validation of the insomnia severity index in cancer patients. *Psychooncology* 2005;14:429–41.
- Morin CM, Belleville G, Bélanger L, *et al*. The insomnia severity index: psychometric indicators to detect insomnia cases and evaluate treatment response. *Sleep* 2011;34:601–8.
- Smith AB, Selby PJ, Velikova G, *et al*. Factor analysis of the hospital anxiety and depression scale from a large cancer population. *Psychol Psychother* 2002;75:165–76.
- Moorey S, Greer S, Watson M, *et al*. The factor structure and factor stability of the hospital anxiety and depression scale in patients with cancer. *Br J Psychiatry* 1991;158:255–9.
- Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361–70.
- Hays RD, Bjorner JB, Revicki DA, *et al*. Development of physical and mental health summary scores from the patient-reported outcomes measurement information system (PROMIS) global items. *Qual Life Res* 2009;18:873–80.
- Revicki DA, Kawata AK, Harnam N, *et al*. Predicting EuroQol (EQ-5D) scores from the patient-reported outcomes measurement information system (PROMIS) global items and domain item banks in a United States sample. *Qual Life Res* 2009;18:783–91.
- Rothrock NE, Hays RD, Spritzer K, *et al*. Relative to the general US population, chronic diseases are associated with poorer health-related quality of life as measured by the patient-reported outcomes measurement information system (PROMIS). *J Clin Epidemiol* 2010;63:1195–204.
- Barile JP, Reeve BB, Smith AW, *et al*. Monitoring population health for healthy people 2020: evaluation of the NIH PROMIS(R) global health, CDC healthy days, and satisfaction with life instruments. *Qual Life Res* 2013;22:1201–11.

- 47 Colagiuri B, Smith CA. A systematic review of the effect of expectancy on treatment responses to acupuncture. *Evid Based Complement Alternat Med* 2012;2012:857804.
- 48 Kalaoukalan D, Cherkin DC, Sherman KJ, et al. Lessons from a trial of acupuncture and massage for low back pain: patient expectations and treatment effects. *Spine* 2001;26:1418–24.
- 49 Keefe JR, Amsterdam J, Li QS, et al. Specific expectancies are associated with symptomatic outcomes and side effect burden in a trial of chamomile extract for generalized anxiety disorder. *J Psychiatr Res* 2017;84:90–7.
- 50 Mao JJ, Armstrong K, Farrar JT, et al. Acupuncture expectancy scale: development and preliminary validation in China. *Explore* 2007;3:372–7.
- 51 Garland SN, Gehrman P, Barg FK, et al. Choosing options for insomnia in cancer effectively (CHOICE): design of a patient centered comparative effectiveness trial of acupuncture and cognitive behavior therapy for insomnia. *Contemp Clin Trials* 2016;47:349–55.
- 52 Robinson-Papp J, George MC, Wongmek A, et al. The quantitative analgesic questionnaire: a tool to capture patient-reported chronic pain medication use. *J Pain Symptom Manage* 2015;50:381–6.
- 53 Chen LX, Mao JJ, Fernandes S, et al. Integrating acupuncture with exercise-based physical therapy for knee osteoarthritis: a randomized controlled trial. *J Clin Rheumatol* 2013;19:308–16.
- 54 Mao JJ, Xie SX, Farrar JT, et al. A randomised trial of electroacupuncture for arthralgia related to aromatase inhibitor use. *Eur J Cancer* 2014;50:267–76.
- 55 Liou KT, Baser R, Romero SAD, et al. Personalized electroacupuncture versus auricular-acupuncture comparative effectiveness (PEACE): a protocol of a randomized controlled trial for chronic musculoskeletal pain in cancer survivors. *Medicine* 2020;99:e20085.
- 56 Mao JJ, Bruner DW, Stricker C, et al. Feasibility trial of electroacupuncture for aromatase inhibitor--related arthralgia in breast cancer survivors. *Integr Cancer Ther* 2009;8:123–9.
- 57 Cheng X. *Chinese acupuncture and moxibustion*. Beijing, China: Foreign Languages Press, 1987.
- 58 Mao JJ, Farrar JT, Armstrong K, et al. De qi: Chinese acupuncture patients' experiences and beliefs regarding acupuncture needling sensation-an exploratory survey. *Acupunct Med* 2007;25:158–65.
- 59 MacDonald G. *Medicine hands: massage therapy for people with cancer*. Simon and Schuster, 2014.
- 60 Walton T. *Medical conditions and massage therapy: a decision tree approach*. Lippincott Williams & Wilkins, 2010.
- 61 Tango T. Power and sample size for the S:T repeated measures design combined with a linear mixed-effects model allowing for missing data. *J Biopharm Stat* 2017;27:963–74.
- 62 Farrar JT, Portenoy RK, Berlin JA, et al. Defining the clinically important difference in pain outcome measures. *Pain* 2000;88:287–94.
- 63 Farrar JT, Pritchett YL, Robinson M, et al. The clinical importance of changes in the 0 to 10 numeric rating scale for worst, least, and average pain intensity: analyses of data from clinical trials of duloxetine in pain disorders. *J Pain* 2010;11:109–18.
- 64 Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J Royal Stat Soc Series B Methodol* 1995;57:289–300.
- 65 Benjamini Y, Yekutieli D. The control of the false discovery rate in multiple testing under dependency. *Ann. Statist.* 2001;29:1165–88.
- 66 Kent DM, Rothwell PM, Ioannidis JPA, et al. Assessing and reporting heterogeneity in treatment effects in clinical trials: a proposal. *Trials* 2010;11:85.
- 67 Henderson NC, Louis TA, Wang C, et al. Bayesian analysis of heterogeneous treatment effects for patient-centered outcomes research. *Health Serv Outcomes Res Methodol* 2016;16:213–33.
- 68 Wang C, Louis TA, Henderson NC, et al. beanz: an R package for Bayesian analysis of heterogeneous treatment effects with a graphical user interface. *J Stat Softw* 2018;85:31.
- 69 Imai K, Ratkovic M. Estimating treatment effect heterogeneity in randomized program evaluation. *Ann Appl Stat* 2013;7:443–70.
- 70 Wager S, Athey S. Estimation and inference of heterogeneous treatment effects using random forests. *J Am Stat Assoc* 2017;1–15.
- 71 Lu K, Luo X, Chen P-Y. Sample size estimation for repeated measures analysis in randomized clinical trials with missing data. *Int J Biostat* 2008;4:Article 9.
- 72 Mao JJ, Bowman MA, Xie SX, et al. Electroacupuncture versus gabapentin for hot flashes among breast cancer survivors: a randomized placebo-controlled trial. *J Clin Oncol* 2015;33:3615–20.
- 73 Mao JJ, Liou KT, Baser RE, et al. Effectiveness of electroacupuncture or auricular acupuncture vs usual care for chronic musculoskeletal pain among cancer survivors. *JAMA Oncol* 2021;7:720.
- 74 Dodd MJ, Cho MH, Cooper BA, et al. The effect of symptom clusters on functional status and quality of life in women with breast cancer. *Eur J Oncol Nurs* 2010;14:101–10.
- 75 Miaskowski C, Cooper BA, Paul SM, et al. Subgroups of patients with cancer with different symptom experiences and quality-of-life outcomes: a cluster analysis. *Oncol Nurs Forum* 2006;33:E79–89.
- 76 Moher D, Hopewell S, Schulz KF, et al. CONSORT 2010 explanation and elaboration: updated guidelines for reporting parallel group randomised trials. *Int J Surg* 2012;10:28–55.
- 77 MacPherson H, Altman DG, Hammerschlag R, et al. Revised standards for reporting interventions in clinical trials of acupuncture (stricta): extending the CONSORT statement. *PLoS Med* 2010;7:e1000261.

Appendix 1: Acupuncture Intervention

Background of the Intervention: The following acupuncture procedures and protocol were developed by Dr. Mao in consultation with experienced acupuncturists in China and U.S. The acupuncture points and techniques were selected to treat musculoskeletal pain and are based on classical and modern foundational acupuncture textbooks written in Chinese and/or English. We have piloted versions of this protocol in our prior research to demonstrate efficacy in pain reduction and adequate safety in the cancer population.

Operating Procedure:

Prior to the treatment session, the acupuncturist will:

- Review relevant medical history, laboratory results, and imaging studies to rule out absolute contraindications and to ensure appropriate precautions are taken.
- Greet patient/support members and escort them into private room.
- Take a focused history on pain and co-morbid symptoms (e.g. general aching, psychological distress, fatigue, or poor sleep).
- Conduct a focused physical examination with close attention to medical equipment (e.g. intravenous lines, chemotherapy ports) and areas of swelling/infection/deformities that may affect treatment protocol. If applicable, incorporate tongue/pulse diagnosis to guide acupuncture point selection.
- Assist patient onto table. Establish comfortable body positioning that is appropriate for treatment approach. Offer pillow, and/or bolster to maximize comfort.
- Instruct patient to adjust their clothing per his/her preferences and as indicated for treatment.

During the treatment session, the acupuncturist will:

- Insert needles by following the acupuncture treatment protocol as described below and in Tables 1 and 2.
- Offer blanket to maximize comfort.
- Dim lights and offer quiet music.

After the treatment session, the acupuncturist will:

- Assist patient to get off table.
- Ask patient to re-assess pain and other symptoms, evaluate for adverse events, and invite feedback to be incorporated into future treatment sessions.
- Complete clinical/research documentation.

Acupuncture Treatment Protocol (Total Duration: 30 Minutes)

- Identify one focal body area that the patient considers to be the most painful (e.g. neck, shoulder, back). This will be the primary area of focus for the entire treatment course.
- Choose at least four acupuncture points from Table 1 to address the primary area of pain. The acupuncturist may use clinical judgment to select additional acupuncture points or local trigger points ("ashi" or tender points) not listed in Table 1. The rationale for choosing unlisted points should be clearly documented. All selected points should be specified in clinical/research documentation.
- Choose at least four acupuncture points from Table 2 to address the patient's co-morbid symptoms. The acupuncturist may use clinical judgment to select additional acupuncture points not listed in Table 2. The rationale for choosing unlisted points should be clearly documented. All selected points should be specified in clinical/research documentation.
- Limit the total number of points to 10-20.
- Sanitize hands and clean the skin at needle insertion sites with alcohol pads using aseptic technique.
- Insert needle to appropriate depth with brief manual stimulation to achieve "De Qi" sensation.
- Connect TENS unit to four points near the primary area of pain by attaching positive/negative leads to the needles. Set electrical frequency at 2 Hz. Turn on TENS unit and gradually increase electrical intensity to appropriate level, i.e. the patient should feel the stimulation, but it should not be painful.
- Set timer for 20 minutes, then leave room.
- Document acupuncture procedure, including points used and total needle count.

- After 20 minutes, enter room, turn off TENS unit and detach leads from needles.
- Sanitize hands and remove needles and wipe any blood with a sterile cotton-tipped applicator.
- Perform a final needle count to ensure all needles were removed.
- Sanitize hands.

Table 1: Acupuncture Point Selection Guide for Musculoskeletal Pain						
Primary Pain Location	Acupuncture Points					
Head / Neck	Jianjing (G.B.21)	Huatuo (Extra)	Luozhen (Extra)	Dazhui (G.V.14)	Fengchi (G.B.20)	
Scapula	Tianzong (S.I.11)	Bingfeng (S.I.12)	Jianwaishu (S.I.14)	Gaohuangshu (U.B.43)		
Shoulder	Jianyu (L.I.15)	Jianliao (S.J.14)	Jianzhen (S.I.9)	Naoshu (S.I.10)	Houxi (S.I.3)	
Elbow	Quchi (L.I.11)	Chize (Lu. 5)	Tianjing (S.J.10)	Waiguan (S.J.5)	Hegu (L.I.4)	
Wrist	Yangchi (S.J.4)	Neiguan (P.C.6)	Daling (P.C.7)	Hegu (L.I.4)	Daling (L.I.5)	Yanglao (S.I.6)
Hand / Finger	Houxi (S.I.3)	Sanjian (L.I.3)	Baxie (Extra)	Hegu (L.I.4)		
Back	Shenshu (U.B.23)	Dachangshu (U.B.25)	Weizhong (U.B.40)	Chengshan (U.B.57)	Huatuo (Extra)	Kunlun (U.B.60)
Hip	Huantiao (G.B.30)	Yinmen (U.B.37)	Juliao (G.B.29)	Quixu (G.B.40)	Fengshi (G.B.31)	
Leg	Chengshan (U.B.57)	Feiyang (U.B.58)	Fengshi (G.B.31)			
Knee	Lianqiu (St.34)	Dubi (St.35)	Xiyan (Extra)	Yanlingquan (G.B.34)	Xiyangguan (G.B.33)	Yinlingquan (Sp.9)
Ankle	Jiexi (St.41)	Shangqui (Sp.5)	Quixu (G.B.40)	Kunlun (U.B.60)	Taixi (K.3)	
Foot / Toe	Gongsun (Sp.4)	Shugu (U.B.65)	Bafeng (Extra)	Taixi (Liv. 3)		
Table 2: Acupuncture Point Selection Guide for Addressing Co-Morbid Symptoms Associated with Pain						
Co-Morbid Symptoms	Acupuncture Points					
General Aching	Houxi (S.I.3)	Shenmai (U.B.62)	Dabao (Sp.21)	Geshu (U.B.17)	Yinlingquan (Sp.9)	Hegu/Taixi (L.I.4/Liv.3)
Psychological Distress	Neiguan (P.C.6)	Taixi (Liv.3)	Yin Tang (Extra)	ShenMen (Auricular)	Baihui (Du.20)	
Fatigue	Sanxinjiao (Sp.6)	Zusanli (St.36)	Qihai (CV6)			
Sleep	Shenmen (Ht.7)	Anmian (Extra)				

Appendix 2: Massage Intervention

Background of the Intervention: The following massage procedures and protocol were developed by experienced licensed oncology massage therapists in collaboration with Dr. Mao. This protocol is designed to treat musculoskeletal pain and is based on gold-standard textbooks in the field of oncology rehabilitation and medical massage. Versions of this protocol have been piloted in prior research to demonstrate efficacy in pain reduction and adequate safety in the cancer population.

Operating Procedure:

Prior to the treatment session, the massage therapist (MT) will:

- Greet patient/support members, escort them into private room, and sanitize hands.
- Take a focused history on pain and co-morbid symptoms.
- Review relevant medical history, laboratory results and imaging studies to rule out absolute contraindications and to ensure appropriate precautions are taken.
- Conduct a focused physical examination with close attention to medical equipment (e.g. intravenous lines, chemotherapy ports) and areas of swelling/infection/deformities that may affect treatment protocol. Visually assess posture and gait.
- Identify co-morbid complaints of the patient e.g. general aching, psychological distress, fatigue, or poor sleep.
- Assist patient onto table/chair. Establish comfortable body positioning that is appropriate for treatment approach. Offer blanket, pillow, and/or bolster to maximize comfort.
- Adjust patient's clothing per his/her preferences and as indicated for treatment.
- Identify patient's lubricant preference, i.e. oil or lotion.
- Dim lights and offer quiet music.

During the treatment session, the MT will:

- Follow the massage treatment protocol as described below in Table 1 and Table 2.
- Assess joints using active and/or passive pain-free range of motion. Use compression to identify and treat hypertonic muscles, tender/trigger points. Assess for trigger points with referral pattern to the site of pain. Use gentle fascial mobilizations to assess and treat fascial restriction in the area of pain, including surgical scars and tissues with a history of radiation treatment.
- Solicit and respond to patient feedback.

After the treatment session, the MT will:

- Assist patient to get off table/chair.
- Ask patient to re-assess pain and other symptoms, evaluate for adverse events, and invite feedback to be incorporated into future treatment sessions.
- Perform clinical/research documentation.

In subsequent treatment sessions, the MT will:

- Get patient feedback about the impact of the previous session in pain and co-morbid complaints.
- Adjust approach to respond to any shifts in primary area of pain if applicable.

Massage Treatment Protocol (Total Duration: 30 Minutes)

- Set silent timer for 5 minute intervals.
- Perform parasympathetic toning protocol for a minimum of 5 minutes. Assess breathing pattern. Provide verbal and physical cues for diaphragmatic breathing. Perform gentle release of

diaphragm and mobilization of ribs. Compress and muscle strip neck muscles. Perform OA release.

- Identify one focal body area that the patient considers to be the most painful (e.g. neck, shoulder, back, leg).
- Perform fascial release for a minimum of 5 minutes using techniques from the menu. Begin with indirect fascial release for tissues proximal to the site of pain. In subsequent sessions the treatment area may broaden in response to assessment and treatment outcomes.
- Perform tender point release for a minimum of 5 minutes using techniques from the menu. Begin with muscles proximal to the site of pain or with related referral patterns, in subsequent sessions the treatment area may broaden in response to assessment and treatment outcomes.
- Remaining time (2 - 3 minutes) can be used for integrative work distal to the pain site if indicated by gait and postural assessment or patient feedback during the treatment.
- End with effleurage (1-2 minutes) towards the heart, the therapist may use lubricant if necessary and appropriate.

Table 1: Recommended Massage Techniques

<ul style="list-style-type: none"> ▪ Palmar Compression (PC) ▪ Digital Compression (DC) ▪ Lifting/Pincer Compression (LC) ▪ Muscle Stripping (MS) ▪ Active Range of Motion (AROM) ▪ Passive Range of Motion (PROM) ▪ Post-Isometric PROM (PI-PROM) ▪ Post-Isometric AROM (PI-AROM) ▪ Positional Release (PR) ▪ Effleurage (E) ▪ Indirect/Gathering Fascial Release (IFR) ▪ Direct/Stretching Fascial Release (DFR) 	<ul style="list-style-type: none"> ▪ Superficial Fascial Release (SFR) ▪ Muscular/Deep Fascial Release (MFR) ▪ Kinetic Fascial Release (KFR) ▪ Long Duration Fascial Release (LDFR) ▪ Short Duration Fascial Release (SDFR) ▪ Compression of trigger/tender point (CTP) ▪ Positional release of trigger/tender point (P RTP) ▪ Local stretch for trigger/tender point (LSTP) ▪ Global/Muscle stretch for trigger/tender point (GSTP)
--	---

Table 2: Massage Guide Organized by Primary Location of Musculoskeletal Pain

Primary Pain Location	Recommended Massage Guide
HEAD/JAW Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.	<ul style="list-style-type: none"> ▪ Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis. ▪ Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate. ▪ Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. ▪ Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol. ▪ Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. ▪ Effleurage with strokes directed toward the heart.
CERVICAL SPINE Assess and treat local tissues first then progress	<ul style="list-style-type: none"> ▪ Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis.

<p>to proximal and distal areas. See muscle guides.</p>	<ul style="list-style-type: none"> ▪ Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate. ▪ Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. ▪ Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol. ▪ Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. ▪ Effleurage with strokes directed toward the heart.
<p>THORACICSPINE Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.</p>	<ul style="list-style-type: none"> ▪ Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis. ▪ Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate. ▪ Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. ▪ Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol. ▪ Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. ▪ Effleurage with strokes directed toward the heart.
<p>SHOULDER Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.</p>	<ul style="list-style-type: none"> ▪ Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis. ▪ Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate. ▪ Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. ▪ Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol. ▪ Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. ▪ Effleurage with strokes directed toward the heart.
<p>SHOULDER GIRDLE Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.</p>	<ul style="list-style-type: none"> ▪ Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis. ▪ Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate. ▪ Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. ▪ Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol.

	<ul style="list-style-type: none"> Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. Effleurage with strokes directed toward the heart.
CHEST/BREAST Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.	<ul style="list-style-type: none"> Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis. Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate. Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol. Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. Effleurage with strokes directed toward the heart.
RIB CAGE Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.	<ul style="list-style-type: none"> Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis. Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate. Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol. Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. Effleurage with strokes directed toward the heart.
UPPER EXTREMITY Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.	<ul style="list-style-type: none"> Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis. Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate. Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol. Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. Effleurage with strokes directed toward the heart.
ABDOMINAL Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.	<ul style="list-style-type: none"> Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis. Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate.

	<ul style="list-style-type: none"> ▪ Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. ▪ Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol. ▪ Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. ▪ Effleurage with strokes directed toward the heart.
LUMBAR Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.	<ul style="list-style-type: none"> ▪ Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis. ▪ Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate. ▪ Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. ▪ Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol. ▪ Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. ▪ Effleurage with strokes directed toward the heart.
SACRAL/PELVIC Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.	<ul style="list-style-type: none"> ▪ Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis. ▪ Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate. ▪ Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. ▪ Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol. ▪ Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. ▪ Effleurage with strokes directed toward the heart.
HIP Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.	<ul style="list-style-type: none"> ▪ Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis. ▪ Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate. ▪ Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping. ▪ Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol. ▪ Assess A/PROM in joints in the area, treat with pain-free joint mobilizations. ▪ Effleurage with strokes directed toward the heart.

LOWER EXTREMITY Assess and treat local tissues first then progress to proximal and distal areas. See muscle guides.	<ul style="list-style-type: none">▪ Assess fascial mobility with superficial fascial release of tissues in the focal area with attention to local scars/fibrosis.▪ Treat restriction with superficial indirect fascial release, tenting tissues over the painful area first. Progress to direct or kinetic fascial release where appropriate.▪ Compress primary muscles to assess for tension, treat hypertonicity with compression and muscle stripping.▪ Assess for trigger/tender points keeping in mind common referral patterns and treat with all appropriate elements of integrated TP release protocol.▪ Assess A/PROM in joints in the area, treat with pain-free joint mobilizations.▪ Effleurage with strokes directed toward the heart.
--	--