



BMJ Open Analysing 11 years of incidence trends, clinicopathological characteristics, and forecasts of colorectal cancer in young and old patients: a retrospective cross-sectional study in an Indonesian national referral hospital

Nur Rahadiani ^{1,2}, Muhammad Habiburrahman ¹, Murdani Abdullah,^{1,3,4} Wifanto Saditya Jeo,^{1,5} Marini Stephanie,^{1,2} Diah Rini Handjari,^{1,2} Ening Krisnuhoni^{1,2}

To cite: Rahadiani N, Habiburrahman M, Abdullah M, *et al.* Analysing 11 years of incidence trends, clinicopathological characteristics, and forecasts of colorectal cancer in young and old patients: a retrospective cross-sectional study in an Indonesian national referral hospital. *BMJ Open* 2022;**12**:e060839. doi:10.1136/bmjopen-2022-060839

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-060839>).

Received 05 January 2022
Accepted 15 August 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Nur Rahadiani;
nur.rahadiani@ui.ac.id

ABSTRACT

Objective To obtain annual incidence trends, understand clinicopathological characteristics, and forecast the future burden of colorectal cancer (CRC) in Indonesia.

Design 11-year retrospective cross-sectional study.

Setting A national referral hospital in Jakarta, Indonesia.

Participants Data from 1584 eligible cases were recorded for trends and forecasting analyses; 433 samples were analysed to determine clinicopathological differences between young (<50 years) and old (≥50 years) patients.

Methods Trend analyses were done using Joinpoint software, expressed in annual percentage change (APC), and a regression analysis was executed to generate a forecasting model. Patients' characteristics were compared using χ^2 or non-parametric tests.

Main outcomes Analysis of trends, forecasting model, and clinicopathological features between the age groups.

Results A significant increase in APC was observed among old patients (+2.38%) for CRC cases. Colon cancer increased remarkably (+9.24%) among young patients; rectal cancer trends were either stable or declining. The trend for right-sided CRC increased in the general population (+6.52%) and old patients (+6.57%), while the trend for left-sided CRC was stable. These cases are expected to be a significant health burden within the next 10 years. Patients had a mean age of 53.17±13.94, 38.1% were young, and the sex ratio was 1.21. Prominent characteristics were left-sided CRC, tumour size ≥5 cm, exophytic growth, adenocarcinoma, histologically low grade, pT3, pN0, inadequately dissected lymph nodes (LNs), LN ratio <0.05, no distant metastasis, early-stage cancer, no lymphovascular invasion, and no perineural invasion (PNI). Distinct features between young and old patients were found in the histological subtype, number of dissected LN, and PNI of the tumour.

Conclusions Epidemiological trends and forecasting analyses of CRC cases in Indonesian patients showed an enormous increase in colon cancer in young patients, a particularly concerning trend. Additionally, young patients

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This is the first retrospective cross-sectional study of Indonesian colorectal cancer (CRC) patients with a substantial data coverage period from 2009 to 2019.
- ⇒ We provide trend analysis to determine changes in the annual incidence of CRC in Indonesia based on age, tumour location, and side involvement of cancer, along with a forecasting model to estimate case patterns over the next 10 years.
- ⇒ This epidemiological study comprehensively analysed the difference in clinicopathological characteristics of CRC in young and old patients.
- ⇒ Data were taken from a single centre and might not be fully representative of other centres in Indonesia. Also, being a retrospective study, this study is susceptible to record bias and data loss from medical record retention and deterioration of pathology slides.
- ⇒ Data that could help explain the CRC trends, such as lifestyle, diet, alcohol use, tobacco use, family history, hereditary cancer syndromes, socioeconomic characteristics, and diagnostic test frequency, were not recorded.

exhibited particular clinicopathological characteristics that contributed to disease severity.

INTRODUCTION

Colorectal cancer (CRC) is the fourth most common cancer globally and is becoming more common in low-income and middle-income countries.¹ CRC is usually diagnosed through endoscopic biopsy or polypectomy. Microscopic examination is conducted to search for invasions. In the new era of personalised medicine, the role of anatomical

pathologists has been dramatically expanded. Their role is no longer limited to providing histopathologic diagnosis but also assessing staging, margins, and prognostic parameters that can only be made available by microscopic examinations such as tumour grade, lymphovascular invasion (LVI), and perineural invasion (PNI). Further research about the pathological characteristics of CRC is essential for treatment approaches and policy-making.

Recent long-term studies discovered that young people under 50 years old are more likely to get colon cancer, primarily in high-income countries.^{2–3} These studies' results suggest that the clinical, histopathological, and prognostic aspects of CRC epidemiology are also expected to encounter worrying changes.⁴ By 2030, the incidence of colon and rectal cancer in young people, for whom routine screening is currently not recommended, is projected to increase by 28–30% and 46–124%, respectively.⁵ Several Asian countries, including China, Japan, India, and South Korea, have also reported a tremendous rise in the number of young patients with CRC.^{2–6} This phenomenon is presumably due to rapid changes in lifestyle, diet, and genetic alterations in high-risk populations, particularly young adults.²

Epidemiological studies on CRC from other parts of Asia, including Southeast Asia, are needed since CRC cases are relatively less researched and are becoming a public health threat. Furthermore, in the population younger than 50, CRC shows a rising incidence and appears to display a more aggressive phenotype with unique genetic profiles, critical differences in somatic gene mutations, and gene methylation.⁷ Distinct molecular carcinogenesis and genomic profiles of CRC in Indonesia drove us to present a broader view of CRC in terms of epidemiology and clinicopathological characteristics,^{8–10} which has not been published by any previous investigation in Indonesia. These knowledge gaps motivated us to research how CRCs have changed from 2009 to 2019 in young patients compared with their older counterparts. We also aimed to obtain annual incidence trends, understand clinicopathological characteristics, and forecast the future burden of CRC in Indonesia.

MATERIALS AND METHODS

Study design, data collection, and selection process

This retrospective cross-sectional study was conducted at the Dr. Cipto Mangunkusumo Hospital, Jakarta, Indonesia, to analyse CRC incidence from 2009 to 2019 using pathological archives and hospital medical records.^{11–13} Data from 2020 were not included to avoid bias due to the COVID-19 pandemic, which caused a decrease in the number of patients with CRC attending the hospital. In total, 1958 patients have had a malignant tumour of the colon or rectum based on International Classification of Diseases, Tenth Revision (ICD-10) topography (C18–C20) and morphology codes (M8140/3, M8480/3, and M8490/3) with adequate biopsy or resection specimens eligible for enrolment in this study.¹⁴ For the analysis of

trends, forecasting, and clinical data, 1584 patients were selected by exclusion criteria (i.e., duplication of inputted cases, change of diagnosis or metastasis), with 433 resection samples undergoing a further analysis of pathological characteristics between two age groups, as shown in figure 1.

Extraction and definition of variables

The variables of age, registration year, sex, tumour site, colonic tumour location, side involvement, tumour subsites, and specimen type were extracted directly from cancer registry data. Data on tumour size, growth pattern, histological subtypes, tumour grade, pathological tumour (pT), node status (pN), adequacy of dissected lymph node (LN), lymph node metastasis (LNM), distant metastasis, staging, LVI, and PNI were retrieved from hospital medical records and pathological reports of patients who underwent surgery.

The young patient population was defined as subjects under 50 years of age, agreeing with previous studies.¹⁵ Pathological specimens of each patient were examined under the microscope by two independent pathologists who recorded the histopathology characteristics of: pathological tumour staging, histological subtypes, growth pattern, tumour grade, LVI, and PNI. We evaluated the number of dissected LNs in agreement with other studies and WHO guidelines, with a minimum of 12 LNs taken for each case.^{16–18} Along with LNs, we also calculated the LN ratio (LNR), defined as the number of positive LNs divided by the number of LNs examined. LNR was a significant predictor of survival in other malignancies and could be classified into subgroups according to the following cutoffs: <0.05 (LNR1), 0.05–0.20 (LNR2), 0.20–0.40 (LNR3), and 0.40–1.00 (LNR4).¹⁹

The tumour site was defined as the location where the primary tumour originated. A category of cancers known as right-sided CRC (RSCRC) originated from the caecum, ascending colon, hepatic flexure, and transverse colon. Meanwhile, left-sided CRC (LSCRC) originated from the splenic flexure, descending colon, sigmoid colon, and rectum.²⁰ Cancer of the caecum, ascending colon or transverse colon was referred to as proximal colon cancer. The descending colon or the sigmoid colon was the sites of distal colon cancer.^{14–21} Tumour size was defined as the largest dimension of the three-dimensional tumour, classified into <5 cm and ≥5 cm. Metastasis (distant metastasis) was confirmed by radiography or pathological diagnostic procedure. The WHO guideline and the American Joint Committee on Cancer eighth edition were the basis for pathological staging.^{17–18} Tumours with a stage of pT3–T4 or a pathological staging of pTNM III–IV were considered to be in the advanced stage.^{17–18} Tumours were also divided into three categories based on their subtypes: adenocarcinoma not otherwise specified (NOS), mucinous adenocarcinoma, and signet-ring cell carcinoma. The tumour growth pattern was classified into exophytic, endophytic, ulcerative, and linitis plastica.²² According to a WHO categorisation based on the percentage of




Figure 1 Study flow diagram for retrospective data collection, selection process, analysis of overall included samples and subgroup analysis of complete data in the final report. CRC, colorectal cancer; ICD-10, International Classification of Diseases, Tenth Revision.

gland formation in the tumour mass, tumour grade was grouped as well differentiated, moderately differentiated, and poorly differentiated.¹⁷ LVI and PNI were defined as the occurrence of each parameter in at least one slide of the pathology specimen sample.²³

Statistical analysis and presentation of data

A complete dataset from biopsy and resection specimens was used to extrapolate the CRC trend over 11 years, establish forecast models, and conduct a comparative analysis of the recorded variables. Missing data from the

retention of the medical records and deteriorated slides were omitted. In order to address the missing data and perform a more thorough analysis of pathological characteristics, this study employed a subgroup analysis for each measured outcome with more complete data from resection-only cases (figure 1).

Data were then recorded and processed using the SPSS V.25.0 statistical software with χ^2 and its alternative tests (Fisher's exact test, Kruskal-Wallis test or Mann-Whitney test). Analysis was performed for the young

and old patient populations for clinicopathological characteristics. The mean value of quantitative parameters (number of positive and dissected LNs, LNR and tumour size) was compared between two age groups with the Student's t-test. Annual incidence rates were quantified using the Joinpoint regression package provided by the US National Cancer Institute Surveillance Research Programme and National Cancer Institute (V.4.9.1.0).²⁴ Joinpoint regression analysis, established by Kim *et al*,²⁵ is a well-known approach used to study varying trends over time with Bonferroni adjustment.²⁶ It automatically joined separated time series of points (years) of cases on a logarithmic scale, expressed the trends as an annual percentage change (APC), and therefore, quantified the short-term increase or decrease between two successive points of change.^{24 25} A Monte Carlo permutation test assessed the significance of changing trends (i.e., APC).²⁷ Joinpoint regression analysis might be employed when the temporal trend of a given quantity (e.g., proportions, rates, counts), such as incidence and mortality (e.g., referring to cancer-related scenarios), was of interest.^{28–30} It is valuable to generate quantitative inferences instead of qualitative ones in epidemiological studies.^{24 31}

This presented study also performed linear and non-linear regression analyses to construct the best-fitted model to forecast the increasing trend of CRC cases in the next 10 years (2020–2029) using Minitab 19.1 (64-bit).^{32–38} The model trend equation to predict CRC cases can be visualised in linear [$Y_t = b_0 + (b_1 * t)$], quadratic [$Y_t = b_0 + b_1 * t + (b_2 * t^2)$], exponential [$Y_t = b_0 + (b_1^t)$], or S-curve (Pearl-Reed logistic) [$Y_t = (10^a) / (b_0 + b_1 * b_2 t)$] functions, with Y_t being the variable, b_0 being a constant, b_1 and b_2 being coefficients, and t as the value of the time unit. The best-fitted model is the model which has the lower values for three of these parameters: MAPE, mean absolute percent error; MAD, mean absolute deviation; and MSD, mean square deviation, or at least for two parameters, or having the lowest value for MAPE.^{36 39 40} The MAPE expresses accuracy as a percentage of the error. The MAD expresses accuracy in the same units as the data, which helps conceptualise the amount of error. The MSD measures the accuracy of the fitted time series. After deciding on the models, we measured the significance of their slope using the analysis of variance (ANOVA) test for curve estimation in SPSS. Statistical analyses with a $p < 0.05$ and a 95% CI for probability were considered significant.

Patient and public involvement statement

It was not possible to involve patients or the public in our research's design, conduction, reporting or dissemination plans. This report complied with the Strengthening the Reporting of Observational Studies in Epidemiology guidelines for observational studies (including cross-sectional studies), as stated in the Research Checklist.⁴¹

RESULTS

Of the 1584 people diagnosed with CRC in this study, males dominated the CRC cases registered in our centre, with a sex ratio (male: female) of 1.21. Distribution based on age groups, as shown in [table 1](#), demonstrated that the highest proportion of CRC was found in ages 51–60 years old; the mean age was 53.17 ± 13.94 years old, with females (52.28 ± 13.98 years old) generally being younger than males (53.90 ± 13.89 years old), $p = 0.021$. Looking at more specific age groupings, the number and proportion of patients' age was: 11–20 (11; 0.7%), 21–30 (81; 5.1%), 31–40 (225; 14.2%), 41–50 (339; 21.4%), 51–60 (432; 27.3%), 61–70 (334; 21.1%), 71–80 (135; 8.5%), 81–90 (20; 1.3%), and ≥ 91 (7; 0.4%). The mean age of the young patient population was surprisingly very young (38.82 ± 7.46 years old). The proportion of young patients in this centre reached 38.1% ($n = 604$) of the total incidence ($n = 1584$). Rectal cancer incidence was higher than colon cancer (64.3% vs. 35.7%). It was roughly equal for the percentage number of proximal and distal colon cancer (49.6% vs. 50.4%). Concerning tumour side involvement, LSCRC was still higher in proportion (82.3%). Of all cases of colon cancer, the sigmoid colon is the most often affected area, accounting for 13%.

[Figure 2](#) elucidates changes in the trend of CRC cases in Indonesian patients (denoted as an APC) over 11 years among all patients, as well as subcategorised by age groups (i.e., young and old patients), anatomical location of the tumour (i.e., colon, rectum or colon plus rectum) and side involvement of CRC (right sided vs. left sided). Using joinpoint regression analysis, a significant APC was observed among all patients, specifically in the annual incidence of colon cancer (+6.38%) and RSCRC (+6.52%). Among young patients, notable APC was only found in colon cancer (+9.24%); meanwhile, in the old patient group, a remarkable APC was noticed in CRC as a whole (+2.38%), colon cancer (+5.11%), and RSCRC (+6.57%). Trend patterns were positive for all tumour locations, except the rectum, which experienced stagnation in old patients (+0.58%) as well as dropped among the general population (−0.09%) and young patients (−0.97%) with $p > 0.05$. More detailed data on the trend analysis of our patients with CRC have been provided in the online supplemental files 1–3.

This study also investigated the increase of colon cancer based on their subsites (i.e., caecum, ascending colon, transverse colon, descending colon, and sigmoid colon), which is visualised in [figure 3](#). Significantly positive APC values were observed highest in the ascending colon (+10.60%), followed by the descending colon (+10.04%), the transverse colon (+9.88%), and the sigmoid colon (+5.84%). The caecum, on the other hand, displayed a slight negative trend with a low APC value (−0.98%, $p > 0.05$).

Additionally, as illustrated in [figure 4](#), several forecasting models of CRC incidences were generated using the best-fitted regression analysis. This approach predicted subsequent 10-year annual incidence rates for CRC cases

Table 1 Clinicopathological characteristics of tumours in young and old patients (n=1584)

Characteristics	Young patients (<50 years) (n=604)		Old patients (≥50 years) (n=980)		All patients (n=1584)		P-value
	N	%	n	%	N	%	
Registration year							0.931
2009	49	8.1	70	7.1	119	7.5	
2010	52	8.6	94	9.6	146	9.2	
2011	47	7.8	83	8.5	130	8.2	
2012	52	8.6	76	7.7	128	8.1	
2013	50	8.3	89	9.1	139	8.8	
2014	69	11.4	103	10.5	172	10.9	
2015	64	10.6	91	9.3	155	9.8	
2016	48	8.0	81	8.3	129	8.1	
2017	47	7.8	91	9.3	138	8.7	
2018	54	8.9	96	9.8	150	9.5	
2019	72	11.9	106	10.8	178	11.2	
Sex							0.056
Male	313	51.8	556	56.7	869	54.9	
Female	291	48.2	424	43.3	715	45.1	
Tumour site							0.002
Colon	187	31.0	379	38.7	566	35.7	
Rectal	417	69.0	601	61.3	1018	64.3	
Colonic tumour location							0.572
Proximal colon	96	51.3	185	48.8	281	49.6	
Distal colon	91	48.7	194	51.2	285	50.4	
Side involvement							0.131
RSCRC	96	15.9	185	18.9	281	17.7	
LSCRC	508	84.1	795	81.1	1303	82.3	
Tumour subsites							0.002
Caecum	20	3.3	58	5.9	78	4.9	
Ascending colon	52	8.6	75	7.7	127	8.0	
Transverse colon	24	4.0	52	5.3	76	4.8	
Descending colon	33	5.5	46	4.7	79	5.0	
Sigmoid	58	9.6	148	15.1	206	13.0	
Rectum	417	69.0	601	61.3	1018	64.3	
Specimen type							0.135
Biopsy	267	44.2	471	48.1	748	46.6	
Resection	337	55.8	509	51.9	846	53.4	

All statistical tests were done using the χ^2 test. The value with bold printed indicates the significant p-value. Percent values (%) were calculated as a percentage of the column total.

LSCRC, left-sided colorectal cancer; RSCRC, right-sided colorectal cancer.

using a specific case equation formula. What stands out in the analysis is that the dominance of colon cancer was expected to occur in the subsequent ten years among all groups (all, young, and old patients) with a significant increase in progression slope ($p < 0.001$, $p = 0.005$, and $p = 0.001$, respectively). Likewise, CRC future trends

in old patients would also steeply increase following the quadratic model ($p = 0.043$). A similar model existed in RSCRC cases, where all patients and old patients were forecasted to grow continually, with the corresponding p-value was 0.018 and 0.006. In contrast, the prediction of rectal cancer in all patients and old patients tended to




Figure 2 Trend analysis using joinpoint regression expressed by APC of CRC incidence among 1584 patients during 11 years period of study classified by tumour locations (colorectal, colon and rectum), and side involvement (RSCRC and LRSCRC) grouping in all, young, and old patients. A positive trend for 2009–2019 was observed among CRC, colon cancer, RSCRC, and LSCRC, while rectal cancer tended to stagnate and decrease in all groups. Colon plus rectum indicated a total incidence of both locations. Plotted lines indicate an APC. *Indicates that the APC significantly differs from zero at the alpha = 0.05 level using the logarithmically transformed data permutation model in joinpoint regression analysis. APC, annual percentage changes; 95%CI, 95% confidence interval; CRC, colorectal cancer; RSCRC, right-sided colorectal cancer; and LRSCRC, left-sided colorectal cancer.




Figure 3 Tumour subsites-specific incidence rate using jointpoint regression expressed by APC of CRC incidence among 1584 patients during 2009–2019 based on anatomical subsites of tumour in the colon. A sharp increase of cases by order in value was found in ascending, descending, transverse and sigmoid colon, respectively, while a gradual decline was observed in the caecum. *Denotes a significant change in APC vs. 0 ($p < 0.05$) using the logarithmically transformed data permutation model in jointpoint regression analysis. APC, annual percentage changes; 95%CI, 95% confidence interval; and CRC, colorectal cancer.

be constant and dropped among young patients in the following period. Although, at a glance, the remaining forecasts appeared will be likely increasing, their slope progression was not significant ($p > 0.05$). The precise number of predicted cases for the next 10 years (2020–2029) can be found in online supplemental files 4–6. As shown in online supplemental file 7 and table 2, the average future burden of CRC from 2020 to 2029 compared with the current 11-year data in all, young, and old patients was ~181 vs. 144 cases/year; ~67 vs. 55 cases/year; and ~113 vs. 89 cases/year, respectively.

As described in table 3, most tumour sizes were equal to or more than 5 cm (61%). Distant metastasis occurred in 6.9% of all cases. Most tumours were exophytic lesions (83.1%), adenocarcinoma NOS (85.2%), well differentiated (67.7%), with a pathological tumour staging of pT3 (66.6%), having inadequately dissected LNs (56.4%) and with category of LNR1 (57.5%), tumour stage IIA (34.2%), early stage (55.2%), without LVI (61.7%), and absence of PNI (88.7%). Comparing young and old patients, there were no significant differences in clinicopathological and histopathological characteristics except for histological subtypes, adequacy of LNs sampling, and PNI. Adenocarcinoma NOS was more prevalent in old patients than in their counterparts, while the mucinous

variant dominated in young patients ($p = 0.043$). Old patients were more likely to have inadequately dissected LN than young patients ($p = 0.004$). Young patients with CRC had more PNI than old patients ($p < 0.001$).

The comparison of means scores between two age groups highlighted in table 2 proves two significant differences in clinicopathological parameters of the tumor. First, the mean age value between the groups of young and old patients was extremely contrasted, with more than 23 years apart ($p < 0.001$). In addition, the average number of dissected LNs was significantly higher in the young patient group than in their older counterparts ($p = 0.004$).

Figure 5 portrays an example of microscopic tissue images from CRC cases. This figure highlights numerous key pathological markers essential for diagnosis, identifying histological patterns, and predicting prognosis.

DISCUSSION

This observational study was conducted to assess clinical trends of CRC over 11 years, forecast the future burden of CRC over the next 10 years and analyse the pathology of 1584 CRC cases in a national referral hospital in Indonesia. The current investigation corroborated previous




Figure 4 Annual incidence trends, the equation for predicting cases, and the forecast number of cases in the next 10 years using the best fitted-model regression analysis (linear, quadratic, exponential growth, or S-shaped curve model) for CRC classified by tumour locations (colorectal, colon, and rectum), and side involvement (RSCRC and LSCRC) in all, young, and old patients. Projection of a positive trend for the period 2020–2029 was observed among CRC, colon cancer, and LSCRC, while rectal cancer tended to stagnate and decrease. RSCRC was forecasted to have an increased burden in all and old patients but tended to decrease in young patients. Blue connected points show actual rates, red loosely dotted connected lines indicate a best-fitted trend, and the green densely connected dotted line indicates the forecasting trend. Y_t is the variable (equation for predicted cases) and t is the time unit (year) value. The significance test results for the slope of curve estimation employing the analysis of variance (ANOVA) statistical test.*Indicates a significant progression slope ($p < 0.05$). CRC, colorectal cancer; RSCRC, right-sided colorectal cancer; LSCRC, left-sided colorectal cancer; MAPE, mean absolute percent error; MAD, mean absolute deviation; and MSD, mean square deviation.

Table 2 Comparison of the mean value of clinicopathological parameters of tumour between young and old patients

Parameters	Mean±SD or number			P-value
	Young patients (<50 years)	Old patients (≥50 years)	All patients	
CRC (cases per year)	54.91±9.07	89.09±10.94	144.00±18.61	
Colon cancer (cases per year)	17.00±6.93	34.45±6.98	51.45±13.05	
Rectal cancer (cases per year)	37.91±5.20	54.64±8.62	92.55±12.40	
RSCRC (cases per year)	8.73±4.10	16.82±4.67	25.55±7.15	
LSCRC (cases per year)	46.18±7.04	72.27±9.33	118.45±14.69	
Age (years old)*	38.82±7.46	62.01±8.65	53.17±13.94	<0.001
Tumour size (cm)†	5.92±3.12	5.99±2.98	5.97±3.02	0.818
Smallest tumour size (cm)†	1.3	1.0	1.0	
Largest tumour size (cm)†	17.0	18.0	18	
Total count of positive LNs†	265	402	667	
Total count of dissected LNs†	1587	2726	4313	
Positive LNs†	1.84±3.37	1.39±2.34	1.54±2.73	0.107
Dissected LNs†	11.02±6.10	9.43±5.04	9.96±5.46	0.004
LNR†	0.18±0.29	0.18±0.30	0.18±0.29	0.964

All data were normally distributed and thus statistical tests were done using the independent and two-tailed student t-tests with equal variances were assumed (In Levene's test, the homogeneity assumption of the variance was met). The value with bold printed indicates the significant p-value.

*Assessed among 1584 patients.

†Assessed among 433 patients.

CRC, colorectal cancer; LNR, lymph node ratio; LNs, lymph nodes; LSCRC, left-sided colorectal cancer; RSCRC, right-sided colorectal cancer.

findings regarding men's predominance in CRC incidence. These findings are possibly due to men being more likely to smoke and drink alcohol (both risk factors for CRC), whereas women have higher levels of endogenous oestrogens, which protect against CRC carcinogenesis.⁴² Our study found that most CRC cases were identified in the middle-aged population, with peak incidence occurring between 51 and 60 years old, consistent with previous findings.⁴³ Female patients had a mean age younger than male patients, consistent with findings from an investigation conducted in Brunei Darussalam.⁴⁴ The definition of 'young patients' in an epidemiological study of CRC is arbitrary; this research employed 50 years as the cut-off age since this is the recommended age for first CRC screening in most screening programmes that have gained global adoption.¹⁵ Early-onset CRC is more likely to arise sporadically in third-world nations and is hypothesized to have a biologically and clinically unique entity, accounting for its aggressive presentation and poor prognosis.^{45–47} We report that CRC incidence among young patients reached nearly 40%, significantly higher than the rate reported in a previous Indonesian study on CRC between 2014 and 2016 with 275 samples (31.3%)⁴⁸, Western countries (7%)⁴⁹, and other Asian studies (6.7–35.5%).^{50 51} Other findings from South Asia were comparable to ours, with CRC incidence in young individuals ranging from 38% to 52%.^{46 52} The increasing proportion of young patients in our population may be influenced by the demographic profile of Indonesia,

which had a high proportion of people aged 50 or lower in 2019 (79.82%).⁵³

Trend analysis of CRC

The jointpoint regression analysis has significant analytic advantages for disease surveillance and, therefore, is valuable to portray trends in CRC incidence over time. This approach has been widely used in many CRC trend reports from several world regions, including North America (e.g., USA⁵⁴ and Canada⁵⁵), Latin America (e.g., Brazil⁵⁶ and Mexico⁵⁷), Europe (e.g., England⁵⁸ and Netherlands⁵⁹), East Asia (e.g., China⁶⁰ and South Korea⁶¹), the Middle East (e.g., Iran⁶² and Lebanon⁶³), and Southeast Asia (e.g., Vietnam⁶⁴ and Thailand⁶⁵). Using this method, our first Indonesian study also successfully identified short-term CRC trend patterns that differed between young and old patients. CRC incidence rates were modestly elevated in all and young patients with APC+2.23% and +1.98%, respectively with p>0.05. Meanwhile, the CRC cases among subjects aged ≥50 were mounting more dramatically, with an APC of +2.38% (p=0.041). A similar conclusion was reached by Pham *et al*⁶⁴ in the elderly Vietnamese population (APC+5.3%; 95% CI 2.8% to 7.9%). Hypothetically, this might happen because older patients were more likely to be included in screening programmes than young patients.⁶⁴ Since the population-based CRC screening programme has not been implemented in routine clinical practice in Indonesia, the actual rate of early-onset CRC might have been undervalued. The

Table 3 Pathological characteristics of tumour in young and old patients who underwent surgical resection with complete data (n=433)

Characteristics	Young patients (<50 years) (n=144)		Old patients (≥50 years) (n=289)		All patients (n=403)		P-value
	n	%	n	%	n	%	
Tumour size							0.559
<5 cm	59	41.0	110	38.1	169	39.0	
≥5 cm	85	59.0	179	61.9	264	61.0	
Growth pattern							0.412
Exophytic	120	83.3	240	83.0	360	83.1	
Endophytic	15	10.4	21	7.3	36	8.3	
Ulcerative	8	5.6	22	7.6	30	6.9	
Linitis plastica	1	0.7	6	2.1	7	1.7	
Histological subtypes							0.043
Adenocarcinoma NOS	115	79.9	254	87.9	369	85.2	
Mucinous	28	19.4	35	12.1	63	14.5	
Signet-ring cell	1	0.7	0	0.0	1	0.3	
Tumour grade							0.591
Well differentiated	97	67.4	196	67.8	293	67.7	
Moderately differentiated	31	21.5	69	23.9	100	23.1	
Poorly differentiated	16	11.1	24	8.3	40	9.2	
Pathological tumour staging							0.895
pT1	3	2.1	8	2.8	11	2.5	
pT2	22	15.3	44	15.2	66	15.2	
pT3	94	65.3	194	67.1	288	66.6	
pT4	25	17.3	43	14.9	68	15.7	
Pathological node status							0.734
pN0	80	55.6	168	58.1	248	57.3	
pN1a	18	12.5	42	14.5	60	13.8	
pN1b	18	12.5	34	11.8	52	12.0	
pN2a	18	12.5	33	11.4	51	11.8	
pN2b	10	6.9	12	4.2	22	5.1	
Adequacy of dissected LNs							0.004
Inadequate (<12)	67	46.5	177	61.2	244	56.4	
Adequate (≥12)	77	53.5	112	38.8	189	43.6	
LNR							0.967
LNR1 (<0.05)	81	56.2	168	58.1	249	57.5	
LNR2 (0.05–0.20)	21	14.6	38	13.1	59	13.6	
LNR3 (0.20–0.40)	15	10.4	28	9.7	43	10.0	
LNR4 (≥0.40)	27	18.8	55	19.1	82	18.9	
Lymph node metastasis							0.610
Yes	80	55.6	168	58.1	248	57.3	
No	64	44.4	121	41.9	185	42.7	
Distant metastasis							0.110
M0	138	95.8	265	91.7	403	93.1	
M1	6	4.2	24	8.3	30	6.9	
Staging							0.431

Continued

Table 3 Continued

Characteristics	Young patients (<50 years) (n=144)		Old patients (≥50 years) (n=289)		All patients (n=403)		P-value
	n	%	n	%	n	%	
I	19	13.2	41	14.2	60	13.8	
IIA	48	33.3	100	34.6	148	34.2	
IIB	12	8.3	19	6.6	31	7.2	
IIIA	6	4.2	10	3.5	16	3.7	
IIIB	39	27.1	83	28.7	122	28.1	
IIIC	15	10.4	16	5.5	31	7.2	
IV	5	3.5	20	6.9	25	5.8	
Degree of staging							0.921
Early stage (I–II)	79	54.9	160	55.4	239	55.2	
Advanced stage (III–IV)	65	45.1	129	44.6	194	44.8	
Lymphovascular invasion							0.314
Negative	84	58.3	183	63.3	267	61.7	
Positive	60	41.7	106	36.7	166	38.3	
Perineural invasion							<0.001
Negative	114	79.2	270	93.4	384	88.7	
Positive	30	20.8	19	6.6	49	11.3	

All statistical tests were done using the χ^2 test. The value with bold printed indicates the significant p-value. Percent values (%) were calculated as a percentage of the column total.
LNR, Lymph node ratio ; LNs, Lymph nodes; NOS, not otherwise specified.

estimated cost of treatment for patients with CRC in Indonesia is US\$116 083.37,⁶⁶ representing 0.000011% of the gross domestic product (GDP) in 2020.⁶⁷ The cost burden of treatment increases significantly as the disease progresses. In terms of screening costs, colonoscopy, and faecal testing range from US\$207 to US\$765 and US\$2.75 to US\$11, respectively. Given the rising CRC incidence and high cost of treatment, but the comparatively low cost of screening, our research implies that Indonesia should adopt a population-based CRC screening programme for high-risk populations, particularly those born after 1980. This early detection will benefit the public health sector and may further reduce the economic burden.

According to WHO, the prediction of CRC incidence in Indonesia from 2020 to 2025 was higher than the APC of trend analyses done in our study (CRC: +17.7% vs. +2.23%; colon cancer: +18.1% vs. +6.38%; and rectal cancer: +17.3% vs. -0.09%).⁶⁸ Further evidence showed that the trend of CRC cases among all Indonesian patients was lower in this study than in a study of patients with CRC in Tunisia from 1994 to 2009 (+2.23% vs. +3.90%).⁶⁹ We also discovered a smaller APC for CRC cases among young patients compared to a study among young Thai patients between 1989 and 2012 (+1.98% vs. +5.70%).⁶⁵ Trend analysis in [figure 2](#) reveals a sharp rise in colon cancer annually among young patients with a higher APC than in old and overall patients (+9.24% vs. +5.11% and +6.38%, respectively). In the last few decades, the incidence of CRC has been increasing in Asia, particularly

in Southeast Asian countries, including Indonesia and Malaysia.⁷⁰ If this trend continues, the number of CRC cases may suddenly overwhelm the healthcare system. Thus, better health policies should be constructed by the government.

The rise of CRC in young patients has not yet been fully elucidated. Early life exposure to the deleterious effects of risk factors, such as frequent smoking, alcohol consumption, obesity, a Western diet, reduced physical activity, and early-life antibiotic exposure, has been thought to increase susceptibility to CRC. The first contributor is smoking, associated with hypermethylation, microsatellite instability, and BRAF mutations in CRC carcinogenesis.⁷¹ Early-life smoking may contribute to the rising incidence of CRC in young individuals.⁷¹ Concerning that fact, 13.4% of Indonesian teenagers (95% CI 12.9% to 13.9%) and 27.3% of young adults (95% CI 26.8% to 27.8%) were found to be daily smokers, respectively.⁷² The risk of CRC is also increased by alcohol consumption, which is positively associated with the risk of cancer of distal colon and rectum among the Asian population.^{73 74} In Indonesia, alcohol consumption rose strikingly from 2000 to 2020, with the current proportion of alcohol consumption in teenagers and young adults being 4.0% (95% CI 3.8% to 4.3%) and 6.4% (95% CI 6.1% to 6.6%), respectively.⁷² Third attributed factor should be obesity, which has been linked with a higher risk of colon cancer in Asians. In Indonesia, obese and overweight individuals comprised roughly 4–8.8% of people aged 13–18 and




Figure 5 Histopathological features of colorectal cancer resection specimen (all in H&E staining). (A) well-differentiated adenocarcinoma NOS ($\times M40$); (B) poorly differentiated adenocarcinoma NOS ($\times M40$); (C) mucinous adenocarcinoma ($\times M40$, inset $\times M100$); (D) signet-ring cell carcinoma ($\times M40$, inset $\times M400$); (E) PT2 stage tumour infiltrating muscular layer ($\times M40$); (F) pT3 stage tumour infiltrating adipose tissue in subserosal layer ($\times M40$); (G) lymphovascular invasion (pointed by red arrow, $\times M40$); (H) perineural invasion (highlighted by yellow arrow, $\times M40$); (I) lymph node metastasis ($\times M100$). NOS, not otherwise specified.

nearly 21% of those over 18.^{72 75} It is not surprising that the obesity epidemic and the rise in colon cancer happen simultaneously. Many behaviours that are thought to cause weight gain, for instance, unhealthy eating habits and sedentary lifestyles, also raise the risk of CRC. Obesity can promote cancer formation through metabolic abnormalities, hyperinsulinaemia, systemic inflammation, and alteration of the gut microbiota.⁷⁶ An upward trend in CRC in Indonesia is also probably due to the acquisition of the Western diet as the fourth contributor.⁷⁷ This lifestyle trend has been seen in Indonesian teenagers who consume inadequate amounts of protein, fruits, and vegetables, but excessive amounts of sodium and fast food.⁷⁸ A recent study found that the de novo introduction of a Western-style high-fat, low-fibre diet induces inflammation and proliferation in the colonic mucosa within 2 weeks.⁷⁹ Increasing obesity is concurrent with reductions in physical activity levels.⁸⁰ A study in Japan revealed an inverse association between physical activity and CRC, and this association was stronger for colon cancer than rectal cancer.⁸¹ This fact agrees with a survey in Indonesia that found 33.5% (95% CI 33.3% to 33.8%) of the

population lacked physical activity in terms of time and frequency standards.⁷² Another related risk factor for CRC among Indonesians is early life and improper antibiotic use.⁸² These two risk factors could change the gut microbiota and metabolic profile, making a person more likely to have obesity later in life.⁸³

What stands out in the trend analysis of this study, is that the incidence rate of colon and rectal cancer was differed remarkably. In contrast with colon cancer which gradually increase (+5.11% to +9.24%), rectal cancer incidence has generally declined in all and young patients (−0.09% and −0.97%, respectively). Rectal cancer also remains stable in the old patients group (+0.58%). These findings could result from the rectum being more easily examined clinically during screening procedures than the colon, making precancerous lesions or suspected tumours easier to detect and removed during clinical examination of the rectum in screening. The negative trend of rectal cancer in this study was contradictory to the WHO's prediction,⁶⁸ but consistent with the trend in Canada.⁸⁴ After 1985, rectal cancer incidence slightly declined, with an APC of −0.38% among Canadians.⁸⁴ It also has been observed

that the trend of CRC subsite distribution progressively shifted to the proximal colon in various countries, such as the USA (1970–2000)⁸⁵, Japan (1974–1994)⁸⁶, and Norway (1962–2006).⁸⁷

Our findings emphasise that colon cancer incidence rose faster than rectal cancer in young patients (APC+9.24% vs. -0.97%), similar to results among Canadian young patients from 1969 to 2010 (APC+6.2% vs. +1.5%).⁸⁸ The APC of colon cancer in our population was higher than in Tunisia (+6.38% vs. +4.5%).⁸⁹ Some well-known risk factors do not exactly give a similar susceptibility towards colon and rectal cancer. The carcinogenic process may be different depending on where it happens.⁸⁴ Diet patterns, physical inactivity, and high body mass index have been linked to a higher risk of colon cancer, but not rectal cancer.^{73 89} Meanwhile, smoking and alcohol consumption have been linked to a higher risk of rectal cancer.^{90 91} Obesity, insulin resistance, and high blood glucose levels are connected with a higher risk of colon cancer because the colon is more insulin-sensitive than the rectum.^{92 93} We also hypothesised that women may have benefited from the preventive effect of hormones against cancer of distal colon and rectum. Endogenous hormones may have protected some women from developing cancer of distal colon and rectum. Increased use of exogenous hormones, such as hormone replacement therapy or oral contraceptives, might also have resulted in further reductions in these cancers.⁴² Respecting to that evidence, 61% of Indonesian women used contraceptive management between 2005 and 2012,⁹⁴ but this preventive effect has not been investigated for proximal (right-sided) colon tumours.⁸⁴

In contrast to earlier findings and the widely held belief that RSCRC is more common in young patients, this study revealed that RSCRC was more frequent in old patients, similar to a study from Germany.⁹⁵ Our results showed that most young patients had lesions in the left colon, in agreement with a hospital-based study in the Memorial Sloan Kettering Cancer Center, in the USA, where their young patients were more likely to have LSCRC.⁹⁶ A study also claimed that LSCRC is more prevalent in men and younger individuals, whereas RSCRC is more common in women and older people.²⁰ Accordingly, although genetic alterations might spread more in young patients and are typically related to the cause of RSCRC, it is still possible that LSCRC predominated in this group.

The trend analyses in figure 2 show that the APC of RSCRC rose statistically significantly among all patients (+6.52%) and old patients (+6.57%) over the study's 11-year period, with the largest APC being noticed in young patients (+6.59%). The causes of these patterns remain unclear; they might be due to inconsistent plotting of several incidences each year to follow a particular joined line to figure out a trend. The rising trend of RSCRC from 2009 to 2019 could be influenced by a lack of genetic counselling addressing age, specific syndromes, and family history in Indonesia, as RSCRC is usually associated with a genetic predisposition.⁹⁷ It is also challenging

to detect nonpolypoid (flat or depressed) tumours, more common in the right colon. These lesions are more likely to include carcinoma but are more difficult to detect and occur more frequently in high-risk individuals.⁹⁸ Higher colonoscopy miss rates may impede screening and identification of precancer and cancer lesions in the right colon, contributing to the rising trend of RSCRC.^{99–101 102} The presented study found that LSCRC had a positive steady trend among all (APC+1.41%), young (APC+1.37%), and old patients (APC+1.46%); all p-values were >0.05, similar to a report from Siegel *et al*¹⁰³ in the USA. The clinical implications of different proportion of side involvement between young and old patients was to the aggressiveness of the disease. RSCRCs are typically bulky, exophytic, polypoid lesions projecting into the lumen and causing significant anaemia. LSCRCs are infiltrating, constricting lesions encircling the lumen, often leading to obstruction.¹⁰⁴ A study implied that LSCRCs are genetically more unstable and phenotypically more aggressive due to distinct molecular biology patterns between RSCRC and LSCRC in DNA euploidy status, KRAS and p53 mutation rates.⁹⁵

Observing more specifically the trend of colon cancer based on its subsites, what can be seen in figure 3 is the significant growth of four of five colon subsites during the study period. The APC of ascending colon rose more quickly than APC in China from 2000 to 2004 (+10.60% vs. +2.25%).⁹¹ The transverse and descending colon had opposite results (+9.88% vs. -1.95% and +10.04% vs. -1.02%, respectively), while the sigmoid colon had a more positive trend (+5.84% vs. +4.19%).¹⁰⁵ Surprisingly, no differences in APC were found in the caecum (-0.98%), which had a slow and steady decline in cases. These trends aligned with the right-sided dominance during 11 years of study. Different parts of the colon may be more or less vulnerable to carcinogens because of biological differences in the intestine.¹⁰⁶ For example, genetic factors may play a significant role in developing proximal colon cancer, but factors like diet, exercise, and hormone use are more likely linked to distal colon cancer.¹⁰⁶

The trend analysis in this study enlightens us to narrow down patients in danger. Given the rapid economic transition and urbanisation occurring in Indonesia, it is possible to generalise the upward CRC incidence trend in a single centre in Jakarta to all of Indonesia,^{107 108} similar to what a study in Vietnam suggested.⁶⁴ However, as this is a single-centre study, the data presented may not be fully representative of other centres. Further research is needed to see if the trend can be reversed, for example, by evaluating current CRC screening standards and lowering the age at which people should begin screening. To reduce the upward trend, more studies are also required to investigate CRC risk factors in Indonesia. Our current study did not record data on risk factors of CRC that might help explain the trend of CRC found in the study. Furthermore, the cross-sectional design of this study did not allow us to establish any causal relationship.



Forecasting the CRC burden

In figure 4, we forecasted the future burden of CRC by performing a fit-model regression analysis to predict colon and rectal cancer incidences along with RSCRC and LSCRC. The model with a significant slope was found in all and old patients with colon cancer and RSCRC. Meanwhile, in young patients, the model with a significant slope was found only for colon cancer. Projection models for CRC, colon cancer, and rectal cancer follow the exponential growth curve pattern in all patients and young patient group. While, in the old patient group, colon cancer and CRC forecasting models use the quadratic model. In contrast, the projection model for rectal cancer follows the linear model in the old patient group. Compared with RSCRC, which follows the quadratic model, LSCRC was more varied, with the overall occurrences following the exponential growth curve, the young patients' incidences following the quadratic model, and the old patients' cases following the S-shaped model (sigmoid) curve.

The best-fitted model for forecasting CRC cases had different clinical implications based on curve shape. Addressing the interpretation of each curve was challenging since little robust research explains forecasting cancer incidence.¹⁰⁹ A linear trend is a forecasting model that develops a linear relationship between time and the response variable (incidence of disease). The linear model observed in rectal cancer among old patients means that cases increase gradually and linearly at a constant rate over time. This model assumption was based on forecast accuracy metrics and was supported by what has been pictured in trend analysis of rectal cancer with stable APC.¹¹⁰ What should be highlighted in this paper is that although the rectum was the most prevalent tumour site, we identified a negative trend or stable growth for this site in both joinpoint analysis and fit-model regression analysis for forecasting, similar to what was predicted in Japan.⁸⁶

Six of 15 scenarios were fitted into the quadratic curve model, a forecasting method that developed a non-linear relationship between time series and the response variable. The quadratic trend resembles a polynomial regression model that accurately captures the data trend.¹¹⁰ Following a quadratic model, the number of RSCRC cases was expected to steeply grow after 2019, particularly among all patients ($p < 0.05$) and old patients ($p < 0.01$). The literature corroborates that RSCRC is associated with several adverse prognostic factors: older age, advanced stage, and mucinous histological subtype.^{20 111 112} On the other hand, the incidence of RSCRC among young patients was projected to plummet until 2029 ($p > 0.05$). The future trend among young patients differs from the past between 2009 and 2019, which exhibited a steady movement. This pattern was similar to a study in the USA, which found that RSCRC increased initially, experienced stagnation, and was projected to fall by 2.3–2.6% annually.⁹⁷ The reasons for different past and future trends of RSCRC might be explained hypothetically by the increased use of colonoscopy in the early 21st century.¹¹⁴

Along with this direction, improved techniques and training for conducting colonoscopy in the right colon to screen, detect and diagnose may contribute to reducing RSCRC lesions among subclinical diseases.¹¹³ Another possible explanation for this result is that in the previous 11-year period, our young patients were dominated by a high proportion of patients with genetic factors, thus resulting in a higher trend of RSCRC cases in the young patients' group.^{8–10} In the next 10 years, the trend is predicted to shift to increasing rates of LSCRCs and RSCRCs otherwise due to greater exposure to specific cancer-related risk factors at the distal subsites.^{64 106 114} It is linked to the increasing adoption of a Westernised lifestyle in Indonesia, as also growing in other Asian countries, is a reasonable ground for this shift.^{84 115} The reasons for conflicting forecasts between young and old patients for CRC in overall and specifically RSCRC cases remain unclear. It might be explained by the complex attributions of risk factors associated with age and side of tumour involvement which has not been scrutinized in this study. Accordingly, further studies are urgently required in Indonesia to identify the contributing factors for the occurrence of CRC in each subsite, thus explaining the different trends based on subsite and side involvement.

Seven cases were forecasted following the exponential growth curve as the best-fitted model. Among these forecast models, two colon cancer cases elucidated significant progression slopes; they were in all patients ($p < 0.001$) and in young patients ($p < 0.01$). Their future trends were identical to past trends in the previous period, even were expected to be skyrocketing. Exponential growth curve has a J-shape, reflecting a growth whose rate is proportional to the size of the population over a specific period. Exponential growth curve modelling is a regression-based method for analysing longitudinal data (i.e., tracking the same sample at different points in time), suited to the projection of trends in one disease entity into a different period. The advantage of growth curve modelling over other methods is that this technique permits the testing of several types of trajectories until the one with the best fit to the data is found, and an output is far more precise than other statistical means.^{116 117} Exponential growth is distinguished by its slow start and, at some point, accelerating growth rate. The exponential growth curve has the fastest growth compared with the S-shaped, quadratic, and linear curves. This pattern causes an explosion of cases, relatively more than the S-shaped, which causes a relatively constant growth rate in the population.

One scenario of LSCRC among old patients following the sigmoid (S-shaped) curve trend model refers to a case whose growth rate decreases with the increasing number of individuals.¹¹⁰ An S-shaped curve is symmetric around the inflection point, which means that the case increases rapidly initially, followed by a slower rate after the inflection point than the rate postulated by the curve. Following this pattern, of movement of LSCRC cases have initial slow growth, reach a growth explosion, then at their upper limit, cases will be gradually steady, consistent

with insignificant slope progression ($p > 0.05$). The S-curve trend model is best for time series that follow a logistic manner.¹¹⁰ However, this model drawback may lead to underestimation and overestimation of the actual disease risk at the lower and upper tails of projected line.¹¹⁸

Projected CRC cases in Indonesia for the next 10 years confirm the future global burden of CRC, which is expected to increase by 60%, to over 2.2 million new cases in 2030.¹¹⁹ Looking specifically at online supplemental file 7, regarding cases predicted for 2020–2029, the burden of CRC remained high in our institution.

Distinct clinical and pathological features in young patients

Young individuals may be more susceptible to CRC due to genetic alterations and dietary changes; hence molecular profiles of young Indonesian patients with CRC have been identified to understand better the specific pathway involved in this group.⁸ Our young cases, mainly found in distal locations for CRC, are not in line with the characteristics of hereditary CRC, primarily found in proximal sites. They also did not follow the conventional pathways of sporadic CRC (the CIN pathway).⁸ Instead, carcinogenesis in these patients seems to have originated with MSI and inflammatory pathways, including cyclooxygenase-2 (COX-2) and nucleus factor κ B (NF- κ B). Also, lower mutation rates of the pro-oncogene KRAS are found among young Indonesian patients.⁸ Sudoyo *et al*¹²⁰ found that 56.5% of CRC cases were positively stained for MSH2 and 16.5% stained for MLH1. Moreover, signet-ring cell carcinoma—an aggressive subtype of CRC that spreads rapidly and is characterised by late symptom manifestations—disproportionately affects young individuals.¹²¹ It is also possible that the differences in the immune systems of young patients could play a role in age-related immunosenescence, T-cell dysfunction, and systemic inflammation.¹²²

Age is crucial due to its impact on prognosis. However, this idea is still debatable; some suggest worse outcomes at a young age,^{123 124} whereas others imply an equal prognosis between young and old age¹²⁵ depending on the staging reported.^{43 124} Contrary to other studies,^{46 126 127} where stage III–IV cancer predominates in the young age group, we found that more than half of our young patients with stage I–II cancer. However, no statistically significant difference in cancer staging between the two age groups was evident, similar to a prior report.¹²⁸ This might reflect increased awareness of the disease among young patients and primary care physicians, better access to colonoscopy, and more widespread use of CT with improved quality. Also, the introduction of national health insurance in the middle of the study period (2014) made access to healthcare more accessible, increasing people's concern for their health. Providing better facilities for cancer diagnosis may result in an inflation of the number of CRC and earlier detection of CRC through screening.¹²⁹ Patients with cancer found through screening show up at a much earlier stage of the disease than those not found through screening. Our study found no distinct clinical

characteristics between young and old patients regarding sex, side involvement, location, site or specimen type. There is no tendency for proximalisation of colon cancer in young patients compared with old patients in our study. Overall, the proximal and distal colon had an equal proportion of all CRC cases. However, if we included rectal cancer in the calculation of distal CRC, the proportion was in line with an extensive colonoscopy survey in Asia, which found that more patients had distal than proximal CRC.¹³⁰

Single institution and population-based studies have found that young patients with CRC have unique tumour locations, stages at presentation and histological features. Our findings were similar to those of these studies.^{131–134} The proportion of rectal cancer among young patients was significantly higher than in their old counterparts; as previously mentioned in an American study, 32% of CRC occurred in the rectum.¹³⁴ Looking more specifically at colon subsites, young patients with CRC mainly have lesions originating from the ascending and descending colon. Meanwhile, the caecum, transverse colon, and sigmoid colon were the most affected sites among old populations. Lesions with poorly defined histological features, such as mucinous and signet ring features, are more likely associated with poor outcomes.¹²³ They are also more resistant to chemotherapy.¹²⁸ Our results showed that the proportion of adenocarcinoma NOS in young patients was less frequent than in old patients, agreeing with a study by Chan *et al*⁵² and Gheju *et al*.¹³⁵ The mucinous histological variant was significantly higher in young than in old patients. Signet-ring cell cancer was only observed in young patients, accounting for only 0.6–1.0% of all CRC cases globally.¹³⁵ Our single patient who has signet-ring cell cancer has the following characteristics: 48 years, female, located in the caecum, right-sided, size 5.5 cm, brown-coloured surface, exophytic, adequate LNR 5/13, pT3N2aM0 (IIIB), no LVI, no PNI, and with poor tumour differentiation. Likewise, only one patient with signet-ring cell carcinoma was also identified in a Romanian study, but that patient was elderly (>50 years).¹³⁵ Signet-ring cancers have intracellular mucin pushing the nucleus to one side and are associated with a more advanced stage at diagnosis, a higher incidence of LVI, LNM, and liver metastases, a higher rate of recurrence, and higher aggression.^{136 137} The literature stated that mucinous histopathology was a significant predictor of poor outcomes and more advanced node stage.¹³⁸

The average number of dissected LNs in our study was lower than that in a recent Romanian study (mean: 9.96 ± 5.46 vs. median: 35.7 LNs removed), indicating that optimal LN sampling was a challenge in our institution.¹³⁵ Meanwhile, the average number of positive LNs per patient was lower than positive cases in Romania (mean: 1.54 ± 2.73 vs. median: 3.7 (1–62)).¹³⁵ The interpretation of LNM is thus more complicated because the number of dissected LNs was not ideal, but the positive number was favourable, which might be masking. More insufficiently removed LNs might result in a higher probability

of positive LNs in actual conditions due to unsuccessful LNs sampling, which could harm the detection of cancer spread. This issue may have an impact on patient staging. In contrast, increasing the number of dissected LNs leads to more accurate information about node status and more effective patient care. In a recent Dutch nationwide study,¹³⁹ authors found that with an increasing number of evaluated nodes, the risk of mortality is decreased, related to a better quality of surgical resection (yielding more LN for the pathologist to assess).

A closer inspection of the dissected LNs in [table 2](#) shows significant differences between the two age groups. The number of adequate LNs dissected in young patients was higher than in old patients, a favourable finding in young patients. Old patients are more likely to receive inadequate LN dissection during operative therapy, given their higher surgical risk for various postoperative complications and comorbid diseases. This concern possibly makes surgeons weigh the risks and benefits of a more thorough LN dissection.¹⁴⁰ Other contributing factors to the number of LNs dissected from resection include the surgeon's technique, bowel resection length, and tumour location.¹⁴¹ Complying with a minimum LN count of 12 is sometimes problematic, challenging, and less applicable. Thus, a novel measurement has been proposed to be used in clinical practice: LNR, a ratio of positive LNs to total dissected LNs. The mean score of LNR in our patients was lower than a median value of LNR in a study in Romania (mean: 0.18 ± 0.29 vs. median: 0.221 (0.139–1)).¹³⁵ This value was in line with the highest proportion of lower-category LNR (LNR1 was 57.5%) in the study analysis, implying favoured results. LNR provides a superior prognostic value than the number of positive nodes alone. A higher LNR is also significantly associated with poorer survival of CRC.¹⁴² Given no statistical difference in the LNR measurement between young and old patients in this study, it might be potential for LNR to be included as a predictive indicator in CRC staging systems for all patients.

This study found a lesser proportion of PNI in all CRC patients than in Elsamany *et al*¹⁴³ (11.3% vs. 24.4%). Nonetheless, we documented a significantly higher proportion of PNI in young patients than in old patients ($p < 0.001$), similar to findings in Zahir *et al*,⁴⁵ showing that 22% of their young CRC patients had positive PNI. PNI is associated with a higher rate of metastatic disease, a greater likelihood of recurrence, and poorer survival.¹⁴⁴ Several studies have also recognised it as a notable independent prognostic factor in CRC multivariate analysis.¹⁴⁴

Although some pathological features exhibited significant differences between the two age groups, no evidence was found for significant differences in tumour size, growth pattern, tumour grade, pT, pN, LNR, LNM, distant metastasis, and LVI. Two-thirds of patients had tumour size ≥ 5 cm, the most significant size being 18 cm. Although some authors believe that tumour size does not affect prognosis, others believe that tumour size partially affects prognosis.^{145 146} Increasing tumour size is

associated with decreased loco-regional control, resulting in an increased risk of malignant potential.¹⁴⁷ More extensive tumours are more likely to be more invasive and invade adjacent organs.¹⁴⁸ Local recurrence was significantly higher in patients with tumours measuring ≥ 5 cm in size, poorly differentiated adenocarcinoma, pT4 stage, and having adjuvant radiotherapy. Moreover, the 5-year overall survival rates in patients with tumours ≥ 5 cm were lower than those with a size < 5 cm (log-rank, $p = 0.001$).¹⁴⁹

According to our findings, the proportion of growth patterns (from highest to lowest, in both age groups) was exophytic, endophytic, ulcerative, and linitis plastica. These findings agree with a previous study in Thailand, which found that fungating and polyp mass (exophytic) were more common compared to ulcerative masses.¹⁴⁹ Our research revealed that exophytic growth patterns were prevalent in all patients and were distributed equally between the two age groups. Ulcerative growth and linitis plastica were much less common, which is favourable since both growth modes entail a worse prognosis. Linitis plastica suggests de novo origin, associated with a reduced proportion of KRAS mutations. Clinically, de novo tumours may represent a more aggressive subtype of CRC with a worse prognosis, poorer disease progression, and higher aggressiveness.¹⁰⁴ These results call for more awareness and persistence in detecting non-polypoid lesions, more intensive monitoring of colonoscopically treated cases, and surgery for selected patients.

Concerning tumour grading, most tumours in both age categories were well differentiated, similar to the results of a study from India.¹³⁸ These findings differed from those of a study by Chan *et al*,⁵² who discovered that both age groups were primarily affected by cases of moderately differentiated tumours. We identified that young patients were more likely to have poorly differentiated CRC than old patients. This finding shows how young patients have predilections for more aggressive tumour biology and implies a poorer prognosis regarding distinct tumour grade and histological subtypes distribution.^{124 150} However, although we found notable differences in the histological subtypes of young and old patients, no evidence was found for a significant association between tumour grade and age.

LVI was detected in almost two-fifths of individuals in this study. This proportion was fewer than in a previous report on Saudi patients (49.5%).¹⁴³ However, it was noticed that positive LVI cases were higher in our young patients than in old ones (41.7% vs. 36.7%). These findings suggest that LVI is a critical histopathological feature that needs to be assessed in every young patient with CRC, since literature mentioned its presence links to worse survival.¹⁴³

In short, all empirical findings related to clinicopathological characteristics of CRC in this study have provided a new understanding of this disease entity in Indonesia. Our study collected CRC data archived in one of Indonesia's national referral hospitals for cancer with a lengthy study period and is therefore the most robust data

accessible in our nation. Its coverage could represent CRC epidemiology on a regional scale since primary data for the entire country is not readily available. Another strength of this study was that we applied an efficient and noteworthy statistical method called joinpoint regression analysis to study the in-depth dynamics of CRC cases in Indonesia.^{31 151 152} This approach has allowed estimation of the magnitude of incidences, testing the movement of cases statistically, and clearly illustrating the direction of CRC trends.^{24 25 153} This study also provided several best-fitted models and computed forecasts that predict future trend patterns statistically.

However, our study should be interpreted with caution in light of the following limitations related to research methodologies. As a retrospective study, the quality of our database depends on the patient records and is subjective to record bias. We also excluded patients from our study due to retention of medical records or microscopic slide deterioration. This research may also have missed some old, frail patients with symptoms of CRC who were treated at home or in nursing homes without further investigation. Furthermore, several drawbacks might also arise concerning joinpoint regression analysis to measure the trend of cases. This method's common impediment was that it only offered a description of the time series based solely on yearly aggregated data¹⁵⁴; thus, it could not draw a causal relationship between possible risk factors that contributed to the findings.¹⁵⁵ As such, we could only hypothesise associations between CRC trends changes highlighted by our data and their possible influential factors supported by existing scientific evidence. Also, relying on the length of the study period, the software could only measure a certain number of year segments at a time.¹⁵³ A longer research term would have offered more freedom to measure the APC in several segmented sequences.¹⁵³ As a result, we could not compare several joinpoint segments to gain additional clarity regarding the impact of a specific intervention or event. The analysis could only be limited to 1 joinpoint because our samples only had 11 data points (i.e., 2009–2019).¹⁵³ To exemplify, given that Indonesia initially implemented universal health coverage in 2014, this limitation might restrict the analysis to distinguish different APCs between 2009–2013 and 2014–2019.

In addition, the projections of future CRC incidence discussed in this study should be carefully interpreted.¹⁰⁹ Predictions of future cancer incidence inherently depend on several uncertain factors, could be part of a larger cycle and may not persist into the future. Our projection of CRC in 2020–2029 was assumed to have similar clinicopathological characteristics as the circumstances observed from 2009 to 2019. Any changes affecting future cancer incidence rates beyond those included in the model's base years could not be statistically calculated by the forecasting models.¹⁵⁶ Dynamic evolutions in the population (e.g., advancing obesity or smoking rates and introducing new screening programmes with more cutting-edge technologies), governmental policy

adjustments, and emerging public health threats (e.g., pandemics) may influence the record of a predictive number of cases.¹⁵⁶ Trends and projections are volatile, and thus we could only forecast cases over a short period (e.g., 10 years in our study) to maintain forecasting accuracy. Moreover, this work did not include population-level data, and the mathematical prediction of cases in this study should be further validated using multicentre datasets.¹⁵⁷ Therefore, population and multicentre epidemiological studies are highly suggested to further predict trends in this disease entity

Despite all methodology-related limitations, our data showed a similar trend to other countries worldwide, primarily Asian countries. The incidence rates fit well into forecasting models, allowing clinicians and policy-makers to predict and anticipate future disease burdens of CRC.

CONCLUSION

This study sets out to assess clinical trends in CRC over 11 years based on tumour locations and side involvement, forecast the future incidence of CRC for the next 10 years, and analyse the clinicopathological profile among Indonesian patients in a single centre. Epidemiological trends and forecasting of CRC cases in Indonesian patients showed an enormous increase, notably for colon cancer, with a particularly concerning trend in young patients. Forecasts for the next 10 years using fit-model regression analysis found a significantly high number of CRC burdens in the future, particularly for colon cancer compared with rectal cancer, which is stable and declining. Additionally, young patients exhibited particular clinicopathological characteristics regarding tumour location, tumour subsites, histological subtypes, adequacy of dissected LNs and PNI, contributing to the disease's severity, aggressiveness, and prognosis. Multidisciplinary policies encompassing specialised screening protocols, extensive educational efforts, and lifestyle adjustments are required immediately to address this perplexing problem.

Author affiliations

¹Faculty of Medicine, Universitas Indonesia, Central Jakarta, DKI Jakarta, Indonesia

²Department of Anatomical Pathology, Dr. Cipto Mangunkusumo Hospital, Central Jakarta, DKI Jakarta, Indonesia

³Division of Gastroenterology, Pancreatobiliary, and Endoscopy, Department of Internal Medicine, Dr. Cipto Mangunkusumo Hospital, Central Jakarta, DKI Jakarta, Indonesia

⁴Human Cancer Research Center, Indonesia Medical Education and Research Institute (IMERI), Faculty of Medicine, Universitas Indonesia, Central Jakarta, DKI Jakarta, Indonesia

⁵Division of Digestive Surgery, Department of Surgery, Dr. Cipto Mangunkusumo Hospital, Central Jakarta, DKI Jakarta, Indonesia

Collaborators None declared.

Contributors NR was the principal investigator of this study, conceptualised the study, acquired funding, and controlled the decision to publish. As a guarantor, NR accepted full responsibility for the overall content of the work. NR and MH did the investigation, designed the methodology, had full access to the data, contributed to the analysis, drafted the paper, and did the project administration. MH was entirely responsible for software utilisation, data cleaning, and visualisation of research

findings. NR, MS, DRH, EK, MA, and WSJ collected the data, provided resources, and validated all data analyses. NR, EK, MA, and WSJ supervised the study process thoroughly. All authors critically revised the manuscript for important intellectual content, and all authors gave final approval for the version to be published.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval Ethical approval (KET-139/UN2.F1/ETIK/PPM.00.02/2020, protocol number: 10-11-1416) was obtained from the Institutional Ethical Review Board (IERB) of the Faculty of Medicine, Universitas Indonesia. General consent for the use of medical record data and residual material had already been obtained, in line with ethical approval.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. All data relevant to the study are included in the article or uploaded as online supplemental information. Raw data was obtained from a third party and are not publicly available. All additional relevant data analyses to the study have been uploaded as online supplemental information. To obtain more details data, please contact our corresponding author.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Nur Rahadiani <http://orcid.org/0000-0003-2342-8133>

Muhammad Habiburrahman <http://orcid.org/0000-0001-6372-8240>

REFERENCES

- Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394–424.
- Siegel RL, Torre LA, Soerjomataram I, et al. Global patterns and trends in colorectal cancer incidence in young adults. *Gut* 2019;68:2179–85.
- Araghi M, Soerjomataram I, Bardot A, et al. Changes in colorectal cancer incidence in seven high-income countries: a population-based study. *Lancet Gastroenterol Hepatol* 2019;4:511–8.
- Guastadisegni C, Colafranceschi M, Ottini L, et al. Microsatellite instability as a marker of prognosis and response to therapy: a meta-analysis of colorectal cancer survival data. *Eur J Cancer* 2010;46:2788–98.
- Bailey CE, Hu C-Y, You YN, et al. Increasing disparities in the age-related incidences of colon and rectal cancers in the United States, 1975–2010. *JAMA Surg* 2015;150:17–22.
- Sung JJY, Chiu H-M, Jung K-W, et al. Increasing trend in young-onset colorectal cancer in Asia: more cancers in men and more rectal cancers. *Am J Gastroenterol* 2019;114:322–9.
- Chang DT, Pai RK, Rybicki LA, et al. Clinicopathologic and molecular features of sporadic early-onset colorectal adenocarcinoma: an adenocarcinoma with frequent signet ring cell differentiation, rectal and sigmoid involvement, and adverse morphologic features. *Mod Pathol* 2012;25:1128–39.
- Abdullah M, Sudoyo AW, Utomo AR, et al. Molecular profile of colorectal cancer in Indonesia: is there another pathway? *Gastroenterol Hepatol Bed Bench* 2012;5:71–8.
- Yusuf I, Pardamean B, Baurley JW, et al. Genetic risk factors for colorectal cancer in multiethnic Indonesians. *Sci Rep* 2021;11:1–9.
- Abdullah M, Meilany S, Trimarsanto H. Genomic profiles of Indonesian colorectal cancer patients [version 1; peer review : awaiting peer review] 2022:1–11.
- Borovecki A, Mlinaric A, Horvat M, et al. Informed consent and ethics Committee approval in laboratory medicine. *Biochem Med* 2018;28:1–9.
- Junod V, Elger B. Retrospective research: what are the ethical and legal requirements? *Swiss Med Wkly* 2010;140:1–9.
- Gill SK, Gupta V, Bansal P. Informed consent status in observational studies with retrospective design: a poor show. *Asian J Pharm Clin Res* 2017;10:480–7.
- Lin JS, Piper MA, Perdue LA, et al. Screening for colorectal cancer: a systematic review for the U.S. preventive services Task force. *Evid Synth no 135* 2016;135:239.
- Mueller M, Schneider MA, Deplazes B, et al. Colorectal cancer of the young displays distinct features of aggressive tumor biology: a single-center cohort study. *World J Gastrointest Surg* 2021;13:164–75.
- Baxter NN, Virnig DJ, Rothenberger DA, et al. Lymph node evaluation in colorectal cancer patients: a population-based study. *J Natl Cancer Inst* 2005;97:219–25.
- WHO. WHO Classification of Tumours: Digestive System Tumours. In: *Who classification of tumours editorial board, editor. classification of tumours*. Geneva, Switzerland: World Health Organization, 2019: 1–537.
- Amin MB, Edge S, Greene F. *Ajcc cancer staging manual*. 8. New York: Springer, 2017: 252–4. <https://www.springer.com/gp/book/9783319406176>
- Akagi Y, Adachi Y, Kinugasa T, et al. Lymph node evaluation and survival in colorectal cancer: review of population-based, prospective studies. *Anticancer Res* 2013;33:2839–48.
- Baran B, Mert Ozupek N, Yerli Tetik N, et al. Difference between left-sided and right-sided colorectal cancer: a focused review of literature. *Gastroenterology Res* 2018;11:264–73.
- Huyghe JR, Harrison TA, Bien SA, et al. Genetic architectures of proximal and distal colorectal cancer are partly distinct. *Gut* 2021;70:1325–34.
- Wittekind C, Oberschmid B, Pathology CC. In: Schwab M, editor. In: Schwab M, ed. *Encyclopedia of cancer*. 925. 11. Berlin Heidelberg: Springer, 2017.
- Chen K, Collins G, Wang H, et al. Pathological features and prognostication in colorectal cancer. *Curr Oncol* 2021;28:5356–83.
- National Cancer Institute. Statistical Research and Applications Branch Joinpoint Trend Analysis Software [Internet]. Surveillance Research Program, 2021. Available: <https://surveillance.cancer.gov/joinpoint/> [Accessed cited 2021 Jun 9].
- Kim HJ, Fay MP, Feuer EJ, et al. Permutation tests for joinpoint regression with applications to cancer rates. *Stat Med* 2000;19:335–51.
- Hung BT, Long NP, Hung LP, et al. Research trends in evidence-based medicine: a joinpoint regression analysis of more than 50 years of publication data. *PLoS One* 2015;10:e0121054.
- Langton S, Lowe D, Rogers SN, et al. The impact of the UK 'two-week rule' on stage-on-diagnosis of oral cancer and the relationship to socio-economic inequalities. *J Cancer Policy* 2019;20:100191–8.
- Tyczynski JE, Berkel HJ. Mortality from lung cancer and tobacco smoking in Ohio (U.S.): will increasing smoking prevalence reverse current decreases in mortality? *Cancer Epidemiology, Biomarkers & Prevention* 2005;14:1182–7.
- John U, Hanke M, mortality Lcirrhosis. Liver cirrhosis mortality, alcohol consumption and tobacco consumption over a 62 year period in a high alcohol consumption country: a trend analysis. *BMC Res Notes* 2015;8:822.
- Doucet M, Rochette L, Hamel D. Incidence, prevalence, and mortality trends in chronic obstructive pulmonary disease over 2001 to 2011: a public health point of view of the burden. *Can Respir J* 2016;2016:7518287:1–10.
- Rea F, Pagan E, Compagnoni MM, et al. Joinpoint regression analysis with time-on-study as time-scale. Application to three Italian population-based cohort studies. *Epidemiol Biostat Public Heal* 2017;14:e12616:1–7.
- Hu YJ, Chen J, Zhong WS, et al. Trend analysis of betel Nut-associated oral cancer and health burden in China. *Chin J Dent Res* 2017;20:69–78.
- Claudio D, Miller A, Huggins A. Time series forecasting in an outpatient cancer clinic using common-day clustering. *IIE Trans Healthc Syst Eng* 2014;4:16–26.
- Burkhamer J, Kriebel D, Clapp R. The increasing toll of adolescent cancer incidence in the US. *PLoS One* 2017;12:1–16.

- 35 Ashoor AS, Kazem AAK, Gore S. An interactive network security for evaluating linear regression models in cancer mortality analysis and Self-Correlation of errors by using Durbin-Watson tests in Babylon/ Iraq. *J Phys Conf Ser* 2021;1804:012127.
- 36 Musa B. Comparison of various models on cancer rate and forecasting. *J Appl Sci Environ Manag* 2017;21:957–9.
- 37 Wah W, Stirling RG, Ahern S, *et al.* Forecasting of lung cancer incident cases at the small-area level in Victoria, Australia. *Int J Environ Res Public Health* 2021;18:5069–13.
- 38 Rashed ER, Eissa ME. Long-Term quantitative assessment of women survivability from cancer: a unique descriptive analysis. *Highlights Biosci* 2020;3:1–8.
- 39 Minitab Inc. Interpret all statistics and graphs for trend analysis [Internet]. Minitab Express Support, 2021. Available: <https://support.minitab.com/en-us/minitab-express/1/help-and-how-to/modeling-statistics/time-series/how-to/trend-analysis/interpret-the-results/all-statistics-and-graphs/> [Accessed 19 Jan 2021].
- 40 Tofallis C. A better measure of relative prediction accuracy for model selection and model estimation. *J Oper Res Soc* 2015;66:1352–62.
- 41 von EE, Altman DG, Egger M. Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ [Internet]* 2007 Oct 18;335:806 LP–8 <http://www.bmj.com/content/335/7624/806.abstract>
- 42 Murphy N, Strickler HD, Stanczyk FZ, *et al.* A prospective evaluation of endogenous sex hormone levels and colorectal cancer risk in postmenopausal women. *J Natl Cancer Inst* 2015;107:djv210.
- 43 O'Connell JB, Maggard MA, Liu JH, *et al.* Do young colon cancer patients have worse outcomes? *World J Surg* 2004;28:558–62.
- 44 Wong MC, Ding H, Wang J, *et al.* Prevalence and risk factors of colorectal cancer in Asia. *Intest Res* 2019;17:317–29.
- 45 Zahir MN, Mahpara E, Rafiq S, *et al.* Clinical features and outcome of sporadic colorectal carcinoma in young adults: a single center, cross sectional analysis. *JCO* 2013;31:e14680.
- 46 Gupta S, Bhattacharya D, Acharya AN, *et al.* Colorectal carcinoma in young adults: a retrospective study on Indian patients: 2000–2008. *Colorectal Dis* 2010;12:e182–9.
- 47 Bhurgri Y, Khan T, Kayani N, *et al.* Incidence and current trends of colorectal malignancies in an unscreened, low risk Pakistan population. *Asian Pac J Cancer Prev* 2011;12:703–8.
- 48 Anthonysamy MA, Indrayani Maker LPL, Gotra IM, *et al.* Prevalence of colorectal carcinoma based on microscopic type, sex, age and anatomical location in Sanglah General Hospital. *Intisari Sains Medis* 2020;11:272–6.
- 49 O'Connell JB, Maggard MA, Livingston EH, *et al.* Colorectal cancer in the young. *Am J Surg* 2004;187:343–8.
- 50 Chiang J-M, Chen M-C, Changchien CR, *et al.* Favorable influence of age on tumor characteristics of sporadic colorectal adenocarcinoma: patients 30 years of age or younger may be a distinct patient group. *Dis Colon Rectum* 2003;46:904–10.
- 51 Nath J, Wigley C, Keighley MRB, *et al.* Rectal cancer in young adults: a series of 102 patients at a tertiary care centre in India. *Colorectal Dis* 2009;11:475–9.
- 52 Chan KK, Dassanayake B, Deen R, *et al.* Young patients with colorectal cancer have poor survival in the first twenty months after operation and predictable survival in the medium and long-term: analysis of survival and prognostic markers. *World J Surg Oncol* 2010;8:1–11.
- 53 Central Bureau of Statistics Indonesia. Total Population by Age Group and Gender, 2019 [in Indonesian]. *BPS Indonesia* 2020 https://www.bps.go.id/indikator/indikator/view_data_pub/0000/api_pub/YW40a21pdTU1cnJxOGt6dm43ZEdoZz09/da_03/3
- 54 Cho MY, Siegel DA, Demb J, *et al.* Increasing Colorectal Cancer Incidence Before and After Age 50: Implications for Screening Initiation and Promotion of "On-Time" Screening. *Dig Dis Sci* 2022;67:4086–91.
- 55 Brenner DR, Heer E, Sutherland RL, *et al.* National trends in colorectal cancer incidence among older and younger adults in Canada. *JAMA Netw Open* 2019;2:e198090.
- 56 Dutra VGP, Parreira VAG, Guimarães RM. Evolution of mortality for colorectal cancer in Brazil and regions, by sex, 1996–2015. *Arq Gastroenterol* 2018;55:61–5.
- 57 Hoffman RM, Espey DK, Rhyne RL, *et al.* Colorectal cancer incidence and mortality disparities in New Mexico. *J Cancer Epidemiol* 2014;2014:1–8.
- 58 Exarchakou A, Donaldson LJ, Girardi F, *et al.* Colorectal cancer incidence among young adults in England: trends by anatomical sub-site and deprivation. *PLoS One* 2019;14:e0225547–13.
- 59 Swartjes H, Brouwer NPM, de Nes LCF, *et al.* Incidence, treatment and relative survival of early-onset colorectal cancer in the Netherlands since 1989. *Eur J Cancer* 2022;166:134–44.
- 60 Liu X, Bi Y, Wang H, *et al.* Different trends in colorectal cancer mortality between age groups in China: an age-period-cohort and joinpoint analysis. *Public Health* 2019;166:45–52.
- 61 Khil H, Kim SM, Hong S, *et al.* Time trends of colorectal cancer incidence and associated lifestyle factors in South Korea. *Sci Rep* 2021;11:1–12.
- 62 Mohammadi G, Akbari ME, Mehrabi Y, *et al.* Analysis of cancer incidence and mortality in Iran using joinpoint regression analysis. *Iran Red Crescent Med J* 2017;19.
- 63 Lakkis NA, El-kibbi O, Osman MH. Colorectal Cancer in Lebanon : Incidence, Temporal Trends, and Comparison to Regional and Western Countries 2021;28:1–12.
- 64 Pham DX, Phung AHT, Nguyen HD. Trends in colorectal cancer incidence in Ho Chi Minh City, Vietnam (1996–2015): joinpoint regression and age-period-cohort analyses. *Cancer Epidemiol* 2021;2022:102113.
- 65 Sarakarn P, Suwanrungruang K, Vatanasapt P, *et al.* Joinpoint analysis trends in the incidence of colorectal cancer in Khon Kaen, Thailand (1989 – 2012). *Asian Pac J Cancer Prev* 2017;18:1039–43.
- 66 Kristina SA, Endarti D, Wiedyaningsih C, *et al.* Estimating the burden of cancer and treatment cost related to alcohol consumption in Indonesia: a descriptive study. *Asian Pac J Cancer Prev* 2018;19:1845–9.
- 67 The World Bank. GDP (current US\$) - Indonesia [Internet], 2021. Available: <https://data.worldbank.org/indicator/NY.GDP.MKTP.CD?locations=ID> [Accessed 11 Sep 2021].
- 68 World Health Organization. Estimated number of new cases from 2020 to 2025, Both sexes, age [0-85+] Indonesia [Internet]. WHO, 2021[cited 2021 Jun 25]. p. Cancer Tomorrow. Available from: https://gco.iarc.fr/tomorrow/en/dataviz/bubbles?types=0&sexes=0&mode=cancer&group_populations=1&multiple_populations=0&multiple_cancers=1&cancers=8_9&populations=360&group_cancers=0&years=2025&single_unit=1000&bar_mode=grouped&key=total&show_bar_mode_prop
- 69 Khiari H, Ben Ayoub HW, Ben KH. Colorectal cancer incidence trend and projections in Tunisia (1994 - 2024). *Asian Pacific J Cancer Prev* 2017;18:2733–9.
- 70 Magaji BA, Moy FM, Roslani AC, *et al.* Descriptive epidemiology of colorectal cancer in University Malaya medical centre, 2001 to 2010. *Asian Pac J Cancer Prev* 2014;15:6059–64.
- 71 Limsui D, Vierkant RA, Tillmans LS, *et al.* Cigarette smoking and colorectal cancer risk by molecularly defined subtypes. *J Natl Cancer Inst* 2010;102:1012–22.
- 72 National Institute of Health Research and Development Indonesia. Basic Health Research [In Indonesian] [Internet]. *Jakarta* 2019. [Epub ahead of print: Available from] <https://www.litbang.kemkes.go.id/laporan-riiset-kesehatan-dasar-risikesdas/>
- 73 Shin A, Joo J, Bak J, *et al.* Site-Specific risk factors for colorectal cancer in a Korean population. *PLoS One* 2011;6:e23196.
- 74 Akhter M, Kuriyama S, Nakaya N, *et al.* Alcohol consumption is associated with an increased risk of distal colon and rectal cancer in Japanese men: the Miyagi cohort study. *Eur J Cancer* 2007;43:383–90.
- 75 Oddo VM, Maehara M, Rah JH. Overweight in Indonesia: an observational study of trends and risk factors among adults and children. *BMJ Open* 2019;9:e031198–14.
- 76 Wong SH, Yu J. Gut microbiota in colorectal cancer: mechanisms of action and clinical applications. *Nat Rev Gastroenterol Hepatol* 2019;16:690–704.
- 77 Yee YK, Tan VPY, Chan P, *et al.* Epidemiology of colorectal cancer in Asia. *J Gastroenterol Hepatol* 2009;24:1810–6.
- 78 Rachmi CN, Jusril H, Ariawan I, *et al.* Eating behaviour of Indonesian adolescents: a systematic review of the literature. *Public Health Nutr* 2021;24:s84–97.
- 79 O'Keefe SJ, Li JV, Lahti L, *et al.* Fat, fibre and cancer risk in African Americans and rural Africans. *Nat Commun* 2015;6:6342.
- 80 Akimoto N, Ugai T, Zhong R, *et al.* Rising incidence of early-onset colorectal cancer - a call to action. *Nat Rev Clin Oncol* 2021;18:230–43.
- 81 Pham NM, Mizoue T, Tanaka K, *et al.* Physical activity and colorectal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. *Jpn J Clin Oncol* 2012;42:2–13.
- 82 Parathon H, Kuntaman K, Widiastoety TH, *et al.* Progress towards antimicrobial resistance containment and control in Indonesia. *BMJ* 2017;358:j3808–5.

- 83 Chen L-W, Xu J, Soh SE, *et al.* Implication of gut microbiota in the association between infant antibiotic exposure and childhood obesity and adiposity accumulation. *Int J Obes* 2020;44:1508–20.
- 84 Gibbons L, Waters C, Mao Y, *et al.* Trends in colorectal cancer incidence and mortality. *Health Rep* 2001;12:41–55.
- 85 Cucino C, Buchner AM, Sonnenberg A. Continued rightward shift of colorectal cancer. *Dis Colon Rectum* 2002;45:1035–40.
- 86 Nakagawa H, Ito H, Hosono S, *et al.* Changes in trends in colorectal cancer incidence rate by anatomic site between 1978 and 2004 in Japan. *Eur J Cancer Prev* 2017;26:269–76.
- 87 Larsen IK, Bray F. Trends in colorectal cancer incidence in Norway 1962–2006: an interpretation of the temporal patterns by anatomic subsite. *Int J Cancer* 2010;126:721–32.
- 88 Patel P, De P. Trends in colorectal cancer incidence and related lifestyle risk factors in 15–49-year-olds in Canada, 1969–2010. *Cancer Epidemiol* 2016;42:90–100.
- 89 Wei EK, Giovannucci E, Wu K, *et al.* Comparison of risk factors for colon and rectal cancer. *Int J Cancer* 2004;108:433–42.
- 90 Cheng J, Chen Y, Wang X, *et al.* Meta-Analysis of prospective cohort studies of cigarette smoking and the incidence of colon and rectal cancers. *Eur J Cancer Prev* 2015;24:6–15.
- 91 Park S-Y, Wilkens LR, Setiawan VW, *et al.* Alcohol intake and colorectal cancer risk in the Multiethnic cohort study. *Am J Epidemiol* 2019;188:67–76.
- 92 Giovannucci E. Insulin and colon cancer. *Cancer Causes Control* 1995;6:164–79.
- 93 Komninou D, Ayonote A, Richie JP, *et al.* Insulin resistance and its contribution to colon carcinogenesis. *Exp Biol Med* 2003;228:396–405.
- 94 Ministry of Health of Indonesia. The Situation of Family Planning in Indonesia [In Indonesian]. *Bul Jendela Data dan Inf Kesehatan* 2013;2:1–44.
- 95 Benedix F, Kube R, Meyer F, *et al.* Comparison of 17,641 patients with right- and left-sided colon cancer: differences in epidemiology, perioperative course, histology, and survival. *Dis Colon Rectum* 2010;53:57–64.
- 96 Quah HM, Joseph R, Schrag D, *et al.* Young age influences treatment but not outcome of colon cancer. *Ann Surg Oncol* 2007;14:2759–65.
- 97 Siegel RL, Ward EM, Jemal A. Trends in colorectal cancer incidence rates in the United States by tumor location and stage, 1992–2008. *Cancer Epidemiol Biomarkers Prev* 2012;21:411–6.
- 98 Soetikno RM, Kaltenbach T, Rouse RV, *et al.* Prevalence of nonpolypoid (flat and depressed) colorectal neoplasms in asymptomatic and symptomatic adults. *JAMA* 2008;299:1027–35.
- 99 Bressler B, Paszat LF, Chen Z, *et al.* Rates of new or missed colorectal cancers after colonoscopy and their risk factors: a population-based analysis. *Gastroenterology* 2007;132:96–102.
- 100 Boursi B, Halak A, Umansky M, *et al.* Colonoscopic screening of an average-risk population for colorectal neoplasia. *Endoscopy* 2009;41:516–21.
- 101 Singh A, Kuo Y-F, Riall TS, *et al.* Predictors of colorectal cancer following a negative colonoscopy in the Medicare population. *Dig Dis Sci* 2011;56:3122–8.
- 102 Imperial R, Ahmed Z, Toor OM, *et al.* Comparative proteogenomic analysis of right-sided colon cancer, left-sided colon cancer and rectal cancer reveals distinct mutational profiles. *Mol Cancer* 2018;17:177.
- 103 Siegel RL, Jemal A, Ward EM. Increase in incidence of colorectal cancer among young men and women in the United States. *Cancer Epidemiol Biomarkers Prev* 2009;18:1695–8.
- 104 Papagiorgis PC, Zizi AE, Tseleni S, *et al.* Clinicopathological differences of colorectal cancers according to tumor origin: Identification of possibly *de novo* lesions. *Biomed Rep* 2013;1:97–104.
- 105 Zhou Q, Li K, Lin G-Z, *et al.* Incidence trends and age distribution of colorectal cancer by subsite in Guangzhou, 2000–2011. *Chin J Cancer* 2015;34:1–7.
- 106 Bufill JA. Colorectal cancer: evidence for distinct genetic categories based on proximal or distal tumor location. *Ann Intern Med* 1990;113:779–88.
- 107 Schröders J, Wall S, Hakimi M, *et al.* How is Indonesia coping with its epidemic of chronic noncommunicable diseases? A systematic review with meta-analysis. *PLoS One* 2017;12:e0179186.
- 108 Angkurawaranon C, Jiraporncharoen W, Chenthanakij B, *et al.* Urbanization and non-communicable disease in Southeast Asia: a review of current evidence. *Public Health* 2014;128:886–95.
- 109 Bray F, Møller B. Predicting the future burden of cancer. *Nat Rev Cancer* 2006;6:63–74.
- 110 Ismail L, Materwala H, Znati T, *et al.* Tailoring time series models for forecasting coronavirus spread: case studies of 187 countries. *Comput Struct Biotechnol J* 2020;18:2972–3206.
- 111 Wang CB, Shahjehan F, Merchea A, *et al.* Impact of tumor location and variables associated with overall survival in patients with colorectal cancer: a Mayo clinic colon and rectal cancer registry study. *Front Oncol* 2019;9:76.
- 112 Warschkow R, Sulz MC, Marti L, *et al.* Better survival in right-sided versus left-sided stage I–III colon cancer patients. *BMC Cancer* 2016;16:554.
- 113 Siegel RL, Fedewa SA, Anderson WF, *et al.* Colorectal cancer incidence patterns in the United States, 1974–2013. *J Natl Cancer Inst* 2017;109:27–32.
- 114 Kuipers EJ, Grady WM, Lieberman D, *et al.* Colorectal cancer. *Nat Rev Dis Primers* 2015;1:15065.
- 115 Center MM, Jemal A, Smith RA, *et al.* Worldwide variations in colorectal cancer. *CA Cancer J Clin* 2009;59:366–78.
- 116 Hardy MA, Bryman A. *Handbook of Data Analysis [Internet]*. London: SAGE Publications Ltd.; 2004: 1–728. <https://us.sagepub.com/en-us/nam/handbook-of-data-analysis/book209824>
- 117 Nini A, Corradini C, Guo D, *et al.* The application of growth curve modeling for the analysis of diachronic corpora. *Lang Dyn Change* 2017;7:102–25.
- 118 Devlin SM, Satagopan JM. Statistical interactions from a growth curve perspective. *Hum Hered* 2016;82:21–36.
- 119 Rawla P, Sunkara T, Barsouk A. Epidemiology of colorectal cancer: incidence, mortality, survival, and risk factors. *Prz Gastroenterol* 2019;14:89–103.
- 120 Sudoyo AW, Hernowo B, Krisnuhoni E, *et al.* Colorectal cancer among young native Indonesians: a clinicopathological and molecular assessment on microsatellite instability. *Med J Indones* 2010;19:245–51.
- 121 Gopalan V, Smith RA, Ho Y-H, *et al.* Signet-ring cell carcinoma of colorectum—current perspectives and molecular biology. *Int J Colorectal Dis* 2011;26:127–33.
- 122 Foster AD, Sivarapatna A, Gress RE. The aging immune system and its relationship with cancer. *Aging health* 2011;7:707–18.
- 123 Chou C-L, Tseng C-J, Shiu Y-L. The impact of young age on the prognosis for colorectal cancer: a population-based study in Taiwan. *Jpn J Clin Oncol* 2017;47:1010–8.
- 124 Lieu CH, Renfro LA, de Gramont A, *et al.* Association of age with survival in patients with metastatic colorectal cancer: analysis from the ARCAD clinical trials program. *J Clin Oncol* 2014;32:2975–82.
- 125 Blanke CD, Bot BM, Thomas DM, *et al.* Impact of young age on treatment efficacy and safety in advanced colorectal cancer: a pooled analysis of patients from nine first-line phase III chemotherapy trials. *J Clin Oncol* 2011;29:2781–6.
- 126 Kansakar P, Singh Y. Changing trends of colorectal carcinoma in Nepalese young adults. *Asian Pac J Cancer Prev* 2012;13:3209–12.
- 127 Fu J, Yang J, Tan Y. Young patients (≤35years old) with colorectal cancer have worse outcomes due to more advanced disease: a 30-year retrospective review. *Med* 2014;93:1–7.
- 128 Kocián P, Svobodová I, Krejčí D, *et al.* Is colorectal cancer a more aggressive disease in young patients? a population-based study from the Czech Republic. *Cancer Epidemiol* 2019;63:101621.
- 129 Dana Kharisma D. Healthcare Access Inequity within a Social Health Insurance Setting: A Risk Faced by Indonesia's Jaminan Kesehatan Nasional (JKN) Program. *Bwp* 2020;3:63–74.
- 130 Byeon J-S, Yang S-K, Kim TI, *et al.* Colorectal neoplasm in asymptomatic Asians: a prospective multinational multicenter colonoscopy survey. *Gastrointest Endosc* 2007;65:1015–22.
- 131 You YN, Xing Y, Feig BW, *et al.* Young-Onset colorectal cancer: is it time to pay attention? *Arch Intern Med* 2012;172:287–9.
- 132 Lee PY, Fletcher WS, Sullivan ES, *et al.* Colorectal cancer in young patients: characteristics and outcome. *Am Surg* 1994;60:607–12.
- 133 Domergue J, Ismail M, Astre C, *et al.* Colorectal carcinoma in patients younger than 40 years of age. Montpellier cancer Institute experience with 78 patients. *Cancer* 1988;61:835–40.
- 134 Ahnen DJ, Wade SW, Jones WF, *et al.* The increasing incidence of young-onset colorectal cancer: a call to action. *Mayo Clin Proc* 2014;89:216–24.
- 135 Gheju A, Jurescu A, Tăban S, *et al.* Different disease characteristics in young patients with colorectal cancer: a large retrospective study in a City in Romania. *J Int Med Res* 2021;49:030006052110166.
- 136 Chew M-H, Yeo S-AE, Ng Z-P, *et al.* Critical analysis of mucin and signet ring cell as prognostic factors in an Asian population of 2,764 sporadic colorectal cancers. *Int J Colorectal Dis* 2010;25:1221–9.
- 137 Börger ME, Gosens MJEM, Jeuken JWM, *et al.* Signet ring cell differentiation in mucinous colorectal carcinoma. *J Pathol* 2007;212:278–86.

- 138 Sudarshan V, Hussain N, Gahine R, *et al.* Colorectal cancer in young adults in a tertiary care hospital in Chhattisgarh, Raipur. *Indian J Cancer* 2013;50:337–40.
- 139 Elferink MAG, Siesling S, Visser O, *et al.* Large variation between hospitals and pathology laboratories in lymph node evaluation in colon cancer and its impact on survival, a nationwide population-based study in the Netherlands. *Ann Oncol* 2011;22:110–7.
- 140 Dubez A, Solymosi N, Schweigert M, *et al.* Time trends and disparities in lymphadenectomy for gastrointestinal cancer in the United States: a population-based analysis of 326,243 patients. *J Gastrointest Surg* 2013;17:611–9.
- 141 Shen SS, Haupt BX, Ro JY, *et al.* Number of lymph nodes examined and associated clinicopathologic factors in colorectal carcinoma. *Arch Pathol Lab Med* 2009;133:781–6.
- 142 Zhang M-R, Xie T-H, Chi J-L, *et al.* Prognostic role of the lymph node ratio in node positive colorectal cancer: a meta-analysis. *Oncotarget* 2016;7:72898–907.
- 143 Elsamany SA, Alzahrani AS, Mohamed MM, *et al.* Clinico-Pathological patterns and survival outcome of colorectal cancer in young patients: Western Saudi Arabia experience. *Asian Pac J Cancer Prev* 2014;15:5239–43.
- 144 Bentzen SM, Balslev I, Pedersen M, *et al.* Time to loco-regional recurrence after resection of Dukes' B and C colorectal cancer with or without adjuvant postoperative radiotherapy. A multivariate regression analysis. *Br J Cancer* 1992;65:102–7.
- 145 Compton CC, Fielding LP, Burgart LJ, *et al.* Prognostic factors in colorectal cancer. College of American pathologists consensus statement 1999. *Arch Pathol Lab Med* 2000;124:979–94.
- 146 Compton C, Fenoglio-Preiser CM, Pettigrew N, *et al.* American joint Committee on cancer prognostic factors consensus conference: colorectal Working group. *Cancer* 2000;88:1739–57.
- 147 Ruiz-Tovar J, Jiménez-Miramón J, Valle A, *et al.* Endoscopic resection as unique treatment for early colorectal cancer. *Rev Esp Enferm Dig* 2010;102:435–41.
- 148 Wu Z-Y, Wan J, Li J-H, *et al.* Prognostic value of lateral lymph node metastasis for advanced low rectal cancer. *World J Gastroenterol* 2007;13:6048–52.
- 149 Chen C-H, Hsieh M-C, Hsiao P-K, *et al.* A critical reappraisal for the value of tumor size as a prognostic variable in rectal adenocarcinoma. *J Cancer* 2017;8:1927–34.
- 150 Shida D, Ahiko Y, Tanabe T, *et al.* Shorter survival in adolescent and young adult patients, compared to adult patients, with stage IV colorectal cancer in Japan. *BMC Cancer* 2018;18:1–8.
- 151 Chaurasia AR. *COVID-19 Trend and Forecast in India: A Joinpoint Regression Analysis*. medRxiv [Internet, 2020. <http://medrxiv.org/content/early/2020/06/03/2020.05.26.20113399.abstract>
- 152 Ajbar S, Asif M, Ajbar AM. Did domestic travel restrictions slow down the COVID-19 pandemic in Saudi Arabia? A joinpoint regression analysis. *J Glob Health Rep* 2021;5:1–9.
- 153 US, Health Dof, Services H. National Institute of Health, National Cancer Institute. Methodology for Characterizing Trends [Internet]. *Online Summary of Trends in US Cancer Control Measures 2022* <https://progressreport.cancer.gov/methodology>
- 154 Guimarães RM, Magalhães MdeAFM, Xavier DR, *et al.* Is it time to talk about the end of social distancing? A joinpoint analysis of COVID-19 time series in Brazilian capitals. *Rev Soc Bras Med Trop* 2020;53:1–7.
- 155 Dragomirescu I, Llorca J, Gómez-Acebo I, *et al.* A join point regression analysis of trends in mortality due to osteoporosis in Spain. *Sci Rep* 2019;9:4264.
- 156 AIHW. Cancer mortality trends and projections: 2014 to 2025 [Internet]. Australian Institute of Health and Welfare, 2017. Available: <https://www.aihw.gov.au/reports/cancer/cancer-mortality-trends-and-projections-2014-to-2025/data> [Accessed 1 Aug 2022].
- 157 González-Padilla DA, España-Navarro R, Subiela JD, *et al.* Is “Movember” an Effective Prostate Cancer Awareness Campaign Beyond the English Language? Insights From Google Trends Among Spanish Speakers. *SIUJ* 2021;2:362–9.