

# BMJ Open Process of pain assessment in people with dementia living in nursing homes: a scoping review protocol

Caroline Kreppen Overen <sup>1,2</sup> Maria Larsson <sup>1</sup>  
Adelheid Hummelvoll Hillestad <sup>2</sup> Siren Eriksen <sup>2,3</sup>

**To cite:** Overen CK, Larsson M, Hillestad AH, *et al.* Process of pain assessment in people with dementia living in nursing homes: a scoping review protocol. *BMJ Open* 2022;**12**:e063230. doi:10.1136/bmjopen-2022-063230

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-063230>).

Received 25 March 2022

Accepted 12 September 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Institute of Health Sciences, Department of Nursing, Karlstad University, Faculty of Health, Science and Technology, Karlstad, Sweden

<sup>2</sup>Bachelor Education in Nursing, Lovisenberg Diaconal University College, Oslo, Norway

<sup>3</sup>Norwegian National Centre for Ageing and Health, Tønsberg, Norway

## Correspondence to

Caroline Kreppen Overen;  
caroline.kreppen.overen@ldh.no

## ABSTRACT

**Introduction** Pain is a common symptom in people with dementia; untreated, it reduces quality of life and causes suffering. People with dementia living in nursing homes most often have dementia in moderate to severe stages. The cognitive impairment, including language and communication difficulties, challenges pain assessment. Since pain is a subjective experience, self-reporting is the gold standard of assessment methods. Healthcare professionals are advised to help people with dementia communicate about their pain. The proposed scoping review is the first step in the development of a systematic pain assessment model for people with dementia living in nursing homes. The scoping review aims to identify, categorise and summarise knowledge on how pain assessment processes in this population are described in the literature, with a special focus on self-reporting.

**Methods and analysis** The scoping review will be conducted following the six-stage framework developed by Arksey and O'Malley, in addition to recent methodological developments. Systematic searches in CINAHL, Embase, Medline and PsycInfo will be conducted. The protocol follows the Preferred Reporting Items for Systematic Reviews and Meta-Analysis Protocols (PRISMA-P) and Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) checklists, and the scoping review will adhere to the PRISMA-ScR checklist. The review will include research that concerns assessment of pain in people with dementia living in nursing homes. Studies will be evaluated for quality and ethical standards. The analysis process will follow Bradbury-Jones *et al*'s PAGER framework. Patterns will be formed using thematic analysis. An overview of advances, gaps, evidence for practice and research recommendations associated with each pattern will be prepared. The research questions and results will be presented to and discussed in a reference group comprising nursing home residents, relatives, healthcare professionals and nursing home managers.

**Ethics and dissemination** The scoping review aims to collect and summarise data from available publications and does not require ethical approval. The final manuscript will be submitted to a peer-reviewed, open-access journal.

**Registration in open science framework** <https://osf.io/8kaf5/>

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This review will use an established scoping review methodology and standardised reporting guidelines.
- ⇒ To minimise the risk of personal biases, two reviewers will independently assess the studies for inclusion or exclusion; if disagreement arises, an additional reviewer will be consulted.
- ⇒ The included studies will be assessed for quality and ethical standards.
- ⇒ The review may miss relevant literature, as it will not include grey literature nor studies not published in English/non-Nordic languages.

## INTRODUCTION

In 2016, dementia was the fifth leading cause of death worldwide,<sup>1</sup> and the palliative perspective is important throughout the whole dementia trajectory.<sup>2</sup> As most people with dementia live their final days in a nursing home or similar,<sup>3 4</sup> healthcare professionals play an essential role in offering quality palliative care in this context. The prevalence of dementia in nursing home residents worldwide differs by location, nation and region.<sup>5</sup> In Norway as many as 80% of nursing home residents have dementia, and the majority has dementia in moderate to severe stages.<sup>6</sup> Moderate to severe stages of dementia have been described as an extended and intensive palliative care phase, often characterised by a loss of independence and autonomy, and reduction in physical and cognitive functions.<sup>7</sup> The trajectory is often unpredictable and palliative care initiation should therefore reflect need, not prognosis.<sup>8</sup> A five-round Delphi study resulted in 57 consensus-based recommendations for optimal palliative care in dementia, of which eight are clinical.<sup>2</sup> One of these clinical domains is symptom relief, considered one of the main aspects of palliative care.<sup>9</sup>

Pain is a common symptom among people with dementia living in nursing homes.<sup>10–13</sup> In a recent study, van Dam *et al* found that



43.3% of participants with dementia had clinically relevant pain scores.<sup>10</sup> Helvik *et al* state that 35.5% of their participants had clinically relevant pain on admittance to a nursing home.<sup>11</sup> A review conducted by Corbett *et al* indicates that 50% of people with dementia regularly experience pain and that the prevalence of pain in nursing home patients might be higher.<sup>12</sup> Pain in this patient group is often related to musculoskeletal, gastrointestinal and cardiac conditions, genitourinary infections, and wounds.<sup>12</sup> Discomfort caused by pain in people with dementia can be expressed as behavioural and psychological symptoms (BPSD), such as agitation, apathy, restlessness or wandering.<sup>2 14</sup> In Norway, the national clinical guidelines for dementia recommend that people with BPSD or other signs of discomfort should be assessed for pain as part of palliative care.<sup>15</sup> Pain assessment is frequently compromised by cognitive impairment,<sup>16</sup> including aspects of language and communication difficulties<sup>17</sup> in nursing home population.

Pain can be defined as *an unpleasant sensory and emotional experience associated with, or resembling that associated with, actual or potential tissue damage*.<sup>18</sup> Pain is a symptom, which is a subjective experience, as opposed to signs, which can be observed.<sup>19 20</sup> These definitions imply that self-reported information is the most appropriate for assessing pain. This represents a challenge in the target population, who may have difficulty communicating their symptoms because of reduced cognitive function<sup>7 21</sup>—their pain may therefore go unrecognised and unmanaged.<sup>13</sup> One recommendation is the systematic use of standardised observational tools and skills to chart pain, which can compensate for patients' lack of verbal communication.<sup>2 15 22 23</sup> However, nurses often rely on experience-based knowledge when interpreting signs of pain, and less-experienced nurses may fail to recognise pain in people with dementia.<sup>24</sup> Moreover, Pautex *et al* argue that the routine use of observational scales in severe dementia may not be justified and that self-assessment can be reliably performed among this population.<sup>25</sup> Achterberg *et al* highlight how self-reporting can also be adapted to individual capabilities during the course of dementia.<sup>13</sup> They recommend an initial use of simple numerical or verbal scales and the later use of 'yes' or 'no' questions; when cognitive and linguistic impairments reach a certain level, an observational tool can be added to the self-report to strengthen the validity of the pain assessment.

Pain management requires continuous mapping, assessment and treatment evaluation.<sup>19</sup> When caring for people with dementia this is complex and challenging.<sup>26–28</sup> It relies on healthcare professionals' knowledge of individuals' normal level of functioning and communication methods.<sup>26 29</sup> Healthcare professionals providing individualised care in nursing homes may be in a unique position to support and help people with dementia communicate their subjective

experience of pain, if they have knowledge of and frequent contact with the residents.<sup>26 29</sup>

Healthcare professionals in nursing homes need tools to systematically manage pain in people with dementia, which consider individual variation in pain expressions and ability to self-report. The proposed scoping review is the first step in developing a care model for systematic pain assessment in people with dementia living in nursing homes, which also includes how healthcare professionals recognise pain and evaluate initiated measures. The development of this model is rooted in the initial steps of The UK Medical Research Council (MRC) framework for developing and evaluating complex interventions.<sup>30</sup> To promote sustainable research and reduce research waste, it is important to obtain a preliminary overview of the field of research.<sup>31</sup> To the best of our knowledge, no study has reviewed the literature on self-reporting in pain assessment processes in people with dementia living in nursing homes, and how healthcare professionals can integrate self-reporting in recognising, assessing and evaluating pain in this group and context. The aim of this scoping review is therefore to identify, categorise and summarise knowledge about these processes from the literature.

## METHODS AND ANALYSIS

The proposed scoping review will follow Arksey and O'Malley's six-stage methodological framework<sup>32</sup> and Levac *et al*'s recommendations for each stage.<sup>33</sup> This will facilitate examination of the research concerning pain assessment in the target population and the identification of knowledge gaps. The Preferred Reporting Items for Systematic Reviews and Meta-Analysis Protocols (PRISMA-P)<sup>34</sup> and the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) checklists<sup>35</sup> were used to prepare this protocol (online supplemental files 1; 2). PRISMA-ScR<sup>35</sup> will be used in the review. The scoping review will be carried out in the period March 2022–September 2023.

### Stage 1: identify the research question(s)

Levac *et al*<sup>33</sup> highlight that research questions as comprehensive and broad as those recommended by Arksey and O'Malley<sup>32</sup> may lack the direction, clarity and focus needed to inform later steps in the research process. The concept and target population have therefore been defined to clarify the scoping review's focus and establish an effective search strategy, combined with a clear objective.<sup>33</sup> A population, concept, context (PCC) framework has informed the research questions (table 1) and will guide the database searches and eligibility criteria.

The following preliminary research questions were developed:

**Table 1** Population, concept, context framework informing research questions and search strategy

Criteria	Determinants
Population	People with dementia
Concept	Pain assessment processes
Context	Nursing home

- ▶ How do healthcare professionals recognise and assess pain in people with dementia living in nursing homes?
- ▶ How are the assessment processes of self-reported pain in the target group described?

In accordance with Arksey and O'Malley, these may be adjusted as the review progresses.

### Stage 2: identify relevant studies

The research questions and key concepts will inform the search strategy. The CINAHL, Embase, Medline and PsycInfo databases will be searched to identify relevant studies. The databases have been selected to cover a comprehensive range of healthcare research. A search strategy will be developed for each database with the assistance of an experienced librarian; these strategies will include medical subject headings (MESH), and search terms and synonyms combined using Boolean operators. The search strategy will consist of three main blocks informed by the PCC framework (table 1): people with dementia (population), pain assessment processes (concept of interest) and nursing home (context). The different search terms in each block will be combined with OR, and the blocks will be combined with AND. The reference lists of included studies will be manually searched. In line with Arksey and O'Malley, the search process will be iterative, and search terms may be adapted as the research team gains familiarity with the literature.<sup>32</sup> A pilot search will be conducted, where the first ~80 references will be reviewed; the search strategy will be

adjusted if needed. A preliminary search was conducted on 11/2/22 (online supplemental file 3).

### Stage 3: select studies


Following Arksey and O'Malley, the scoping review will identify all relevant literature regardless of study design, to obtain a broad picture of the existing research on the chosen topic.<sup>32</sup> Similarly, no time limit for publication will be specified. The inclusion and exclusion criteria are presented below (table 2)—these may be revised as the study progresses,<sup>32</sup> and any revised criteria will be applied to all citations. The selection process will be documented in a PRISMA-flowchart<sup>36</sup> (figure 1), including reasons for exclusion.<sup>34</sup> Duplicates will be removed using endnote and the duplicates not detected by endnote will be removed manually as the abstracts are reviewed. If the relevance of a study is unclear from the title and abstract, the full article will be reviewed. Traditionally, scoping reviews do not include secondary research, such as literature reviews. However, literature reviews will be included; as interventions targeting pain management in people with dementia may have been developed based on literature reviews, it would be inappropriate to exclude articles that could help answer the research questions.

Study selection will begin with a review of the title and abstract. If these correspond to the research questions and aim, a full-text review will be conducted. The studies will be reviewed by at least two researchers; in line with Levac *et al*, the research team will meet to discuss study inclusion and exclusion decisions in the beginning, middle and final stages of the abstract review process, and refine the search strategy as needed.<sup>33</sup> At least two reviewers will independently review full-text articles for inclusion; if disagreement arises, an additional reviewer will be consulted to determine final inclusion.

Levac *et al* argue that identifying gaps in the existing literature without assessing the quality of the included studies may lead to false conclusions about the nature

**Table 2** Preliminary eligibility criteria guiding study selection

	Eligibility criteria	
	Inclusion criteria	Exclusion criteria
Source	Peer-reviewed journals Published in English, Norwegian, Swedish or Danish	Grey literature
Population	People with dementia (eg, patients, service users or residents)	Mixed samples (eg, mild cognitive impairment/cognitive impairment + dementia) Dementia in people with Downs syndrome Cognitive impairment not caused by dementia
Context	Nursing home	
Concept	Literature that describes: How healthcare professionals (including nurses, nurse assistants and doctors) recognise, assess and evaluate pain in the population Whether and/or how self-reporting of pain is integrated in pain assessment processes in the population How healthcare professionals can support people with dementia in self-reporting of pain	
Study design	All study designs	Editorials, commentaries or letters, discussion papers, opinion papers and non-empirical studies



**Figure 1** Overview of study selection process using PRISMA flow diagram.<sup>36</sup> \*Consider, if feasible to do so, reporting the number of records identified from each database or register searched (rather than the total number across all databases/registers). \*\*If automation tools were used, indicate how many records were excluded by a human and how many were excluded by automation tools.

and extent of those gaps; they also assert that quality assessment of the included studies will increase the likelihood that findings will be useful for practice.<sup>33</sup> Study quality will therefore be assessed using appropriate appraisal tools, for example, the Critical Appraisal Skills Programme (CASP)<sup>37</sup> and Mixed Methods Appraisal Tool (MMAT).<sup>38</sup> An additional researcher will be included in the decision making if disagreement or uncertainty arises around the quality assessment.

#### Stage 4: charting data

This stage involves ‘charting’ key items of information obtained from the included studies by sorting material according to relevant issues and themes. In the proposed scoping review, this will be a mixture of general and specific information relating to study design and relevant findings. The charting process is also considered an iterative process, which means that the researchers may continuously update the data-charting form. In line with Levac *et al*, two researchers will independently extract data from the first five studies using the data-charting form and determine together whether the approach is consistent with the research questions and aim.<sup>33</sup> A preliminary data-charting form has been developed based on Arksey

#### Box 1 Data-charting form

- ⇒ Author, date and country
- ⇒ Study title
- ⇒ Aim, objective and/or research questions
- ⇒ Ethical assessment (financial support, conflicts of interest, informed consent, research committee approval, data protection)
- ⇒ Study context
- ⇒ Participant characteristics
- ⇒ Sampling method
- ⇒ Design and methods
- ⇒ Relevant findings

and O’Malley’s template (Box 1).<sup>32</sup> Ethical mapping is included in the data-charting form, in response to Weingarten *et al*’s emphasis on increasing ethical awareness in reviews.<sup>39</sup> Articles that do not adhere to ethical standards will be excluded.

#### Stage 5: collate, summarise and report results

As Arksey and O’Malley point out, unlike systematic reviews, scoping reviews do not synthesise evidence but instead provide an overview of the reviewed material.<sup>32</sup> In this stage, an overview and summary of the extracted information will therefore be prepared and presented, following the PAGER framework,<sup>40</sup> which consists of five categories: patterns, advances, gaps, evidence for practice and research recommendations. Patterns, or key themes, will be formed by using thematic analysis<sup>41</sup> of key findings from each study included in the review. We will then create an overview of advances, gaps, evidence for practice and research recommendations associated with each pattern.<sup>40</sup> The advances, gaps and research recommendations will guide further research needed to develop the pain assessment model, and the evidence for practice will guide the content of the model. Throughout the process, there will be regular meetings of the research group to discuss and agree on aspects of the analytical process and how the findings should best be presented.

#### Stage 6: consult with reference group

This scoping review is the first step in developing a care model for systematic pain assessment in people with dementia living in nursing homes. Correspondingly, a reference group will be formed, consisting of nursing home residents, relatives, healthcare professionals and nursing home managers. Arksey and O’Malley recommend consulting with practitioners and consumers to validate findings and make the research more useful for practice.<sup>32</sup> The findings will therefore be presented to and discussed with the reference group. In addition, the research team is part of a larger group of researchers, with whom the findings will also be discussed.

#### Patient and public involvement

The proposed scoping review’s research questions and aim will be presented to and discussed with the reference group, as will the findings. These latter will support the

development of an intervention promoting systematic pain management for people with dementia living in nursing homes.

### Ethics and dissemination

As the scoping review will not involve the collection of primary empirical data, ethical approval is unnecessary.<sup>42</sup> However, following Weingarten *et al*,<sup>39</sup> who state that ethical assessments of included studies should be conducted, ethical considerations are included in the data-charting process. Studies that do not adhere to ethical standards will be excluded. Findings from the scoping review will be published in an open-access, peer-reviewed journal. The scoping review is an important step in developing a pain assessment model for people with dementia living in nursing homes. Findings will enable the identification of existing models or interventions that may be further developed and tailored to the nursing home context, preventing research waste.<sup>31</sup>

**Acknowledgements** We would like to thank Annelie Ekberg-Andersson and Linda Borg—Information Specialists at Karlstad University—for their contribution to the search strategy presented in this protocol.

**Contributors** CKO was responsible for the preliminary study design, conceptualised the review approach, and led the writing of this protocol. ML, AHH and SE contributed to the protocol's development and approved the final version. SE led the supervision of the protocol's preparation.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not applicable.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### ORCID iDs

Caroline Kreppen Overen <http://orcid.org/0000-0002-2270-4557>

María Larsson <http://orcid.org/0000-0003-0417-6161>

Adelheid Hummelvoll Hillestad <http://orcid.org/0000-0002-5780-1699>

Siren Eriksen <http://orcid.org/0000-0002-5541-0934>

### REFERENCES

- Nichols E, Szoek CE, Vollset SE, *et al*. Global, regional, and national burden of Alzheimer's disease and other dementias, 1990–2016: a systematic analysis for the global burden of disease study 2016. *Lancet Neurol* 2019;18:88–106.
- van der Steen JT, Radbruch L, Hertogh CPM, *et al*. White paper defining optimal palliative care in older people with dementia: a

- Delphi study and recommendations from the European association for palliative care. *Palliat Med* 2014;28:197–209.
- Houttekier D, Cohen J, Bilsen J, *et al*. Place of death of older persons with dementia. A study in five European countries. *J Am Geriatr Soc* 2010;58:751–6.
- Mitchell SL, Teno JM, Miller SC, *et al*. A national study of the location of death for older persons with dementia. *J Am Geriatr Soc* 2005;53:299–305.
- Seitz D, Purandare N, Conn D. Prevalence of psychiatric disorders among older adults in long-term care homes: a systematic review. *Int Psychogeriatr* 2010;22:1025–39.
- Helvik A-S, Engedal K, Benth Jüratè Šaltytė, *et al*. Prevalence and severity of dementia in nursing home residents. *Dement Geriatr Cogn Disord* 2015;40:166–77.
- Hanson E, Hellström A, Sandvide Åsa, *et al*. The extended palliative phase of dementia - An integrative literature review. *Dementia* 2019;18:108–34.
- Livingston G, Huntley J, Sommerlad A, *et al*. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet* 2020;396:413–46.
- Radbruch L, De Lima L, Knaut F, *et al*. Redefining palliative Care-A new consensus-based definition. *J Pain Symptom Manage* 2020;60:754–64.
- van Dam PH, Caljouw MAA, Slettebø DD, *et al*. Quality of life and pain medication use in persons with advanced dementia living in long-term care facilities. *J Am Med Dir Assoc* 2019;20:1432–7.
- Helvik A-S, Bergh S, altytė Benth J. Pain in nursing home residents with dementia and its association to quality of life. *Aging Ment Health* 2021;1–11.
- Corbett A, Husebo B, Malcangio M, *et al*. Assessment and treatment of pain in people with dementia. *Nat Rev Neurol* 2012;8:264–74.
- Achterberg W, Lautenbacher S, Husebo B, *et al*. Pain in dementia. *Pain Rep* 2020;5:e803.
- Tible OP, Riese F, Savaskan E, *et al*. Best practice in the management of behavioural and psychological symptoms of dementia. *Ther Adv Neurol Disord* 2017;10:297–309.
- Helsedirektoratet. Nasjonal faglig retningslinje for demens Oslo: Helsedirektoratet; 2017. Available: <https://www.helsedirektoratet.no/retningslinjer/demens#apiUrl> accessed 18.02.22.
- Sirsch E, Lukas A, Drebenstedt C. Guideline Workgroup (Schmerzassessment bei älteren Menschen in Der vollstationären Altenhilfe, AWMF registry 145-001). pain assessment for older persons in nursing home care: an evidence-based practice guideline. *J Am Med Dir Assoc* 2020;21:149–63.
- Woodward M. Aspects of communication in Alzheimer's disease: clinical features and treatment options. *Int Psychogeriatr* 2013;25:877–85.
- International Association for the Study of Pain. Pain terms and definitions, 2011. Available: <https://www.iasp-pain.org/resources/terminology/#pain> [Accessed 21 Mar 2022].
- Dodd M, Janson S, Facione N, *et al*. Advancing the science of symptom management. *J Adv Nurs* 2001;33:668–76.
- Lenz ER, Pugh LC, Milligan RA, *et al*. The middle-range theory of unpleasant symptoms: an update. *ANS Adv Nurs Sci* 1997;19:14–27.
- Ellis-Smith C, Evans CJ, Bone AE, *et al*. Measures to assess commonly experienced symptoms for people with dementia in long-term care settings: a systematic review. *BMC Med* 2016;14:1–12.
- Husebo BS, Ostelo R, Strand LI. The MOBID-2 pain scale: reliability and responsiveness to pain in patients with dementia. *Eur J Pain* 2014;18:1419–30.
- Torvik K, Kaasa S, Kirkevoll Øyvind, *et al*. Validation of Doloplus-2 among nonverbal nursing home patients - an evaluation of Doloplus-2 in a clinical setting. *BMC Geriatr* 2010;10:1–9.
- Dowding D, Lichtner V, Alcock N, *et al*. Using sense-making theory to aid understanding of the recognition, assessment and management of pain in patients with dementia in acute hospital settings. *Int J Nurs Stud* 2016;53:152–62.
- Pautex S, Michon A, Guedira M, *et al*. Pain in severe dementia: self-assessment or observational scales? *J Am Geriatr Soc* 2006;54:1040–5.
- De Witt Jansen B, Brazil K, Passmore P. Exploring healthcare assistants' role and experience in pain assessment and management for people with advanced dementia towards the end of life: a qualitative study. *BMC Palliat Care* 2017;16:1–11.
- De Witt Jansen B, Brazil K, Passmore P, *et al*. Nurses' experiences of pain management for people with advanced dementia approaching the end of life: a qualitative study. *J Clin Nurs* 2017;26:1234–44.
- De Witt Jansen B, Brazil K, Passmore P, *et al*. 'A tool doesn't add anything'. The importance of added value: Use of observational pain



- tools with patients with advanced dementia approaching the end of life—a qualitative study of physician and nurse experiences and perspectives. *Int J Geriatr Psychiatry* 2018;33:1346–54.
- 29 Monroe TB, Parish A, Mion LC. Decision factors nurses use to assess pain in nursing home residents with dementia. *Arch Psychiatr Nurs* 2015;29:316–20.
- 30 Skivington K, Matthews L, Simpson SA, *et al.* Framework for the development and evaluation of complex interventions: gap analysis, workshop and consultation-informed update. *Health Technol Assess* 2021;25:1–132.
- 31 Chalmers I, Glasziou P. Avoidable waste in the production and reporting of research evidence. *The Lancet* 2009;374:86–9.
- 32 Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 2005;8:19–32.
- 33 Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implementation Science* 2010;5:1–9.
- 34 Moher D, Shamseer L, Clarke M, *et al.* Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1–9.
- 35 Tricco AC, Lillie E, Zarin W, *et al.* PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018;169:467–73.
- 36 Page MJ, McKenzie JE, Bossuyt PM, *et al.* The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Int J Surg* 2021;88:105906.
- 37 CASP Checklist. Casp critical appraisal skills programme, 2018. Available: <https://casp-uk.b-cdn.net/wp-content/uploads/2018/01/CASP-Qualitative-Checklist-2018.pdf> accessed 25.08.21.
- 38 Hong QN, Fàbregues S, Bartlett G, *et al.* The mixed methods appraisal tool (MMAT) version 2018 for information professionals and researchers. *Education for Information* 2018;34:285–91.
- 39 Weingarten MA, Paul M, Leibovici L. Assessing ethics of trials in systematic reviews. *BMJ* 2004;328:1013–4.
- 40 Bradbury-Jones C, Aveyard H, Herber OR. Scoping reviews: the PAGER framework for improving the quality of reporting. *International Journal of Social Research Methodology* 2021:1–14.
- 41 Braun V, Clarke V, Braun V. *Thematic analysis : a practical guide*. Los Angeles, California: SAGE, 2022.
- 42 Helseforskningsloven. *Lov Om medisinsk OG helsefaglig forskning: Helse- OG omsorgsdepartementet*, 2008.