

# BMJ Open Factors associated with higher quality of clinical practice guidelines and their recommendations for the pharmacological treatment of depression: a systematic review

Franciele Cordeiro Gabriel <sup>1</sup>, Daniela Oliveira de Melo <sup>2</sup>,  
Airtón Tetelbom Stein <sup>3,4</sup>, Géssica Caroline Henrique Fontes-Mota <sup>1</sup>,  
Itamires Benício dos Santos <sup>2</sup>, Camila da Silva Rodrigues <sup>2</sup>,  
Mônica Cristiane Rodrigues <sup>5</sup>, Tatiane Bomfim Ribeiro <sup>6</sup>,  
Renério Fráguas <sup>7</sup>, Ivan D Florez <sup>8,9,10</sup>, Diogo Telles-Correia <sup>11</sup>,  
Eliane Ribeiro <sup>1</sup>

**To cite:** Gabriel FC, de Melo DO, Stein AT, *et al.* Factors associated with higher quality of clinical practice guidelines and their recommendations for the pharmacological treatment of depression: a systematic review. *BMJ Open* 2023;**13**:e067390. doi:10.1136/bmjopen-2022-067390

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-067390>).

Received 11 August 2022  
Accepted 11 March 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Franciele Cordeiro Gabriel;  
[francordegabriel@gmail.com](mailto:francordegabriel@gmail.com)

## ABSTRACT

**Objective** The objective of this study was to assess the quality of clinical practice guidelines (CPGs) for the pharmacological treatment of depression along with their recommendations and factors associated with higher quality.

**Design** We conducted a systematic review that included CPGs for the pharmacological treatment of depression in adults.

**Data sources** We searched for publications from 1 January 2011 to 31 December 2021, in MEDLINE, Cochrane Library, Embase, PsycINFO, BVS and 12 other databases and guideline repositories.

**Eligibility criteria for selecting studies** We included CPGs containing recommendations for the pharmacological treatment of depression in adults at outpatient care setting, regardless of whether it met the U.S. National Academy of Medicine criteria, or not. If a CPG included recommendations for both children and adults, they were considered. No language restriction was applied.

**Data extraction and synthesis** Data extraction was also conducted independently and in duplicate, a process that was validated in a previous project. The quality of the CPGs and their recommendations were assessed by three independent reviewers using Appraisal of Guidelines for Research and Evaluation (AGREE II) and Appraisal of Guidelines for Research and Evaluation-Recommendations Excellence (AGREE-REX). A CPG was considered to be of high quality if AGREE II Domain 3 was ≥60%; while their recommendations were considered high if AGREE-REX Domain 1 was ≥60%.

**Results** Seventeen out of 63 (27%) CPGs were classified as high quality, while 7 (11.1%) had high-quality recommendations. The factors associated with higher-scoring CPGs and recommendations in the multiple linear regression analyses were 'Handling of conflicts of interest', 'Multiprofessional team' and 'Type of institution'. 'Inclusion of patient representative in the team' was also associated with higher-quality recommendations.

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ To increase the reliability of the quality assessment of the guidelines and recommendations, three reviewers independently conducted the assessment using both AGREE II and AGREE-REX evaluation tools. Before the assessment, the appraisers underwent rigorous training to ensure consistency in their evaluations.
- ⇒ The study was based on a comprehensive literature search on the pharmacological treatment of depression conducted in 17 databases using a sensitive strategy.
- ⇒ The inclusion of studies published in different languages made it difficult to include all documents present on the websites of specific institutions.

**Conclusions** The involvement of professionals from diverse backgrounds, the handling of conflicts of interest, and the inclusion of patients' perspectives should be prioritised by developers aiming for high-quality CPGs for the treatment of depression.

## INTRODUCTION

Depression is a serious mental health problem that causes severe professional, economic, social and personal incapacitation.<sup>1 2</sup> The Global Burden of Disease Study 2017 estimated that more than 264 million people worldwide were affected by depression.<sup>3</sup> Moreover, according to the WHO, approximately 800 000 cases of suicide per year are attributable to this disease.<sup>4</sup> The prevalence of depression has increased considerably in the last few years,<sup>5 6</sup> overloading healthcare systems.<sup>7</sup> The COVID-19 pandemic increased

the prevalence of mental illness even further, generating an extra need of resources to overcome the burden of mental health disorders.<sup>8</sup> While a growing number of depression cases are being recognised and treated, rates of poor outcomes at 1 year in community populations remain high.<sup>9</sup> The optimisation of health resources for the treatment of depression by implementing evidence-based health interventions is challenging,<sup>10</sup> but necessary. In this scenario, adherence to high-quality clinical practice guidelines (CPGs) is vital.

CPGs are documents that contain recommendations for the optimisation of patient care, developed through the systematic review of evidence and analysis of the risks, benefits and costs of interventions for each clinical health condition.<sup>11</sup> However, the potential benefits of a CPG depend on its quality. Only high-quality CPGs have the potential of facilitating expected positive outcomes in care of patients with depression, facilitating the clinical decision-making process, enhancing the education process of patients and professionals on the best practices, reducing unnecessary clinical variability, and improving the cost-effectiveness of healthcare.<sup>12</sup>


Nevertheless, there are some challenges associated with high-quality CPGs. The development of rigorous CPGs is a time-consuming and expensive task. High-quality CPGs need to be supported by systematic reviews which

require significant time, effort and technical capacity to complete.<sup>11</sup> Guidelines need to be developed by a multi-professional, independent team of experts that do not have competing interests. They also need to be continuously updated in response to new and relevant evidence, their development should be transparent and reproducible, and they need to consider patients' values and preferences.<sup>11 13</sup> Furthermore, the final documents of the guidelines need to be clear, well-organised and user-friendly.<sup>12 14</sup>

With the aim of providing support to guideline developers and users, some instruments have been developed to evaluate CPG quality. The most used tool for this purpose is the Appraisal of Guidelines for Research and Evaluation (AGREE II).<sup>13 15 16</sup> AGREE II has been validated and translated into several languages and provides online training and a clearly written user manual. This popular instrument enables a broad CPG quality assessment,<sup>17</sup> however it does not consider the quality of its recommendations.

To address this knowledge gap, the AGREE team developed an evidence-based AGREE II add-on, called the Appraisal of Guidelines for Research and Evaluation-Recommendations Excellence (AGREE-REX), which enables the critical assessment of the recommendations of CPGs by their developers and users.<sup>18–20</sup> It is important

#### Figure legend



Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *Br Med J*. 2021;372:n71. doi: 10.1136/bmj.n71. For more information, visit: <http://www.prisma-statement.org/>.

**Figure 1** Flowchart of clinical practice guidelines (CPG) selection.

**Table 1** Characteristics of the included clinical practice guidelines (CPGs)

Characteristic, no. (%)	n=63
Handling of conflicts of interest	11 (17.5)
Multiprofessional team	23 (36.5)
Inclusion of patient representative in the team	6 (9.5)
Governmental funding	24 (38.1)
Type of institution or organisation	
Independent researcher/University	19 (30.2)
Professional society	25 (39.7)
Governmental	19 (30.2)
Year of publication	
2011 to 2015	27 (42.9)
2016 to 2021	36 (57.1)

to note that AGREE-REX assesses the clinical credibility of the recommendations, meaning that it checks whether the recommendations have covered the key elements that make them more applicable to a particular context.<sup>17</sup> AGREE-REX considers that for a recommendation to be both high-quality and reliable, it must consider the values of the CPG developers and policy makers, in addition to patient preferences, and additional factors such as the clinical applicability and purpose of the CPG.<sup>18</sup> AGREE-REX complements the AGREE II evaluation by providing an evaluation of key elements that underpin the development of the recommendations. Therefore, for a complete evaluation of CPG quality, it is important to appraise both the general quality of the CPGs and its recommendations' quality.

**Table 2** Clinical practice guidelines (CPGs) descriptive statistics for AGREE II and AGREE-REX Scores, n=63

Instrument domain	Mean±SD	Median (IQR)
AGREE II		
Scope and purpose	62.4±19.4	63 (46–78)
Stakeholder involvement	40.9±23.4	39 (22–59)
Rigour of development	39.4±26.4	35 (14–63)
Clarity of presentation	68.4±18.5	70 (57–83)
Applicability	29.3±21.3	23 (13–39)
Editorial independence	50.0±24.5	53 (33–72)
AGREE-REX		
Clinical applicability	36.3±19.2	35 (19–50)
Values and preferences	17.7±14.0	14 (7–25)
Implementability	35.1±15.7	33 (25–44)
The CPGs included were published between 1 January 2011 and 31 January 2021.		
AGREE II, Appraisal of Guidelines for Research and Evaluation; AGREE-REX, Appraisal of Guidelines for Research and Evaluation-Recommendations Excellence; SD, standard deviation.		

Some studies have already assessed the quality of CPGs for the pharmacological treatment of depression using AGREE II. However, the main focus of these studies was not on the quality of the documents or the factors related to high-quality. Furthermore, patient characteristics were either more restricted or included other chronic conditions, which did not specifically address depression.<sup>21</sup> To our knowledge, no study has assessed the quality of recommendations for the pharmacological treatment of depression using AGREE-REX.

By identifying the factors associated with high-quality CPG recommendations for the pharmacological treatment of depression, we can evaluate the areas for improvement that can help developers enhance their processes and create superior quality CPGs and recommendations. Therefore, our study aimed to assess the quality of CPGs for the treatment of depression and their recommendations and identify the factors associated with higher quality.

## METHODS

The methods for this systematic review have already been previously reported in our published protocol<sup>22</sup> and are only briefly described here. Our methodology involved: (1) The identification of CPGs, (2) Extraction of CPGs' characteristics, (3) Appraisal of CPGs' quality using AGREE II, (4) Appraisal of the quality of CPG recommendations with AGREE-REX, and (5) Analysis of factors associated with the quality of CPGs and their recommendations.

The identification and appraisal of CPGs was initially conducted to elaborate a comparison of high-quality recommendations for the pharmacological treatment of depression. As a post hoc analysis, we then explored the factors associated with high-quality guidelines and recommendations.

We searched for publications from 1 January 2011 to 31 December 2021, in MEDLINE, Cochrane Library, Embase, PsycINFO, BVS (Virtual Health Library Regional Portal) and 12 other databases and guideline repositories (online supplemental material). The obtained citations were screened to select potentially eligible articles, which were obtained as full texts and reviewed to assess inclusion. Both previous steps were performed independently and in duplicate. Data extraction was also conducted independently and in duplicate, a process that was validated in a previous project.<sup>21–23</sup>

The quality of the CPGs was appraised by a team of multidisciplinary researchers trained according to a previously published protocol.<sup>22</sup> A final consolidated score ranging from 0% to 100% was obtained per AGREE II and AGREE-REX domains. More information on the methodology is provided in the published protocol.<sup>22</sup> Data were summarised by standard statistical methods and analysed using simple and multiple linear regression. Data were processed and analysed with IBM SPSS v.25.0. The detailed methodology is outlined in in online

**Table 3** Contribution of CPG characteristics to AGREE II Domain 3 and AGREE-REX Domain 1 Scores (n=63)

Characteristics	Univariate analysis			Multivariable analysis		
	B	95% CI	P value	B	95% CI	P value
AGREE II Domain 3						
Handling of conflicts of interest	43.2	29.2 to 57.2	<0.001	20.4	7.9 to 32.9	0.002
Multiprofessional team	41.1	31.7 to 50.4	<0.001	22.2	10.2 to 34.2	<0.001
Inclusion of patient representative in the team	36.8	15.7 to 57.8	0.001	3.7	−11.5 to 18.9	0.627
Governmental funding	20.2	7.3 to 33.1	0.003	0.3	−12.5 to 13.1	0.964
Type of institution						
Independent researcher/university	Ref.	–	–	Ref.	–	–
Professional society	31.1	18.5 to 43.7	<0.001	17.8	7.6 to 27.9	0.001
Governmental	42.2	28.7 to 55.6	<0.001	19.8	4.5 to 35.1	0.012
Publication year >2015	−3.3	−16.9 to 10.4	0.631	0.5	−7.6 to 8.7	0.898
AGREE-REX Domain 1						
Handling of conflicts of interest	30.0	19.7 to 40.3	<0.001	12.8	3.4 to 22.2	0.008
Multiprofessional team	27.0	19.6 to 34.5	<0.001	11.2	2.2 to 20.3	0.016
Inclusion of patient representative in the team	31.3	16.7 to 45.9	<0.001	11.1	−0.3 to 22.6	0.056
Governmental funding	11.9	2.4 to 21.5	0.016	1.0	−8.7 to 10.6	0.841
Type of institution						
Independent researcher/university	Ref.	–	–	Ref.	–	–
Professional society	25.4	16.4 to 34.4	<0.001	17.0	9.4 to 24.6	<0.001
Governmental	29.1	19.5 to 38.7	<0.001	15.2	3.7 to 26.7	0.010
Publication year >2015	−1.6	−11.5 to 8.2	0.740	1.0	−5.2 to 7.1	0.748

AGREE II, Appraisal of Guidelines for Research & Evaluation; Domain 3, Rigour of Development; AGREE-REX, AGREE Recommendations Excellence; Domain 1, Clinical Applicability; B, linear regression coefficient representing the variable absolute impact (ie, increase or decrease) on the score; CI, confidence interval; P value, statistical significance.

supplemental material 1, while the reasons for excluding CPGs are shown in online supplemental material 2.

### Patient and public involvement

No patient was involved.

## RESULTS

### CPG identification

We retrieved 5063 documents from the search and removed 419 duplicates; thus, we screened 4644 references. We discarded non-relevant references and retrieved 174 full texts to check their eligibility (two full texts could not be retrieved). A total of 126 documents were excluded and 48 documents were included after the full-text review. Moreover, we identified 15 documents from the guidelines' repositories. Ultimately, 63 CPGs were included in this study (see [figure 1](#)).

### Characteristics of CPGs

[Table 1](#) provides a summary of the characteristics of the included CPGs. Most were published after 2015 (36, 57.1%), 11 (17.5%) reported how conflicts of interest were handled and only 6 (9.5%) included patient representatives in the development team.

### Appraisal of the quality of the guidelines and their recommendations

[Table 2](#) presents descriptive statistics for the AGREE II and AGREE-REX Scores. Among the 63 CPGs, domains with low mean scores on the AGREE II ( $\leq 50\%$ ) were 'Stakeholder Involvement', 'Rigour of Development' and 'Applicability', while for the AGREE-REX, the mean scores in all domains were below 50%. The results of the appraisal of the CPGs' domains using the AGREE II and AGREE-REX are shown in [table 1](#) in the online supplemental material 3. [Table 3](#) presents the contribution of CPG characteristics to AGREE II Domain 3 and AGREE-REX Domain 1 scores.

Seventeen (27.0%) CPGs were classified as high quality according to AGREE II (Domain 3, Rigour of Development  $\geq 60\%$ ).<sup>24–40</sup> Moreover, seven (11.1%) CPGs were also considered to have high-quality recommendations (AGREE-REX Domain 1, Clinical Applicability  $\geq 60\%$ ).<sup>24–31</sup> All CPGs classified as being of high quality by AGREE-REX were also considered high quality according to AGREE II (online supplemental material).

### Factors associated with higher quality

In the univariate analyses, all factors except publication year >2015 were found to have statistically significant



associations with AGREE II Domain 3 and AGREE-REX Domain 1. For AGREE II Domain 3, the multiple linear regression analysis showed that 'Handling of conflicts of interest', 'Multiprofessional team' and 'Type of institution' were statistically significant factors associated with increased scores. Regarding 'Type of institution', both 'Professional society' and 'Governmental institutions' presented higher scores when compared with 'Independent researcher/University'. The remaining factors (ie, 'Inclusion of patient representative in the team', 'Governmental funding' and 'Publication yr.>2015') did not reach statistical significance.

When considering AGREE-REX Domain 1, the same three factors were deemed to have a significant association with higher scores in the multiple linear regression model: 'Handling of conflicts of interest', 'Multiprofessional team' and 'Type of institution'. Similarly, for AGREE-REX Domain 1, institutions represented by 'Independent researcher/University' were those found to have lower scores when compared with 'Professional society' and 'Governmental institutions'. However, it is important to note that 'Inclusion of patient representative in the team' reached borderline statistical significance ( $p=0.056$ ) and was associated with higher AGREE-REX Domain 1 scores.

## DISCUSSION

In this systematic review, we identified 63 CPGs for the treatment of depression in adults. We found that few guidelines were classified as high quality according to AGREE II (17/63, 27.0%) and even fewer were considered to have high-quality recommendations according to AGREE-REX (7/63, 11.1%). The factors associated with higher-scoring CPGs (AGREE II Domain 3) and recommendations (AGREE-REX Domain 1) in our multiple linear regression analyses were 'Handling of conflicts of interest', 'Multiprofessional team' and 'Type of institution', with 'Professional society' and 'Governmental institutions' presenting higher scores when compared with 'Independent researcher/University'. Additionally, 'Inclusion of patient representative in the team' was found to have borderline statistical significance ( $p=0.056$ ) with higher AGREE-REX Domain 1 Scores, which are related to improved clinical applicability.

In the multivariable analysis, 'Inclusion of patient representative in the team' was no longer significant in the model analysing AGREE II Domain 3 (Rigour of Development). This finding could be explained by the small number of CPGs with this particular characteristic declared, which may produce less reliable estimates and wide CIs in a more saturated model. In addition, 'Governmental funding' lost its statistical significance, probably because it may be a surrogate marker of the involvement of 'Governmental institutions'.

We identified several domains with suboptimal performance as evaluated with the AGREE II and AGREE-REX tools. With the AGREE II tool, the worst-scored domains

were 'Applicability', 'Rigour of development' and 'Stakeholder involvement'. These findings show that CPGs may not always be based on the best evidence. Many currently available CPGs do not improve the quality of care for patients and may lead to the waste of scarce resources.<sup>41–43</sup> In other words, low scores in AGREE II Domain 3 (Rigour of Development) may indicate that most pharmacological treatments for depression did not report or perform the methodological processes expected for high-quality guidelines. Therefore, these CPGs had low or no use of appropriate development methods, selection and synthesis of evidence, and recommendations.

With the AGREE-REX tool, all three domains—'Clinical applicability', 'Values and preferences' and 'Implementability'—received low scores. The domain with the worst quality was 'Values', signalling that CPGs did not address the preferences of professionals, policy makers, developers or patient representatives in their recommendations. The values of policy makers are frequently missing, although some CPGs mention the term 'equity'—one of the values and preferences—as an important concept and as a factor considered in its development. Low scores in Domain 2 (Values) have also been found when applying AGREE-REX in CPGs for different health conditions.<sup>20</sup> Considering that therapeutic failure and adverse events in depression treatment are not uncommon, considering the values of patients, professionals, developers and policy/decision makers might be central to ensuring the effectiveness of a CPG.

In AGREE-REX Domain 1 (Clinical Applicability), we identified that the CPGs failed to clearly report the analysis of the quality of the studies, develop a list of relevant treatment outcomes (eg, quality of life, symptomatic remission, response) and appoint a patient representative as a team member. Finally, regarding AGREE-REX Domain 3 (Implementability), CPGs did not mention the anticipated impact when implementing recommendations and in the formal analysis of costs beyond the definitions of audit criteria to verify such implementation. Finally, low scores in AGREE II Domain 5 (Applicability) and AGREE-REX Domain 3 (Implementability) reveal a lack of consideration to implement strategies for clinical practice, which may lead to an ineffective interpretation of the best available evidence in practice.

We found that the handling of conflicts of interest was relevant in determining a high-quality status for a CPG and its recommendations. Such results call attention to previous reports supporting that simply declaring conflicts of interest is not enough.<sup>44</sup> Declaring potential conflicts of interests is not sufficient to avoid bias in the CPG development. Handling these conflicts by removing participants with conflicts from specific discussions, from voting or from the guideline group is essential to maintain rigour and transparency in the development process. The mean score of the 'Editorial independence' domain in AGREE II, the domain that includes how to handle conflicts of interest, was 50.0, contrasting with the mean scores of the highest-scored domains: 'Clarity of presentation' with

68.4 and 'Scope and purpose' with 62.4. This indicates a need for increasing attention to improve 'Handling of conflicts of interest' when developing a CPG.

We also found that a 'Multiprofessional team' was associated with higher quality in CPGs and their recommendations. These results may support the implementation of multiprofessional teams working together and sharing different practices and knowledge may offer improved results for patients, organisations and health systems.<sup>45 46</sup>

According to our analyses, 'Inclusion of patient representative in the team' may also be important for the quality of recommendations. Considering patients unique views, preferences and values regarding treatment benefits and harms enriches CPGs, helps minimise disease stigmatisation and improves adherence to treatment.<sup>47-50</sup>

A study by Zafra-Tanaka *et al*<sup>51</sup> analysed the quality and characteristics of 11 CPGs for depression published from January 2014 to May 2018 using the AGREE II instrument. Their findings revealed that <50% of these CPGs (5/11) have shared their search strategies or listed the studies used to develop the recommendations (4/11).<sup>51</sup> These information gaps make it more difficult to understand the possible biases such as potential conflicts of interest in the formulation of the recommendations provided in CPGs, thereby undermining health professionals' trust in these guidelines. Only 18% of the CPGs (2/11) have included patient representatives in their development team. Another relevant finding of this study was that only 27% (3/11) of the CPGs had a score of  $\geq 70\%$  in AGREE II Domain 3 (methodological rigour).

In addition, in a systematic review conducted by our group, the Chronic Diseases and Informed Decisions Research Group (CHRONIDE),<sup>21</sup> the methodological rigour was evaluated using AGREE II for 421 CPGs for treating chronic non-communicable diseases (including depression) in primary healthcare, <25% of those CPGs (99/421) were considered as having high quality. Of all the CPGs evaluated in the study, 31 CPGs were for treating depression. Of these, only 45% (14/31) were considered high quality ( $\geq 60\%$  in AGREE II Domain 3). Remarkably, no associations between geographical region and quality of CPGs were found, unlike the findings of other studies.<sup>52 53</sup> Inclusion of >20 authors, being developed by government institutions and disclosure of financing support were associated with a higher quality based on logistic regression analysis. The authors raised a point that the government institutions have more financial and human resources for developing CPGs, which is generally a lengthy and costly process. Moreover, the study also revealed that Domain 4 (Applicability) received lower scores in the AGREE II assessment, either for the CPGs to treat depression or for the CPGs as a whole.

The findings reported by Zafra-Tanaka *et al*<sup>51</sup> and Molino *et al*<sup>21</sup> (the CHRONIDE Group) demonstrate that professionals and policy makers should know that only few CPGs for adults with depression demonstrate high developmental rigour. In addition, the relevance of including the patient's viewpoints, highlighted by Zafra-Tanaka *et*

*al*,<sup>51</sup> converges to low quality of applicability detected by Molino *et al*.<sup>21</sup> Thus, the findings indicate that healthcare policy makers should invest in improving the developmental rigour of CPGs to attain the confidence of professionals using them and, on the implementation, and monitoring of recommendations to ensure their applicability. Following these principles, the role of CPGs as a tool to promote evidence-based health is safeguarded.

This article has several strengths. First, we conducted a comprehensive search across 17 CPG databases and repositories. To ensure a comprehensive quality assessment of the guidelines and recommendations, three independent raters were trained rigorously and used both AGREE II and AGREE-REX evaluation tools. This contributed to increase the reliability of quality assessment. Moreover, the use of AGREE-REX is still incipient and at the time of writing fewer than 10 studies have used this tool for the appraisal of the quality of CPGs recommendations.

Nevertheless, this study also has some limitations. To better evaluate the guidelines and their recommendations, whenever possible, we reviewed their supplemental materials and the methodological guidance of their institutions. However, the search for these documents was conducted in the organisations' websites, and we may have missed additional relevant documents that may impact the final evaluations. Moreover, the inclusion of studies published in different languages made it difficult to include all documents present on the websites of specific institutions.

Another aspect is that although most included CPGs were published after the release of The Diagnostic and Statistical Manual of Mental Disorders V (DSM-V) in 2013, some of them may have relied on DSM-IV definitions. However, since the evaluation of CPG quality does not specifically involve diagnostic criteria, it is unlikely that this could influence the results observed in this study.

Finally, as there were very few CPGs classified as high quality, we obtained wide CIs for the associations, which creates uncertainty regarding the estimates. Therefore, caution is recommended when interpreting our results. However, this possible limitation does not lessen the relevance of our work since, at least to our knowledge, this is the first study to assess the quality of recommendations of CPGs for the treatment of depression and to explore the factors associated with higher-quality CPGs and their recommendations.

## CONCLUSION

We identified 63 CPGs for the pharmacological treatment of depression in adults, with 27.0% classified as high quality and 11.1% as having high-quality recommendations. The factors 'Handling of conflicts of interest', 'Multiprofessional team' and 'Type of institution' were significantly associated with higher quality in AGREE II Domain 3 and AGREE-REX Domain 1, followed by 'Inclusion of patient representative in the team', which may have an important role in AGREE-REX Domain 1.

CPG developers should be aware of the above characteristics to obtain more reliable and implementable recommendations. They should focus on improving quality as a whole and, more emphatically, on developing better recommendations rather than creating new ones with similar limitations.

#### Author affiliations

<sup>1</sup>Departamento de Farmácia, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, São Paulo, Brazil

<sup>2</sup>Departamento de Ciências Farmacêuticas, Universidade Federal de São Paulo, São Paulo, Brazil

<sup>3</sup>Departamento de Saúde Pública, Universidade Federal de Ciências da Saúde de Porto Alegre, Porto Alegre, Brazil

<sup>4</sup>Hospital Conceição, Porto Alegre, Brazil

<sup>5</sup>Faculdade de Saúde Pública, Universidade de São Paulo, São Paulo, Brazil

<sup>6</sup>Departamento de Epidemiologia, Escola de Saúde Pública, Universidade de São Paulo, São Paulo, Brazil

<sup>7</sup>Departamento e Instituto de Psiquiatria, Hospital das Clínicas, Faculdade de Medicina da Universidade de São Paulo (IPq-HC-FM-USP), Divisão de Psiquiatria e Psicologia Hospital Universitário (HU), Laboratório de Investigações Médicas – 21 (LIM 21) FM, Universidade de São Paulo, São Paulo, Brazil

<sup>8</sup>School of Rehabilitation Sciences, McMaster University, Hamilton, Ontario, Canada

<sup>9</sup>Department of Pediatrics, University of Antioquia, Medellín, Colombia

<sup>10</sup>Clínica Las Américas, AUNA, Medellín, Colombia

<sup>11</sup>Clínica Universitária de Psiquiatria e Psicologia Médica, Faculdade de Medicina da Universidade de Lisboa, Lisboa, Portugal

**Twitter** Ivan D Florez @ivand\_florez

**Acknowledgements** The authors thank Ana Paula Callejo and Beatriz Minguzzi for their invaluable assistance in helping them better understand the method of appraisal for CPGs.

**Contributors** FCG, DdM, ATS, IDF, DTC and ER conceived the main idea behind this manuscript. FCG, IDF, ATS and ER wrote the manuscript. GCHF-M, IBdS, CdSR, MCR, RF and TR critically revised the manuscript and made important intellectual contributions to its development. All authors read and approved the final version of the manuscript. FCG is responsible for the overall content as guarantor.

**Funding** This work was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPQ) (no. 141811/2020–0) and the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Programa de Pesquisa para o SUS-Gestão Compartilhada em Saúde (PPSUS) (no. 2019/03883-7).

**Competing interests** IDF is the current leader of the AGREE collaboration.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not applicable.

**Ethics approval** Not applicable.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All data relevant to the study are included in the article or uploaded as supplementary information.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Franciele Cordeiro Gabriel <http://orcid.org/0000-0002-4375-3729>

Daniela Oliveira de Melo <http://orcid.org/0000-0001-8613-7953>

Airton Tetelbom Stein <http://orcid.org/0000-0002-8756-8699>

Géssica Caroline Henrique Fontes-Mota <http://orcid.org/0000-0003-1986-9155>

Itamires Benício dos Santos <http://orcid.org/0000-0002-8693-3121>

Camila da Silva Rodrigues <http://orcid.org/0000-0001-7733-5149>

Mônica Cristiane Rodrigues <http://orcid.org/0009-0004-1743-0995>

Tatiane Bomfim Ribeiro <http://orcid.org/0000-0003-4434-748X>

Renério Fráguas <http://orcid.org/0000-0002-3052-066X>

Ivan D Florez <http://orcid.org/0000-0002-0751-8932>

Diogo Telles-Correia <http://orcid.org/0000-0003-4217-2823>

Eliane Ribeiro <http://orcid.org/0000-0003-0886-368X>

#### REFERENCES

- Chisholm D, Sweeny K, Sheehan P, et al. Scaling-up treatment of depression and anxiety: a global return on investment analysis. *Lancet Psychiatry* 2016;3:415–24.
- Thompson C. *Affective disorders: the instruments of psychiatric research*. London: John Wiley & Sons, 2010.
- GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the global burden of disease study 2017. *Lancet* 2018;392:1789–858.
- World Health Organization. *Depression and other common mental disorders: global health estimates [internet]*. Geneva: WHO, 2017. Available: <http://apps.who.int/iris/bitstream/10665/254610/1/WHO-MSD-MER-2017.2-eng.pdf>
- Moreno-Agostino D, Wu Y-T, Daskalopoulou C, et al. Global trends in the prevalence and incidence of depression: a systematic review and meta-analysis. *J Affect Disord* 2021;281:235–43.
- Murray CJL, Barber RM, Foreman KJ, et al. Global, regional, and national disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188 countries, 1990–2013: quantifying the epidemiological transition. *Lancet* 2015;386:2145–91.
- Cipriani A, Furukawa TA, Salanti G, et al. Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: a systematic review and network meta-analysis. *Lancet* 2018;391:1357–66.
- COVID-19 Mental Disorders Collaborators. Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *Lancet* 2021;398:1700–12.
- Spijker J, Bijl RV, de Graaf R, et al. Determinants of poor 1-year outcome of DSM-III-R major depression in the general population: results of the Netherlands mental health survey and incidence study (nemesi). *Acta Psychiatr Scand* 2001;103:122–30.
- Collins PY, Patel V, Joestl SS, et al. Grand challenges in global mental health. *Nature* 2011;475:27–30.
- Institute of Medicine. *Clinical practice cpqs we can trust (US)*. Washington, DC: National Academies Press, 2011.
- Brignardello-Petersen R, Carrasco-Labra A, Guyatt GH. How to interpret and use a clinical practice guideline or recommendation: users' guides to the medical literature. *JAMA* 2021;326:1516–23.
- AGREE Next Steps Consortium. The AGREE-II instrument [electronic version]. [place unknown]: AGREE. 2017. Available: <http://www.agreetrust.org> [Accessed 7 May 2018].
- Swinglehurst D. Evidence-Based guidelines: the theory and the practice. *Evidence-Based Healthcare and Public Health* 2005;9:308–14.
- Hoffmann-Eßer W, Siering U, Neugebauer EAM, et al. Guideline appraisal with agree II: systematic review of the current evidence on how users handle the 2 overall assessments. *PLoS One* 2017;12:e0174831.
- Hoffmann-Eßer W, Siering U, Neugebauer EAM, et al. Guideline appraisal with agree II: online survey of the potential influence of agree II items on overall assessment of guideline quality and recommendation for use. *BMC Health Serv Res* 2018;18:143.
- Brouwers MC, Kho ME, Browman GP, et al. Agree II: advancing Guideline development, reporting and evaluation in health care. *CMAJ* 2010;182:E839–42.
- AGREE-REX Research Team. The appraisal of CPGs research & evaluation-recommendation excellence (AGREE-REX) [internet]. AGREE. 2017. Available: <https://www.agreetrust.org/wp-content/uploads/2019/04/AGREE-REX-2019.pdf> [Accessed 17 May 2019].



- 19 Brouwers MC, Spithoff K, Kerkvliet K, *et al.* Development and validation of a tool to assess the quality of clinical practice guideline recommendations. *JAMA Netw Open* 2020;3:e205535.
- 20 Florez ID, Brouwers MC, Kerkvliet K, *et al.* Assessment of the quality of recommendations from 161 clinical practice guidelines using the appraisal of guidelines for research and evaluation-recommendations excellence (AGREE-REX) instrument shows there is room for improvement. *Implement Sci* 2020;15:79.
- 21 Molino C de GRC, Leite-Santos NC, Gabriel FC, *et al.* Factors associated with high-quality guidelines for the pharmacologic management of chronic diseases in primary care: a systematic review. *JAMA Intern Med* 2019;179:553–60.
- 22 Gabriel FC, Stein AT, de Melo DO, *et al.* Recommendations for the pharmacological treatment of treatment-resistant depression: a systematic review protocol. *PLoS One* 2022;17:e0267323.
- 23 Chronide.org [internet]. [place unknown]: chronide. 2019. Available: <https://www.chronide.org/>
- 24 Ministerio de Salud (COL). Detección temprana y diagnóstico del episodio depresivo y trastorno depresivo recurrente en adultos: atención integral de los adultos con diagnóstico de episodio depresivo o trastorno depresivo recurrente: guía de práctica clínica (bogotá: minsalud). 2013. Available: <http://www.iets.org.co/reportes-iets/Documentacin%20Reportes/Gu%C3%ADa.Completa.Depresion.2013.pdf> [Accessed 30 Jun 2017].
- 25 Ministerio de Salud. *Depresión en personas de 15 años y más: CL*. Santiago: Minsal, 2013.
- 26 National Institute for Health and Care Excellence. *Depression in adults: recognition and management*. London: NICE, 2009. Available: <https://www.nice.org.uk/guidance/cg90/evidenc>
- 27 Härter M, Schorr S, Schneider F, eds. *S3-leitlinie: nationale versorgungsleitlinie unipolare depression langfassung*. 2nd ed. Berlin: Springer, 2015.
- 28 Instituto de Evaluación de Tecnologías en Salud e Investigación (PE). *Guía de práctica clínica para el tamizaje y el manejo del episodio depresivo leve en el primer nivel de atención*. Lima: EsSalud, 2019. Available: [http://www.essalud.gob.pe/ietsi/pdfs/tecnologias\\_sanitarias/GPC\\_Depresion\\_V\\_Extensa.pdf](http://www.essalud.gob.pe/ietsi/pdfs/tecnologias_sanitarias/GPC_Depresion_V_Extensa.pdf)
- 29 American Psychological Association. *Clinical practice guideline for the treatment of depression across three age cohorts*. Washington, DC: American Psychological Association, 2019. Available: <https://www.apa.org/depression-guideline/guideline.pdf>
- 30 Management of Major Depressive Disorder Working Group. *VA/dod clinical practice guideline for the management of major depressive disorder*. 3rd ed. Washington, DC: United States Department of Veterans Affairs, US Department of Defence, 2016.
- 31 Trangle M, Gursky J, Haight R, *et al.* *Depression in primary care*. Bloomington, MN: ICSI, 2016. Available: [https://www.icsi.org/CPGs\\_more/catalog\\_CPGs\\_and\\_more/catalog\\_CPGs/catalog\\_behavioral\\_health\\_CPGs/depression](https://www.icsi.org/CPGs_more/catalog_CPGs_and_more/catalog_CPGs/catalog_behavioral_health_CPGs/depression)
- 32 Kaiser Permanente care management institute. *Diagnosis and treatment of depression in adults: 2012 clinical practice guideline*. Oakland, CA: KPCMI, 2012.
- 33 García-Herrera P, Nogueras MEV, Muñoz CM, *et al.* *Guía de práctica clínica para el tratamiento de la depresión en atención primaria*. Málaga: Hospital Regional Universitario "Carlos Haya, 2011.
- 34 Working group of the clinical practice guideline on the management of depression in adults. *Clinical practice guideline on the management of depression in adults*. Madrid: Ministerio de Sanidade, Servicios Sociales e Igualdad, 2014.
- 35 Registered nurses' association of Ontario. *Delirium, dementia, and depression in older adults: assessment and care*. 2nd ed. Toronto, ON: RNAO, 2016.
- 36 Qaseem A, Barry MJ, Kansagara D, *et al.* Nonpharmacologic versus pharmacologic treatment of adult patients with major depressive disorder: a clinical practice guideline from the American College of physicians. *Ann Intern Med* 2016;164:350–9.
- 37 Instituto Mexicano del Seguro Social. *Guía de práctica clínica gpc: diagnóstico y tratamiento de la depresión en el adulto mayor en el primer nivel de atención*. Ciudad de México: IMSS, 2011.
- 38 Secretaría de Salud, S. (MX). *Diagnóstico y tratamiento del trastorno depresivo en el adulto*. Ciudad de México: Secretaría de Salud, 2015.
- 39 Secretaría de Salud, S. (MX). *Diagnóstico y tratamiento del trastorno depresivo en el adulto: intervenciones de enfermería para la detección, atención y control de la depresión en el adulto mayor en los tres niveles de atención*. Ciudad de México: Secretaría de Salud, 2016.
- 40 Chua HC, Chan LL, Chee KS, *et al.* *Depression*. Singapore: Ministry of Health, 2012. Available: <http://www.smj.org.sg/sites/default/files/5302/5302cpg1.pdf>
- 41 Alonso-Coello P, Irfan A, Solà I, *et al.* The quality of clinical practice guidelines over the last two decades: a systematic review of guideline appraisal studies. *Qual Saf Health Care* 2010;19:e58.
- 42 Bancos I, Cheng T, Prokop LJ, *et al.* Endocrine clinical practice guidelines in North America. A systematic assessment of quality. *J Clin Epidemiol* 2012;65:520–5.
- 43 Don-Wauchope AC, Sievenpiper JL, Hill SA, *et al.* Applicability of the agree II instrument in evaluating the development process and quality of current national Academy of clinical biochemistry guidelines. *Clin Chem* 2012;58:1426–37.
- 44 MacQueen G, Santaguida P, Keshavarz H, *et al.* Systematic review of clinical practice guidelines for failed antidepressant treatment response in major depressive disorder, dysthymia, and subthreshold depression in adults. *Can J Psychiatry* 2017;62:11–23.
- 45 Hopkins D. *Framework for action on interprofessional education and collaborative practice*. Geneva: WHO, 2010.
- 46 Oosterhuis WP, Bruns DE, Watine J, *et al.* Evidence-Based guidelines in laboratory medicine: principles and methods. *Clin Chem* 2004;50:806–18.
- 47 Armstrong MJ, Mullins CD, Gronseth GS, *et al.* Impact of patient involvement on clinical practice guideline development: a parallel group study. *Implement Sci* 2018;13:55.
- 48 Tong A, Tunnicliffe DJ, Lopez-Vargas P, *et al.* Identifying and integrating consumer perspectives in clinical practice guidelines on autosomal-dominant polycystic kidney disease. *Nephrology (Carlton)* 2016;21:122–32.
- 49 Cronin RM, Mayo-Gamble TL, Stimpson S-J, *et al.* Adapting medical guidelines to be patient-centered using a patient-driven process for individuals with sickle cell disease and their caregivers. *BMC Hematol* 2018;18:12.
- 50 Rocha FL, Hara C, Paprocki J. Doença mental e estigma. *Rev Med Minas Gerais* 2015;25:590–6.
- 51 Zafra-Tanaka JH, Goicochea-Lugo S, Villarreal-Zegarra D, *et al.* Characteristics and quality of clinical practice guidelines for depression in adults: a scoping review. *BMC Psychiatry* 2019;19:76.
- 52 Owolabi M, Olowoyo P, Miranda JJ, *et al.* Gaps in hypertension guidelines in low- and middle-income versus high-income countries: a systematic review. *Hypertension* 2016;68:1328–37.
- 53 Bayona H, Owolabi M, Feng W, *et al.* A systematic comparison of key features of ischemic stroke prevention guidelines in low- and middle-income vs. high-income countries. *J Neurol Sci* 2017;375:360–6.



## Supplemental Material 1 – Methods details

### Theoretical rationale for choosing variables of interest to assess their association with the quality of CPGs and their recommendations

- 1) **Handling of conflicts of interest:** In order to have impartial and reliable recommendations, a CPG must not only declare conflicts but also have a handling of conflicts of interest policy. This policy is commonly achieved by editorial independence. Although we have not used in our analyses a variable directly related to the AGREE II domain “Editorial independence”, we have assessed “Handling of conflicts of interest”, which could be considered a proxy of this domain and is a fundamental characteristic for unbiased recommendations.[1]
- 2) **Multiprofessional team:** The development of high-quality CPGs is frequently associated with multidisciplinary teams in order to incorporate different knowledge and apply them in the recommendations. [1,2]
- 3) **Inclusion of patient representative in the team:** This characteristic usually occurs in clinical practice guidelines (CPGs) of higher methodological quality. In addition, interested party’s participation (*i.e.*, patients) is fundamental to the development of acceptable recommendations (Guidelines International Network, 2020). [3]
- 4) **Governmental funding:** Government-funded teams are more likely to have a larger number of professionals involved and more resources allocated allowing a longer and thoroughly review. Developing high quality CPGs usually requires these resources. [1,4]
- 5) **Type of institution:** The type of institution was considered in the model since this variable has been mentioned by previous studies to be associated with higher scores on the AGREE II “rigour of development”. [4]
- 6) **Publication year after 2015:** We hypothesized that newer CPGs and updated or revised versions tend to be associated with higher “rigour of development and highest quality” scores of the recommendations. [4]

### Identifying and selecting CPGs

For the selection of published CPGs, we searched for documents published from January 1, 2011 to December 31, 2021. A period longer than 10 years was considered because some CPGs are not systematically updated. In the occurrence of two or more versions of the same guideline we included only the most recent updated version. The following databases were searched: MEDLINE (via PubMed), Cochrane Library, Embase, PsycINFO, and BVS. Additional searches were performed in the following websites: Australian Clinical Practice Guidelines, National Committee for Health Technology Incorporation (Brazilian Government), Canadian Agency for Drugs and Technologies in Health, Canadian Medical Association, Chilean Ministry of Health, Colombian Ministry of Health and Social Protection, Guidelines International Network, Institute for Clinical Systems Improvement, Portal Guía Salud, Scottish Intercollegiate Guidelines Network, and the National Institute for Health and Care Excellence, Guideline Central, ECRI library. Specific details related to the databases used in this study and the terms used in these repositories are shown below:

#### Medline (via PubMed)

(((((("Guideline" [Publication Type] OR "CPGs as Topic"[Mesh] OR "Practice Guideline" [Publication Type] OR "Health Planning CPGs"[Mesh]) OR "Clinical Protocols"[Mesh])) OR ("Consensus Development Conference, NIH" [Publication Type] OR "Consensus Development

Conference" [Publication Type] OR "Consensus"[Mesh])) OR "Standard of Care"[Mesh]))  
 "Guideline" [Publication Type] OR "CPGs as Topic"[Mesh] OR "Practice Guideline"  
 [Publication Type] OR "Health Planning CPGs"[Mesh]) OR "Clinical Protocols"[Mesh])) OR  
 ("Consensus Development Conference, NIH" [Publication Type] OR "Consensus Development  
 Conference" [Publication Type] OR "Consensus"[Mesh])) OR "Standard of Care"[Mesh]))))  
 AND (("Depressive Disorder"[Mesh] OR "Depressive Disorder, Major"[Mesh] OR Depressive  
 Disorders OR Disorder, Depressive OR Disorders, Depressive OR Neurosis, Depressive OR  
 Depressive Neuroses OR Depressive Neurosis OR Neuroses, Depressive OR Depression,  
 Endogenous OR Depressions, Endogenous OR Endogenous Depression OR Endogenous  
 Depressions OR Depressive Syndrome OR Depressive Syndromes OR Syndrome, Depressive  
 OR Syndromes, Depressive OR Depression, Neurotic OR Depressions, Neurotic OR Neurotic  
 Depression OR Neurotic Depressions OR Melancholia OR Melancholias OR Unipolar  
 Depression OR Depression, Unipolar OR Depressions, Unipolar OR Unipolar Depressions))  
 Embase:  
 #1 'practice guideline'/mj OR 'consensus development'/exp/mj OR 'clinical protocol'/mj #2  
 'depression'/exp  
 #3 #1 AND #2

### Cochrane Library

# 1—MeSH descriptor: [Guideline] explode all trees  
 # 1—MeSH descriptor: [Consensus] explode all trees  
 # 1—MeSH descriptor: [Clinical Protocols] explode all trees  
 # 1—#1 OR #2 OR #3  
 # 1—MeSH descriptor: [Depression] explode all trees

### PsychINFO

((Any Field: (depression))) AND ((Any Field: (guideline)) OR (Any Field: (consensus)) OR  
 (Any Field: ("clinical protocol"))) AND Year: 2011 To 9999  
 ((Any Field: (depression))) AND ((Any Field: (guideline)) OR (Any Field: (consensus)) OR (Any  
 Field: ("clinical protocol"))) AND Year: 2011 To 9999

### BVS

((guideline) OR (consensus) OR (clinical protocol)) AND (depression) AND (db:("LILACS" OR  
 "IBECs" OR "WHOLIS" OR "BDENF" OR "BINACIS" OR "INDEXPSI" OR "BIGG" OR  
 "BBO" OR "CUMED")) AND (year\_cluster:[2011 TO 2021])

We included documents between January 1, 2011, to December 31, 2021; containing recommendations for the pharmacological treatment of depression in adults at outpatient care setting, regardless of whether it met the U.S. National Academy of Medicine. If a CPG included recommendations for both children and adults, they were considered. No language restriction was applied; CPGs that were in languages other than Portuguese, English, or Spanish were translated into Portuguese by a professional translation service. We excluded CPGs for specific populations (e.g., treatment of pregnant women); local CPGs developed by hospitals or originations intended to be applied only at a local level and treatment of depression in comorbidity with specific diseases (e.g., depression in patients with diabetes).

Retrieved references were exported to the online platform Rayyan® reference manager. [5] After removal of duplicates, references were screened by two independent researchers. Then searched for the full texts, and these were reviewed in duplicate. Discrepancies between researchers were resolved by consensus. When no consensus was reached, a third evaluator was involved.

## Extracting the characteristics of the CPGs

A Google Form was used for the extraction of general CPG data. The process of data extraction by two independent researchers was validated in a previous project [4] focused on osteoporosis CPGs conducted by our team. [6] We extracted the following independent variables: year of publication of the most recent version of the CPGs (2011 to 2014/2015 to 2021); type of institution (Governmental or University; Professional society), inclusion of a patient representative in the team (yes/no), multiprofessional team (yes - different professions/no - only one profession), governmental funding (yes/no), and policy for handling of conflicts of interest (yes/no). The type of institution classification and the funding variables were defined as governmental, even if the governmental institution had partnership with other types of institutions. These factors were selected because they are included in the AGREE II [7], AGREE-REX [8] and IOM. [9] Additionally, they are commonly found in literature articles [4,10] related to guideline's quality.

## Appraisal of the CPGs quality

AGREE II is a reliable and validated tool composed by 23 items, clustered in six domains. The appraisal team comprised multidisciplinary researchers, including pharmacists, nurses, and public health professionals previously trained according to protocol. [11] The training consisted of reading the AGREE II manual. [7] Subsequently, the evaluators appraised the CPG quality on chronic pain [12], Gaucher disease [13], and for the treatment of obesity. [14] A discussion was held with previously trained evaluators on the evaluations made. In the next stage of training, the team appraised the most recent two CPGs for hyperthyroidism [15] and urinary tract infection. [16]

The appraisers assigned a score from 1 to 7 for each AGREE II item (following a 7-point Likert scale). Each guideline was appraised by three appraisers as suggested by the AGREE Next Steps Consortium [7] using the AGREE-PLUS Platform. [17] A difference of two or more points in the individual items' [4,11,18] scores was considered discrepant and was resolved by consensus between the appraisers to obtain the final score. A final consolidated score was obtained from 0-100% per domain as suggested by the AGREE II Manual.

## Appraisal of the quality of the recommendations of the CPGs

We used the AGREE-REX instrument to appraise the quality of the CPG recommendations. All recommendations were grouped and analysed for each CPG, according to one of the options recommended in the AGREE-REX manual. [8] AGREE-REX consists of nine items clustered into three domains: clinical applicability, values and preferences, and implementability. The same team of appraisers that assessed the quality of the CPGs using AGREE II also assessed the quality of the recommendations. The team was also trained for this evaluation. The three appraisers assigned a score from 1 to 7 for each item (following a 7-point Likert scale). When there was a discrepancy of two points or more, the evaluators discussed it until they reached consensus. A final consolidated score was obtained from 0-100% per AGREE-REX domain as suggested by the AGREE-REX Manual. [8] The scores of each evaluator were inserted in a Google Form®.

## Statistical data analysis

Quantitative data were described using mean, standard deviation, median, and interquartile range. Categorical variables were presented as frequencies and percentages.



The CPGs were considered high-quality if they scored  $\geq 60\%$  in the AGREE II Rigour of Development domain, as it has been previously recommended. [4,10,19-22] The CPGs' recommendations were considered high-quality if they scored  $\geq 60\%$  in the AGREE-REX Clinical Applicability domain which considers the appropriateness of recommendations for clinical practice, patient needs, and the intended impact of guideline implementation.

Since the outcomes of high-quality CPGs and high-quality recommendations showed only 17 and 7 events, respectively, instead of analysing the binary characteristics we decided to model the scores directly. We used linear regression analysis and with this approach we were able to estimate the impact that each studied factor had on the AGREE II domain 3 and AGREE-REX domain 1 scores. The regression coefficients were obtained with a simple unadjusted model (univariate) and with an adjusted model (multivariable). Results with p values below 0.05 were deemed statistically significant. Data were processed and analyzed with IBM-SPSS version 25.0.

**Supplemental material 2 – Reasons for excluding clinical practice guideline (CPG)**

Reference of the excluded CPG	Reasons for exclusion
1. Institute for Clinical Systems Improvement. Adult Depression in Primary Care. Bloomington, MN: ICSI, 2016.	Duplicate
2. Grinspun, D, Bajnok I, Rey M. Delirium, Dementia, and Depression in Older Adults: Assessment and Care. Toronto, Canada: Registered 'urses' Association of Ontario, 2016.	Duplicate
3. Boltz M (Ed.). Evidence-based Geriatric Nursing Protocols for Best Practice. New York, NY: Springer, 2012.	Duplicate
4. MICHIGAN MEDICINE. Ambulatory unipolar depression guideline. [Ann Arbor]: UM; 2021. Available from: <a href="https://michmed-public.policystat.com/policy/8093108/latest/">https://michmed-public.policystat.com/policy/8093108/latest/</a> .	Duplicate
5. Álvarez Ariza M, Atienza Merino G, Ávila González MJ, <i>et al</i> . GPC sobre el Manejo de la Depresión en el Adulto. Madrid, Spain: Ministerio de Sanidad, Servicios Sociales e Igualdad, 2014.	Duplicate
6. National Guideline Clearinghouse. Depression (Singapore). Rockville, MD: Agency for Healthcare Research and Quality (AHRQ), 2012. Available from: <a href="https://www.guideline.gov/summaries/summary/39324">https://www.guideline.gov/summaries/summary/39324</a> .	Duplicate
7. Management of Major Depressive Disorder Working Group. VA/DoD clinical practice guideline for the management of major depressive disorder. 3rd ed. Washington, DC: US Department of Veterans Affairs, US Department of Defense, 2016.	Duplicate
8. Austin M-P, Hight N, The Expert Working Group. Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline. Melbourne, Australia: Centre of Perinatal Excellence, 2017.	Focused on specific populations
9. McDermott B, Baigent M, Chanen A, <i>et al</i> . Clinical Practice Guidelines: Depression in Adolescents and Young Adults. Melbourne, Australia: Agency for Healthcare Research and Quality, 2010.	Focused on specific populations
10. Michigan Quality Improvement Consortium Guideline. Primary Care Diagnosis and Management of Adults with Depression. Detroit, MI: MQIC; 2018. Available from: <a href="http://mqic.org/guidelines.htm">http://mqic.org/guidelines.htm</a> .	Duplicate
11. National Institute for Health and Clinical Excellence. Depression in Children and Young People: Identification and Management in Primary, Community and Secondary Care. Leicester, UK: British Psychological Society, 2005.	Focused on specific populations
12. Connolly KR, Thase ME. If at first you don't succeed: a review of the evidence for antidepressant augmentation, combination and switching strategies. <i>Drugs</i> 2011, 71(1):43-64. DOI:10.2165/11587620-000000000-00000.	Duplicate
13. Kennedy SH, Lam RW, McIntyre RS, <i>et al</i> . Canadian network for mood and anxiety treatments (CANMAT), 2016, clinical CPGs for the management of adults with major depressive disorder: section 3: pharmacological treatments. <i>Can J Psychiatry</i> . 2016, 61(9)540-560. DOI: <a href="https://doi.org/10.1177/0706743716659417">10.1177/0706743716659417</a> .	Duplicate

Reference of the excluded CPG	Reasons for exclusion
14. Taylor RW, Marwood L, Oprea E, <i>et al.</i> Pharmacological Augmentation in Unipolar Depression: A Guide to the Guidelines. <i>Int J Neuropsychopharmacol.</i> 2020, 23(9):587-625. DOI:10.1093/ijnp/pyaa033.	Duplicate
15. Voineskos D, Daskalakis ZJ, Blumberger DM. Management of Treatment-Resistant Depression: Challenges and Strategies. <i>Journal of Clinical Psychopharmacology</i> . 2020;16:221-234. Published 2020 Jan 21. DOI: 10.2147/NDT.S198774.	Duplicate
16. Samochowiec J, Dudek D, Kucharska Mazur J, Murawiec S, Rymaszewska J, Cubała WJ, Heitzman J, Szulc A, Bała M, Gątecki P. Pharmacological treatment of a depressive episode and recurrent depressive disorder - guidelines of the Polish Psychiatric Association and the National Consultant for Adult Psychiatry. <i>Psychiatr Pol.</i> 2021 Mar 12;55(2):235-259. English, Polish. DOI: 10.12740/PP/ OnlineFirst/132496. Epub 2021 Mar 12. PMID: 34365477.	Duplicate
17. MICHIGAN MEDICINE. Ambulatory unipolar depression guideline. [Ann Arbor]: UM; 2021. Available from: <a href="https://michmed-public.policystat.com/policy/8093108/latest/">https://michmed-public.policystat.com/policy/8093108/latest/..</a>	Duplicate
18. MALAYSIAN HEALTH TECHNOLOGY ASSESSMENT SECTION. Management of major depressive disorder. 2nd ed. Putrajaya: MaHTAS, 2019. Available from: <a href="http://www.psychiatry-malaysia.org/">http://www.psychiatry-malaysia.org/...</a>	Duplicate
19. Malhi GS, Bell E, Boyce P, Bassett D, Berk M, Bryant R, Gitlin M, Hamilton A, Hazell P, Hopwood M, Lyndon B, McIntyre RS, Morris G, Mulder R, Porter R, Singh AB, Yatham LN, Young A, Murray G. The 2020 Royal Australian and New Zealand College of psychiatrists clinical practice guidelines for mood disorders: Bipolar disorder summary. <i>Bipolar Disord.</i> 2020 Dec, 22(8):805-821. DOI: 10.1111/bdi.13036. PMID: 33296123.	Subject matter
20. Moeller SB, Gbyl K, Hjorthøj C, Andreasen M, Austin SF, Buchholtz PE, Fønss L, Hjerrild S, Hogervorst L, Jørgensen MB, Ladegaard N, Martiny K, Meile J, Packness A, Sigaard KR, Straarup K, Straszek SPV, Soerensen CH, Welcher B, Videbech P. Treatment of difficult-to-treat depression - clinical guideline for selected interventions. <i>Nord J Psychiatry.</i> 2022 Apr, 76(3):177-188. DOI: 10.1080/08039488.2021.1952303. Epub 2021 Aug 28. PMID: 34455900.	Duplicate
21. Kennedy SH, Lam RW, McIntyre RS, <i>et al.</i> Canadian network for mood and anxiety treatments (CANMAT) 2016 clinical guidelines for the management of adults with major depressive disorder: section 3. pharmacological treatments. <i>Can J Psychiatry</i> 2016, 61:540-60.	Duplicate
22. National Institute for Health and Care Excellence. Depression in Adults: Recognition and Management. 2009. Available from: <a href="https://www.nice.org.uk/guidance/cg90/evidence">https://www.nice.org.uk/guidance/cg90/evidence</a> .	Duplicate



Reference of the excluded CPG	Reasons for exclusion
23. Ministerio de Salud (CL). Depresión em personas de 15 años y más. Santiago: Minsal, 2013.	Duplicate
24. Ministerio de Salud (COL). Detección temprana y diagnóstico del episodio depresivo y trastorno depresivo recurrente en adultos: atención integral de los adultos con diagnóstico de episodio depresivo o trastorno depresivo recurrente: Guía de práctica clínica. Bogotá: Minsalud; 2013 [cited 2017 Jun 30]. Available from: <a href="http://www.iets.org.co/reportes-iets/Documentacin%20Reportes/Gu%C3%ADa. Completa. Depresion, 2013.pdf">http://www.iets.org.co/reportes-iets/Documentacin%20Reportes/Gu%C3%ADa. Completa. Depresion, 2013.pdf</a>	Duplicate
25. Álvarez Ariza M, Atienza Merino G, Ávila González MJ, <i>et al.</i> GPC sobre el Manejo de la Depresión en el Adulto. Madrid, Spain: Ministerio de Sanidad, Servicios Sociales e Igualdad, 2014.	Duplicate
26. National Guideline Clearinghouse. Depression (Singapore). Rockville, MD: Agency for Healthcare Research and Quality (AHRQ); 2012. Available from: <a href="https://www.guideline.gov/summaries/summary/39324..">https://www.guideline.gov/summaries/summary/39324..</a>	Duplicate
27. American Psychological Association. (2019). Clinical practice guideline for the treatment of depression across three age cohorts. Retrieved from <a href="https://www.apa.org/depression-guideline">https://www.apa.org/depression-guideline</a> .	Duplicate
28. Bennabi D, Charpeaud T, Yrondi A, Genty JB, Destouches S, Lancrenon S, Alaïli N, Bellivier F, Bougerol T, Camus V, Dorey JM, Doumy O, Haesebaert F, Holtzmann J, Lançon C, Lefebvre M, Moliere F, Nieto I, Rabu C, Richieri R, Schmitt L, Stephan F, Vaiva G, Walter M, Leboyer M, El-Hage W, Llorca PM, Courtet P, Aouizerate B, Haffen E. Clinical guidelines for the management of treatment-resistant depression: French recommendations from experts, the French Association for Biological Psychiatry and Neuropsychopharmacology and the fondation FondaMental. BMC Psychiatry. 2019 Aug 28, 19(1):262. DOI: 10.1186/s12888-019-2237-x. PMID: 31455302, PMCID: PMC6712810.	Duplicate
29. Bennabi D, Yrondi A, Charpeaud T, Genty JB, Destouches S, Lancrenon S, Allaili N, Bellivier F, Bougerol T, Camus V, Doumy O, Dorey JM, Haesebaert F, Holtzmann J, Lançon C, Lefebvre M, Moliere F, Nieto I, Rabu C, Richieri R, Schmitt L, Stephan F, Vaiva G, Walter M, Leboyer M, El-Hage W, Aouizerate B, Haffen E, Llorca PM, Courtet P. Clinical guidelines for the management of depression with specific comorbid psychiatric conditions French recommendations from experts (the French Association for Biological Psychiatry and Neuropsychopharmacology and the fondation FondaMental). BMC Psychiatry. 2019 Jan 30, 19(1):50. DOI: 10.1186/s12888-019-2025-7. PMID: 30700272, PMCID: PMC6354367.	Duplicate
30. Malhi, G. S. <i>et al.</i> Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for mood disorders: major depression summary. Medical Journal of Australia, Sydney, v. 208, n. 5, 2018. DOI <a href="http://doi.org/10.5694/mja17.00659">http://doi.org/10.5694/mja17.00659</a> . Available from: <a href="https://www.mja.com.au/journal/2018/208/4/royal-australian-and-new-zealand-college-psychiatrists-clinical-practice">https://www.mja.com.au/journal/2018/208/4/royal-australian-and-new-zealand-college-psychiatrists-clinical-practice</a> .	Duplicate

Reference of the excluded CPG	Reasons for exclusion
31. Nonpharmacologic Versus Pharmacologic Treatment of Adult Patients With Major Depressive Disorder: A Clinical Practice Guideline From the American College of Physicians, 2016. <a href="https://www.acpjournals.org/doi/10.7326/M15-2570">https://www.acpjournals.org/doi/10.7326/M15-2570</a> .	Duplicate
32. Evidence-based guidelines for treating depressive disorders with antidepressants: A revision of the 2008 British Association for Psycho. Available from: <a href="https://www.bap.org.uk/pdfs/BAP_Guidelines-Antidepressants.pdf">https://www.bap.org.uk/pdfs/BAP_Guidelines-Antidepressants.pdf</a> . pharmacologyguidelines. Journal of Psychopharmacology, 2015, v. 29(5) 459-525 DOI: 10.1177/0269881115581093 <a href="http://jop.sagepub.com">jop.sagepub.com</a> .	Duplicate
33. National Institute for Health and Care Excellence. Common Mental Health Disorders. Identification and Pathways to Care. London, UK: NICE, 2011.	Did not include pharmacological treatment
34. Patten SB. Updated CANMAT guidelines for treatment of major depressive disorder. Can J Psychiatry 2016, 61(9):504-5. DOI:10.1177/0706743716660034.	Did not include pharmacological treatment
35. Rush AJ, Aaronson ST, Demyttenaere K. Difficult-to-treat depression: a clinical and research roadmap for when remission is elusive. Aust N Z J Psychiatry 2019, 53(2):109-18. DOI:10.1177/0004867418808585.	Did not include pharmacological treatment
36. Repetitive transcranial magnetic stimulation for depression NICE, 2015. Available from: <a href="https://www.nice.org.uk/guidance/ipg542/resources/repetitive-transcranial-magnetic-stimulation-for-depression-pdf-1899871923433669">https://www.nice.org.uk/guidance/ipg542/resources/repetitive-transcranial-magnetic-stimulation-for-depression-pdf-1899871923433669</a> .	Did not include pharmacological treatment
37. Bennabi D, Haffen E. Transcranial Direct Current Stimulation (tDC): A Promising Treatment for Major Depressive Disorder? Brain Sci. 2018, 8(5):81. Published 2018 May 6. DOI:10.3390/brainsci8050081.	Did not include pharmacological treatment
38. Thompson W, Adcock L. Intensive Day Treatment Programs for Mental Health Treatment: A Review of Clinical Effectiveness, Cost-Effectiveness, and Guidelines [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2017 Sep 15. Available from: <a href="https://www.ncbi.nlm.nih.gov/books/NBK513364/">https://www.ncbi.nlm.nih.gov/books/NBK513364/</a> .	Did not include pharmacological treatment
39. Ho C, Adcock L. Short-Term Psychodynamic Psychotherapy for the Treatment of Mental Illness: A Review of Clinical Effectiveness and Guidelines [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2017 Oct 12. PMID: 30234931.	Did not include pharmacological treatment
40. Palylyk-Colwell E, Argáez C. Telehealth for the Assessment and Treatment of Depression, Post-Traumatic Stress Disorder, and Anxiety: Clinical Evidence [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2018 Apr 10. Disponible en: <a href="https://www.ncbi.nlm.nih.gov/books/NBK532701/">https://www.ncbi.nlm.nih.gov/books/NBK532701/</a> .	Did not include pharmacological treatment
41. Pohar R, Farrah K. Repetitive Transcranial Magnetic Stimulation for Patients with Depression: A Review of Clinical Effectiveness, Cost-Effectiveness and Guidelines – An Update [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2019 Jun 28. PMID: 31433608.	Did not include pharmacological treatment

Reference of the excluded CPG	Reasons for exclusion
42. Ho C, Severn M. e-Therapy Interventions for the Treatments of Patients with Depression: A Review of Clinical Effectiveness [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2018 May 16. Available from: <a href="https://www.ncbi.nlm.nih.gov/books/NBK532214/">https://www.ncbi.nlm.nih.gov/books/NBK532214/</a> .	Did not include pharmacological treatment
43. CANADIAN AGENCY FOR DRUGS AND TECHNOLOGIES IN HEALTH. Internet-delivered cognitive behavioural therapy for major depressive disorder and anxiety disorders. [Ottawa, ON]: CADTH, 2019. Available from: <a href="https://www.cadth.ca/brief-internet-delivered-cognitive-behavioural-therapy-major-depressive-disorder-and-anxiety">https://www.cadth.ca/brief-internet-delivered-cognitive-behavioural-therapy-major-depressive-disorder-and-anxiety</a> .	Did not include pharmacological treatment
44. Frye MA. Clinical practice: bipolar disorder--a focus on depression. N Engl J Med 2011, 364(1):51-9.	Subject matter
45. Andersen BL, DeRubeis RJ, Berman BS, <i>et al</i> . Screening, assessment, and care of anxiety and depressive symptoms in adults with cancer: an American Society of Clinical Oncology guideline adaptation. J Clin Oncol 2014, 32(15):1605-19. DOI:10.1200/JCO.2013.52.4611.	Focused on specific populations
46. American Medical Directors Association. Guideline Summary: Depression in the Long Term Care Setting. Columbia, MD: AMDA; 2011.	Subject matter
47. Li M, Kennedy EB, Byrne N, <i>et al</i> . The management of depression in patients with cancer: a clinical practice guide. J OncolPract 2016, 12(8):747-56. DOI:10.1200/JOP.2016. 011072.	Subject matter
48. Ostacher MJ, Tandon R, Suppes T. Florida best practice psychotherapeutic medication guidelines for adults with bipolar disorder: a novel, practical, patient-centered guide for clinicians. J Clin Psychiatry 2016, 77(7):920-6. DOI:10.4088/JCP.15cs 09841. Available from: <a href="http://www.embase.com/search/results?subaction=viewrecord&amp;from=export&amp;id=L611538719">http://www.embase.com/search/results?subaction=viewrecord&amp;from=export&amp;id=L611538719</a>	Subject matter
49. Pfennig A, Bschor T, Falkai P, <i>et al</i> . The diagnosis and treatment of bipolar disorder: recommendations from the current S3 guideline. Dtsch Arztebl Int 110(6):92-100. DOI:10.3238/arztebl. 2013.0092. 2013.	Subject matter
50. Rosenblat JD, McIntyre RS. Treatment recommendations for DSM-5-defined mixed features. CNS Spectr 2017, 22(2):147- 54. DOI:10.1017/S1092852916000432.	Subject matter
51. HEALTHCARE IMPROVEMENT SCOTTLAD. Scottish palliative care guidelines: depression. Edinburgh: NHSScotland, 2019. Available from: <a href="https://www.palliaticareguidelines.scot.nhs.uk/guidelines/symptom-control /Depression.aspx">https://www.palliaticareguidelines.scot.nhs.uk/guidelines/symptom-control /Depression.aspx</a> .	Subject matter
52. Assessment and Interventions for Perinatal Depression. Available from: <a href="https://rnao.ca/sites/rnao-ca/files/bpg/Perinatal_Depression_FINAL_web_0.pdf">https://rnao.ca/sites/rnao-ca/files/bpg/Perinatal_Depression_FINAL_web_0.pdf</a> Registered Nurses' Association of Ontario. Assessment and Interventions for Perinatal Depression. 2nd ed. Toronto (ON): Registered Nurses' Association of Ontario, 2018.	Focused on specific populations



Reference of the excluded CPG	Reasons for exclusion
53. Carneiro LF, Mota MP, Schuch F, Deslandes A, Vasconcelos-Raposo J. Portuguese and Brazilian guidelines for the treatment of depression: exercise as medicine. <i>Braz J Psychiatry</i> . 2017 Aug 30, 40(2):210-211. DOI: 10.1590/1516-4446-2017-2272. PMID: 28876380; PMCID: PMC6900767.	Did not include pharmacological treatment
54. JOBST, A. European Psychiatric Association Guidance on psychotherapy in chronic depression across Europe. <i>European Psychiatry</i> , v. 33, n. 2016, p. 18-36. DOI <a href="http://doi.org/10.1016/j.eurpsy.2015.12.003">http://doi.org/10.1016/j.eurpsy.2015.12.003</a> . Available from: <a href="http://www.europsy.net/app/uploads/2016/05/Psychotherapy-in-chronic-depression-across-Europe.pdf">http://www.europsy.net/app/uploads/2016/05/Psychotherapy-in-chronic-depression-across-Europe.pdf</a> .	Did not include pharmacological treatment
55. FSRH CEU Statement: Response to published study: Association of Hormonal Contraception with Depression (October 2016) <a href="file:///C:/Users/Usuario/Downloads/association-of-hc-with-depression%20(1).pdf">file:///C:/Users/Usuario/Downloads/association-of-hc-with-depression%20(1).pdf</a> .	Was not a CPG
56. Practice Guidelines for the Prevention, Detection, and Management of Respiratory Depression Associated with Neuraxial Opioid Administration: An Updated Report by the American Society of Anesthesiologists Task Force on Neuraxial Opioids and the American Society of Regional Anesthesia and Pain Medicine*. <i>Anesthesiology</i> 2016, 124:535–552 DOI: <a href="https://doi.org/10.1097/ALN.0000000000000975">https://doi.org/10.1097/ALN.0000000000000975</a> .	Subject matter
57. Manipulating the sleep-wake cycle and circadian rhythms to improve clinical management of major depression DOI:10.1186/1741-7015-11-79. <i>BMC Medicine</i> 2013, 11:79.	Subject matter
58. NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE. Space from depression for treating adults with depression. [London]: NICE, 2020. Available from: <a href="https://www.nice.org.uk/advice/mib215/resources/space-from-depression-for-treating-adults-with-depression-pdf-2285965453227973">https://www.nice.org.uk/advice/mib215/resources/space-from-depression-for-treating-adults-with-depression-pdf-2285965453227973</a> .	Subject matter
59. Goracci A, Forgione RN, De Giorgi R, Coluccia A, Cuomo S, Fagiolini A. Practical guidance for prescribing trazodone extended-release in major depression. <i>Expert Opin Pharmacother</i> . 2016, 17(3):433-41. DOI:	Specific drugs
60. Kandasamy S, Campbell K. Psychotherapy and Pharmacotherapy for Major Depressive Disorder and Generalized Anxiety Disorder: A Rapid Qualitative Review [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2019 Jun 4. PMID: 33074608.	Was not a CPG
61. Antidepressants in Elderly Patients with Major and Minor Depression: A Review of Clinical Effectiveness and Guidelines [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2015 Aug 17. Available from: <a href="https://www.ncbi.nlm.nih.gov/books/NBK315889/">https://www.ncbi.nlm.nih.gov/books/NBK315889/</a> .	Was not a CPG

Reference of the excluded CPG	Reasons for exclusion
62. WELLS, C., CAMPBELL, K. Antidepressants for adults with depression using opioid medications: clinical effectiveness and guidelines. Ottawa, ON: CADTH, 2018 Feb. (CADTH rapid response report: summary of abstracts).	Was not a CPG
63. Kim J, Mierzwinski-Urban M. Ketamine for Treatment-Resistant Depression or Post-Traumatic Stress Disorder in Various Settings: A Review of Clinical Effectiveness, Safety, and Guidelines [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2017 Mar 1. PMID: 29533569.	Focused on specific populations
64. TADROUS, M., LÉSELEUC, L. Drugs for major depressive disorder. Ottawa, ON: CADTH, 2020 Feb. (CADTH technology review: focused critical appraisal; n. 24). Available from: <a href="https://www.cadth.ca/major-depressive-disorder-focused-critical-appraisal-network-meta-analysis">https://www.cadth.ca/major-depressive-disorder-focused-critical-appraisal-network-meta-analysis</a> .	Was not a CPG
65. Diagnosing, Screening, and Monitoring Depression in the Elderly: A Review of Guidelines [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2015 Sep 8. PMID: 26468558.	Was not a CPG
66. E-NOTAS. Modelos e intervenciones de colaboración entre atención primaria y salud mental en la atención a los pacientes diagnosticados de depresión: resultados de una meta-revisión. [Oviedo]: 2014. Available from: <a href="https://enotas.astursalud.es/-/modelos-e-intervenciones-de-colaboraci%C3%B3n-entre-atenci%C3%B3n-primaria-y-salud-mental-en-la-atenci%C3%B3n-a-los-pacientes-diagnosticados-de-depresi%C3%B3n-resultados-de-una-meta-revisi%C3%B3n-">https://enotas.astursalud.es/-/modelos-e-intervenciones-de-colaboraci%C3%B3n-entre-atenci%C3%B3n-primaria-y-salud-mental-en-la-atenci%C3%B3n-a-los-pacientes-diagnosticados-de-depresi%C3%B3n-resultados-de-una-meta-revisi%C3%B3n-</a> .	Was not a CPG
67. MacDonald E, Horton J. Bupropion for Major Depressive Disorder or Persistent Depressive Disorder (Dysthymia) [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health; 2021 Apr. Available from: <a href="https://www.ncbi.nlm.nih.gov/books/NBK571946/">https://www.ncbi.nlm.nih.gov/books/NBK571946/</a> .	Was not a CPG
68. Poitras V, Visintini S. Desvenlafaxine versus Venlafaxine for the Treatment of Adult Patients with Major Depressive Disorder: A Review of the Comparative Clinical and Cost-Effectiveness [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2017 Oct 25. PMID: 29889482.	Was not a CPG
69. YATHAM, L. Cognitive heterogeneity in bd: validity, trajectories and biological mechanisms symposium [Symposium Abstracts; SY1]. Bipolar Disorders, Copenhagen v. 23, n. S1, p. 15-47. DOI <a href="https://doi.org/10.1111/bdi.13086">https://doi.org/10.1111/bdi.13086</a> . Available from: <a href="https://onlinelibrary.wiley.com/doi/epdf/10.1111/bdi.13086">https://onlinelibrary.wiley.com/doi/epdf/10.1111/bdi.13086</a> .	Was not a CPG
70. CANADIAN AGENCY FOR DRUGS AND TECHNOLOGIES IN HEALTH. Antidepressants administered using weight-based dosing strategies for depression: clinical effectiveness and guidelines. [Ottawa, ON]: CADTH; 2020. Available from: <a href="https://www.cadth.ca/antidepressants-administered-using-weight-based-dosing-strategies-depression-clinical-effectiveness">https://www.cadth.ca/antidepressants-administered-using-weight-based-dosing-strategies-depression-clinical-effectiveness</a> .	Was not a CPG

Reference of the excluded CPG	Reasons for exclusion
71. Li KX, Loshak H. Intravenous Ketamine for Adults with Treatment-Resistant Depression or Post-Traumatic Stress Disorder: A Review of Clinical Effectiveness, Cost-Effectiveness and Guidelines [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2019 Oct 24. PMID: 31873996.	Was not a CPG
72. Testosterone for depression, PTSD, or fibromyalgia: Clinical effectiveness and guidelines. Ottawa: CADTH, 2017, Feb. (CADTH rapid response report: reference list).	Was not a CPG
73. CANADIAN AGENCY FOR DRUGS AND TECHNOLOGIES IN HEALTH. <b>Monoamine oxidase inhibitors in elderly patients:</b> clinical effectiveness and guidelines. [Ottawa, ON]: CADTH, 2015. Available from: <a href="https://www.cadth.ca/sites/default/files/thisfile/htis/aug-2015/RA0792%20MAOIs%20in%20the%20Elderly%20Final.pdf">https://www.cadth.ca/sites/default/files/thisfile/htis/aug-2015/RA0792%20MAOIs%20in%20the%20Elderly%20Final.pdf</a> .	Was not a CPG
74. Alexopoulos GS. Pharmacotherapy for late-life depression. J Clin Psychiatry 2011, 72(1):e04. DOI:10.4088/JCP.7085tx2cj.	Was not a CPG
75. Andreescu C, Reynolds CF III. Late-life depression: evidencebased treatment and promising new directions for research and clinical practice. Psychiatr Clin North Am 2011, 34(2):335-55. DOI:10.1016/j.psc.2011.02.005.	Was not a CPG
76. Arnow BA, Steidtmann D, Blasey C, <i>et al.</i> The relationship between the therapeutic alliance and treatment outcome in two distinct psychotherapies for chronic depression. J Consult Clin Psychol 2013, 81(4):627-38. DOI:10.1037/a0031530.	Was not a CPG
77. Ayub-Dargél A, Masson M, Henry C. The RANZCP guidelines: managing mood disorders in the real world. Aust N Z J Psychiatry 2016, 50(12): 1198-9. DOI:10.1177/0004867416676373.	Was not a CPG
78. Bland P. Raising standards of care for patients with depression. Practitioner 2011, 255(1740):21-5.	Was not a CPG
79. Boffin N, Bossuyt N, Declercq T, <i>et al.</i> Incidence, patient characteristics and treatment initiated for GP-diagnosed depression in general practice: results of a 1-year nationwide surveillance study. Fam Prac 2012, 29(6):678-87. DOI:10.1093/fampra/cms024.	Was not a CPG
80. Bohra HM, Novak M. Depression in patients with chronic kidney disease. CANNT J 2015, 25(3):34-8.	Was not a CPG
81. Cohen A. The QOF, NICE, and depression. Br J Gen Pract 2011, 61(590):549. DOI:10.3399/bjgp11X593785.	Was not a CPG
82. Baumann S, Benson-Martin J, Cossie Q, <i>et al.</i> SASOP treatment guidelines for psychiatric disorders: eminence or evidence based? S Afr JPsychiatr 2019, 20(2):a529. DOI 10.4102/sajpsychiatry.v20i2.529.	Was not a CPG
83. Cosgrove L, Shaughnessy AF, Wheeler EE, <i>et al.</i> The American Psychiatric Association's guideline for major depressive disorder: a commentary. Psychother Psychosom 2012, 81(3):186-8. DOI:10.1159/000335523.	Was not a CPG

Reference of the excluded CPG	Reasons for exclusion
84. Cuijpers P. Effective therapies or effective mechanisms in treatment guidelines for depression? <i>Depress Anxiety</i> 2013, 30(11):1055-7. DOI:10.1002/da.22205.	Was not a CPG
85. Cuijpers P. Combined pharmacotherapy and psychotherapy in the treatment of mild to moderate major depression? <i>JAMA Psychiatry</i> 2014, 71(7):747-8. DOI:10.1001/jamapsychiatry. 2014. 277.	Was not a CPG
86. De Coteau PA, Byrne CD, Russell, V. The HSE/ICGP guidelines on the management of depression and anxiety disorders in primary care. <i>Ir Med J</i> 2012, 105(7):251.	Was not a CPG
87. Desseilles M, Witte J, Chang TE, <i>et al.</i> Assessing the adequacy of past antidepressant trials: a clinician's guide to the antidepressant treatment response questionnaire. <i>J Clin Psychiatry</i> 2011, 72(8):1152-4. DOI:10.4088/JCP. 11ac07225.	Was not a CPG
88. Canadian Agency for Drugs and Technologies in Health. Diagnosing, Screening, and Monitoring Depression in the Elderly: A Review of Guidelines. Ottawa, ON: CADTH, 2015.	Was not a CPG
89. Canadian Agency for Drugs and Technologies in Health. Antidepressants in Elderly Patients with Major and Minor Depression: A Review of Clinical Effectiveness and Guidelines. Ottawa, ON: CADTH, 2015.	Was not a CPG
90. Gensichen J, Härter M, Klesse C, <i>et al.</i> Germany's national clinical practice guideline (S3) for unipolar depression—What is important for family practice? <i>ZFA</i> 2011, 87(5):223-30. DOI:10.3238/zfa.2011.0223.	Was not a CPG
91. GEOFFROY, P. A., BELLIVIER, F. The RANZCP mood disorders guidelines: an easy step-by-step toolbox for daily practice. <i>Australian and New Zealand Journal of Psychiatry</i> , Melbourne, v. 50, n. 10, p. 1014-5. DOI Available from: <a href="https://journals.sagepub.com/doi/10.1177/0004867416667829">https://journals.sagepub.com/doi/10.1177/0004867416667829</a> .	Was not a CPG
92. Gitlin M. The Royal Australian and New Zealand College of Psychiatrists clinical guidelines for mood disorders: kudos and quarrels. <i>Aust NZ Psychiatry</i> 2016, 50(10):937-8. DOI:10.1177/0004867416668038.	Was not a CPG
93. Grobler, G. An overview of depression treatment guidelines. Abstracts 2nd African College of Neuropsychopharmacology Congress 30-31 July 2016 Stellenbosch, Western Cape, South Africa. <i>Acta Neuropsychiatr</i> 2016, 28(s3):1-15. DOI:10.1017/neu.2016.37.	Was not a CPG
94. Heilmann KE, Wagner M, Riedel-Heller S, <i>et al.</i> Treating late life depression with antidepressants: a summary of recommendations in international guidelines. <i>Fortschr Neurol Psychiatr</i> 2015, 83(7):381-91. DOI:10.1055/s-0035-1553315.	Was not a CPG
95. Horgan D, Dodd S. Combination antidepressants—use by GPs and psychiatrists. <i>Aust Fam Physician</i> 2011, 40(6):397-400.	Was not a CPG



Reference of the excluded CPG	Reasons for exclusion
96. KASPER, S. Editorial. World Journal of Biological Psychiatry, Glasgow, v. 14, n. 5, p. 333. DOI <a href="https://doi.org/10.3109/15622975.2013.819703">https://doi.org/ 10.3109/15622975.2013.819703</a> . Available from: <a href="https://www.tandfonline.com/doi/full/10.3109/15622975.2013.819703">https://www.tandfonline.com/doi/full/10.3109/15622975.2013.819703</a> .	Was not a CPG
97. Kendall T, Pilling S, Glover N, <i>et al</i> . Guidelines in mental health - national and international perspectives. Int Rev Psychiatry 2011, 23(4):314-7. DOI:10.3109/09540261.2011.607431.	Was not a CPG
98. Kongsuk T. Clinical practice guideline major depressive disorder for general practitioners. Value Health 2013, 16(7): A695. DOI:10.1016/j.jval.2013.08.2091.	Was not a CPG
99. KURIAN, B. T., GRANNEMANN, B., TRIVEDI, M. H. Feasible evidence-based strategies to manage depression in primary care. <b>Current Psychiatry Reports</b> , Philadelphia, v. 14, n. 4, p. 370-375. DOI: <a href="https://doi.org/10.1007/s11920-0120290-y">https://doi.org/10.1007/s11920-0120290-y</a> . Available from: <a href="https://link.springer.com/article/10.1007/s11920-012-0290-y">https://link.springer.com/article/10.1007/s11920-012-0290-y</a> .	Was not a CPG
100. Laux G. Update treatment of depression—S3 Guideline, internet-based psychotherapy, antidepressants and driving ability. Nervenheilkunde 2016, 35(10):691-6. DOI:10.1055/s-0037-1616433.	Was not a CPG
101. FACULTY OF SEXUAL AND REPRODUCTIVE HEALTHCARE OF THE ROYAL COLLEGE OF OBSTETRICIANS AND GYNEACOLOGISTS. The relationship between progestin hormonal contraception and depression: a systematic review. [London]: 6 March 2018. Available from: <a href="https://www.fsrh.org/news/fsrh-ceu-statement-response-to-study-on-progestogen-only/">https://www.fsrh.org/news/fsrh-ceu-statement-response-to-study-on-progestogen-only/</a> .	Was not a CPG
102. FACULTY OF SEXUAL AND REPRODUCTIVE HEALTHCARE OF THE ROYAL COLLEGE OF OBSTETRICIANS AND GYNEACOLOGISTS. FSRH CEU Statement: response to study on progestogen-only contraception use and depression [London]: FSRH; 8 March 2018. Available from: <a href="https://www.fsrh.org/news/fsrh-ceu-statement-response-to-study-on-progestogen-only/">https://www.fsrh.org/news/fsrh-ceu-statement-response-to-study-on-progestogen-only/</a> .	Was not a CPG
103. Leadholm AKK, Rothschild AJ, Nolen WA <i>et al</i> . The treatment of psychotic depression: is there consensus among guidelines and psychiatrists? J Affect Disord 145(2): 214-20. DOI:10.1016/j.jad.2012.07.036.	Focused on specific populations
104. Malhi G, Oakley-Browne M, Hay P. Clinical practice guidelines project (CPG project) overview. Aust NZ JPsychiatr 2015, 49(Suppl 1):30. DOI:10.1177/0004867415578344.	Was not a CPG
105. Manning JS, Jackson WC. Providing guideline-concordant assessment and monitoring for major depression in primary care. J Clin Psychiatry 2015, 76(1):e3. DOI:10.4088/JCP.13013tx7c.	Was not a CPG
106. Morris DW, Trivedi MH. Measurement-based care for unipolar depression. Curr Psychiatry Rep 2011, 13(6):446-58. DOI:10.1007/s11920-011-0237-8.	Was not a CPG

Reference of the excluded CPG	Reasons for exclusion
107. Nelson JC. Foreword. CNSDrugs 2013, 27:3-4. DOI:10.1007/s40263-012-0027-9.	Was not a CPG
108. Nutt DJ. Highlights of the international consensus statement on major depressive disorder. J Clin Psychiatry 2011, 72(6):e21. DOI:10.4088/JCP.9058tx2c.	Was not a CPG
109. Ogasawara K, Ozaki N. Review of the new treatment guideline for major depressive disorder by the Japanese Society of Mood Disorders. Brain Nerve 2012, 64(10):1159-65. In Japanese.	Was not a CPG
110. Oldham J. Fine-tuning our treatment strategies. J Psychiatr Pract 2011, 17(3):157. DOI:10.1097/01.pra.0000398408.13750.8d.	Was not a CPG
111. Pai N. Are the Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for mood disorders meeting the needs of clinicians? Aust N Z J Psychiatry 2016, 50(10):1015-6. DOI:10.1177/0004867416667828.	Was not a CPG
112. Patkar AA, Pae C-U. Atypical antipsychotic augmentation strategies in the context of guideline-based care for the treatment of major depressive disorder. CNS Drugs 2013, 27(Suppl 1):S29-S37. DOI:10.1007/s40263-012-0031-0.	Was not a CPG
113. Rawlins M. Ten years of NICE mental health guidelines. Int Rev Psychiatry 2011, 23(4):311-3. DOI:10.3109/09540261.2011.606804.	Was not a CPG
114. Reisdorf S. Revision of the national health care guidelines on unipolar depression. MedMonatss Pharm 2016, 39(4):171-2.	Was not a CPG
115. Roberge P, Fournier L, Brouillet H, <i>et al.</i> A provincial adaptation of clinical practice guidelines for depression in primary care: a case illustration of the ADAPTE method. J Eval Clin Pract 2015, 21(6):1190-8. DOI:10.1111/jep.12404.	Was not a CPG
116. Schuklenk U, Vathorst SVD. Treatment-resistant major depressive disorder and assisted dying. J Med Ethics 2015, 41(8), 577-83. DOI:10.1136/medethics-2014-102458.	Was not a CPG
117. Schulte-Körne G, Krick K. In reply. Dtsch Arztebl Int 2014, 111(18):330. DOI:10.3238/arztebl.2014.0330c.	Was not a CPG
118. Tomba E, Fava GA. Treatment selection in depression: the role of clinical judgment. Psychiatr Clin North Am 2012, 35(1):87-98. DOI:10.1016/j.psc.2011.11.003.	Was not a CPG
119. TREUER, T., <i>et al.</i> Use of antidepressants in the treatment of depression in Asia: guidelines, clinical evidence, and experience revisited. <i>Asia-Pacific Psychiatry</i> , Richmond, v. 5, n. 4, p. 219-230. DOI <a href="https://doi.org/10.1111/appy.12090">https://doi.org/10.1111/appy.12090</a> . Available from: <a href="https://onlinelibrary.wiley.com/doi/10.1111/appy.12090">https://onlinelibrary.wiley.com/doi/10.1111/appy.12090</a> .	Was not a CPG
120. Tundo A, Calabrese JR, Proietti L, <i>et al.</i> Short-term antidepressant treatment of bipolar depression: are ISBD recommendations useful in clinical practice? J Affect Disord 2015, 171:155-60. doi:10.1016/j.jad.2014.09.019.	Focused on specific populations

Reference of the excluded CPG	Reasons for exclusion
121. Van Avendonk M, van Weel-Baumgarten E, van der Weele G, <i>et al.</i> Summary of the Dutch College of General Practitioners' practice guideline "Depression". Ned Tijdschr Geneeskde 2012, 156(38):A5101.	Was not a CPG
122. Vinberg M, Levinsen MF, Kessing LV. Treatment-resistant depression is treatable. Ugeskr Laeger 2011, 173(9):651-4.	Was not a CPG
123. Xiang YT, Hu C, Wang G, <i>et al.</i> Prescribing patterns of antidepressants, antipsychotics and mood stabilizers in bipolar patients misdiagnosed with major depressive disorder in China. Hum Psychopharmacol 2012, 27(6):626-31. DOI:10.1002/hup.2262.	Was not a CPG
124. YANG, J. <i>et al.</i> Experiences and barriers to implementation of clinical practice guideline for depression in Korea. BMC Psychiatry, London, v. 13, p. 150. DOI: <a href="https://doi.org/10.1186/1471-244X-13-150">https://doi.org/10.1186/1471-244X-13-150</a> . Available from: <a href="https://bmcp psychiatry.biomedcentral.com/articles/10.1186/1471-244X-13-150">https://bmcp psychiatry.biomedcentral.com/articles/10.1186/1471-244X-13-150</a> .	Was not a CPG
125. Zimmerman M. Symptom severity and guideline-based treatment recommendations for depressed patients: implications of DSM-5's potential recommendation of the PHQ-9 as the measure of choice for depression severity. Psychother Psychosom 2012, 81(6):329-32. DOI:10.1159/000342262.	Was not a CPG
126. Zimmerman M, Martinez JH, Friedman M, <i>et al.</i> How can we use depression severity to guide treatment selection when measures of depression categorize patients differently? J Clin Psychiatry 2012, 73(10):1287-91. DOI:10.4088/JCP.12m07775.	Was not a CPG
127. Herrmann M, Gensichen J. Guidelines in general practice: the example depression. Psychiatrie 2015, 66:119-24.	Was not a CPG
128. Kuroki T, Tanaka T. [Threshold of application of antidepressant drugs for treatment of depressive disorder]. Seishin Shinkeigaku Zasshi 2015, 117(4):269-76. In Japanese.	Was not a CPG
129. Semba J. [Proper use of novel antidepressants in psychiatric clinical practice]. Seishin Shinkeigaku Zasshi. 2014, 116(2):138-143. In Japanese.	Was not a CPG
130. Sforzini, L., Worrell, C., Kose, M. <i>et al.</i> A Delphi-method-based consensus guideline for definition of treatment-resistant depression for clinical trials. Mol Psychiatry (2021). <a href="https://doi.org/10.1038/s41380-021-01381-x">https://doi.org/10.1038/s41380-021-01381-x</a> .	Did not include pharmacological treatment
131. McAllister-Williams RH, Arango C, Blier P, Demyttenaere K, Falkai P, Gorwood P, Hopwood M, Javed A, Kasper S, Malhi GS, Soares JC, Vieta E, Young AH, Papadopoulos A, Rush AJ. The identification, assessment and management of difficult-to-treat depression: An international consensus statement. J Affect Disord. 2020 Apr 15; 267:264-282. DOI: 10.1016/j.jad.2020.02.023. Epub 2020 Feb 7. PMID: 32217227.	Was not a CPG

Reference of the excluded CPG	Reasons for exclusion
132. Lu, S., Li, W., Oldenburg, B. <i>et al.</i> Cultural adaptation of the mental health first aid guidelines for depression used in English-speaking countries for China: a Delphi expert consensus study. <i>BMC Psychiatry</i> 20, <b>336</b> (2020). <a href="https://doi.org/10.1186/s12888-020-02736-4">https://doi.org/10.1186/s12888-020-02736-4</a> .	Was not a CPG
133. Taylor RW, Marwood L, Oprea E, DeAngel V, Mather S, Valentini B, Zahn R, Young AH, Cleare AJ. Pharmacological Augmentation in Unipolar Depression: A Guide to the Guidelines. <i>Int J Neuropsychopharmacol.</i> 2020 Dec 3, 23(9):587-625. DOI: 10.1093/ijnp/pyaa033. PMID: 32402075; PMCID: PMC7710919.	Was not a CPG
134. Andrade C. Oral racemic ketamine for common clinical contexts in patients with major depressive disorder: An important intervention that treatment guidelines may never include. <i>Bipolar Disord.</i> 2022 Mar, 24(2):113-114. DOI: 10.1111/bdi.13155. Epub 2021 Nov 23. PMID: 34816542.	Was not a CPG
135. CANADIAN AGENCY FOR DRUGS AND TECHNOLOGIES IN HEALTH. Accessing mental health support outside of a therapist's office: iCBT for major depression and anxiety disorders. [Ottawa, ON]: CADTH, [2020]. Available from: <a href="https://www.cadth.ca/events/evidence-action-accessing-mental-health-support-outside-therapists-office-icbt-major">https://www.cadth.ca/events/evidence-action-accessing-mental-health-support-outside-therapists-office-icbt-major</a> .	Subject matter
136. NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE. Common mental health problems: identification and pathways to care. <b>London: NICE, 2011. Available from:</b> <a href="https://www.nice.org.uk/guidance/cg123">https://www.nice.org.uk/guidance/cg123</a> .	Subject matter
137. HAUTE AUTORITÉ DE SANTÉ. Dangerosité psychiatrique: étude et évaluation des facteurs de risque de violence hétéro-agressive chez les personnes ayant des troubles schizophréniques ou des troubles de l'humeur. Saint-Denis, 2011. Available from: <a href="https://www.has-sante.fr/jcms/c_1069220/fr/dangerosite-psychiatrique-etude-et-evaluation-des-facteurs-de-risque-de-violence-hetero-agressive-chez-les-personnes-ayant-des-troubles-schizophreniques-ou-des-troubles-de-l-humeur">https://www.has-sante.fr/jcms/c_1069220/fr/dangerosite-psychiatrique-etude-et-evaluation-des-facteurs-de-risque-de-violence-hetero-agressive-chez-les-personnes-ayant-des-troubles-schizophreniques-ou-des-troubles-de-l-humeur</a> .	Subject matter
138. Arbejdsgruppen vedr. den nationale kliniske retningslinje for farmakologisk behandling af bipolar lidelse. National klinisk retningslinje for farmakologisk behandling af bipolar lidelse: supplerende vedligeholdelsesbehandling efter depression. Version 27. maj 2014 udg. København: Sundhedsstyrelsen, 2014. 111 s.	Subject matter
139. Shelton RC, Trivedi MH. Using algorithms and computerized decision support systems to treat major depression. <i>J Clin Psychiatry.</i> 2011 Dec, 72(12):e36. DOI: 10.4088/JCP.10027tx4cc. PMID: 22244032.	Was not a CPG
140. Lam RW, Parikh SV, Michalak EE, Dewa CS, Kennedy SH. Canadian Network for Mood and Anxiety Treatments (CANMAT) consensus recommendations for functional outcomes in major depressive disorder. <i>Ann Clin Psychiatry.</i> 2015 May, 27(2):142-9. PMID: 25954941.	Subject matter



Reference of the excluded CPG	Reasons for exclusion
141. Human growth hormone for depression, post-traumatic stress disorder, or fibromyalgia: Clinical effectiveness and guidelines. Ottawa: CADTH, 2017 Feb. (CADTH rapid response report: reference list).	Subject matter
142. Yamada K. [Evidence of treatment for depressive episodes of bipolar disorder]. Seishin Shinkeigaku Zasshi 2011, 113(9):873-9. In Japanese.	Subject matter
143. Depression. Adapted Evidence-based Guideline. Ukraine, 2014.	Was not a CPG
144. Siu AL, US Preventive Services Task Force (USPSTF). Screening for depression in adults: US Preventive Services Task Force recommendation statement. JAMA 2016, 315(4):380-7. DOI:10.1001/jama.2015.18392.	Did not include pharmacological treatment
145. Thase ME. Recommendations for screening for depression in adults. JAMA 2016, 315(4):349-50. DOI:10.1001/jama.2015. 18406.	Did not include pharmacological treatment
146. CANADIAN AGENCY FOR DRUGS AND TECHNOLOGIES IN HEALTH. Internet-Delivered Cognitive Behavioural Therapy (iCBT) programs for major depression and anxiety disorders in Canada. Ottawa, ON: CADTH, 2020. Available from: <a href="https://www.cadth.ca/internet-delivered-cognitive-behavioural-therapy-icbt-programs-major-depression-and-anxiety">https://www.cadth.ca/internet-delivered-cognitive-behavioural-therapy-icbt-programs-major-depression-and-anxiety</a>	Did not include pharmacological treatment
147. Guided Versus Unguided Internet-Delivered Cognitive Behavioural Therapy for Major Depressive Disorder and Anxiety Disorders: Comparative Clinical Effectiveness. Ottawa: CADTH, 2018 Oct. (CADTH rapid response report: summary of abstracts).	Did not include pharmacological treatment
148. Pohar R, Arg��ez C. Acceptance and Commitment Therapy for Post-Traumatic Stress Disorder, Anxiety, and Depression: A Review of Clinical Effectiveness [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2017 Aug 28. PMID: 30226695.	Did not include pharmacological treatment
149. CANADIAN AGENCY FOR DRUGS AND TECHNOLOGIES IN HEALTH. Mindfulness interventions for the treatment of post-traumatic stress disorder, generalized anxiety disorder, depression, and substance use disorders: a review of the clinical effectiveness and guidelines. Ottawa, ON: CADTH, 2015. Available from: <a href="https://www.cadth.ca/mindfulness-interventions-treatment-post-traumatic-stress-disorder-generalized-anxiety-disorder">https://www.cadth.ca/mindfulness-interventions-treatment-post-traumatic-stress-disorder-generalized-anxiety-disorder</a> .	Did not include pharmacological treatment
150. Yoga for the Treatment of Post-Traumatic Stress Disorder, Generalized Anxiety Disorder, Depression, and Substance Abuse: A Review of the Clinical Effectiveness and Guidelines [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2015 Jun 22. Available from: <a href="https://www.ncbi.nlm.nih.gov/books/NBK304560/">https://www.ncbi.nlm.nih.gov/books/NBK304560/</a> .	Did not include pharmacological treatment

Reference of the excluded CPG	Reasons for exclusion
151. Health Quality Ontario. Internet-Delivered Cognitive Behavioural Therapy for Major Depression and Anxiety Disorders: A Health Technology Assessment. Ont Health Technol Assess Ser. 2019- Feb 19, 19(6):1-199. PMID: 30873251; PMCID: PMC6394534.	Did not include pharmacological treatment
152. Bright Light Therapy for Depression: Clinical Effectiveness. Ottawa: CADTH, 2019 Aug. (CADTH rapid response report: summary of abstracts).	Did not include pharmacological treatment
153. Physical activity and support programs for mild to moderate major depressive disorder. (CADTH reference list). Ottawa: CADTH, 2021 November.	Did not include pharmacological treatment
154. CANADIAN AGENCY FOR DRUGS AND TECHNOLOGIES IN HEALTH. Diagnosing, screening, and monitoring depression in the elderly: a review of guidelines. Ottawa, ON: CADTH, 2015.	Did not include pharmacological treatment
155. CANADIAN AGENCY FOR DRUGS AND TECHNOLOGIES IN HEALTH. Stellate ganglion block for the treatment of post-traumatic stress disorder, depression, and anxiety. Ottawa, ON: CADTH, 2021. Available from: <a href="https://www.cadth.ca/stellate-ganglion-block-treatment-post-traumatic-stress-disorder-depression-and-anxiety">https://www.cadth.ca/stellate-ganglion-block-treatment-post-traumatic-stress-disorder-depression-and-anxiety</a> .	Did not include pharmacological treatment
156. Pharmacogenomic testing in depression: a review of clinical effectiveness, cost-effectiveness, and guidelines. Ottawa: CADTH, 2020 Jan. (CADTH rapid response report: summary with critical appraisal).	Did not include pharmacological treatment
157. CANADIAN AGENCY FOR DRUGS AND TECHNOLOGIES IN HEALTH. <b>Repetitive transcranial magnetic stimulation for depression</b> : a review of the clinical effectiveness, cost-effectiveness, and guidelines. Ottawa, ON: CADTH, 2021. Available from: <a href="https://www.cadth.ca/repetitive-transcranial-magnetic-stimulation-depression-review-clinical-effectiveness-cost">https://www.cadth.ca/repetitive-transcranial-magnetic-stimulation-depression-review-clinical-effectiveness-cost</a> .	Did not include pharmacological treatment
158. Health Quality Ontario. Internet-Delivered Cognitive Behavioural Therapy for Major Depression and Anxiety Disorders: A Health Technology Assessment. Ont Health Technol Assess Ser. 2019 Feb 19, 19(6):1-199. PMID: 30873251; PMCID: PMC6394534.	Did not include pharmacological treatment
159. NATIONAL KLINISK RETNINGSLINJE FOR NON-FARMAKOLOGISK BEHANDLING AF UNIPOLAR DEPRESSION. Available from: <a href="https://www.sst.dk/da/udgivelser/2016/~media/0216564bb8aa4d40b7dbaf21e9acf403.ashx">https://www.sst.dk/da/udgivelser/2016/~media/0216564bb8aa4d40b7dbaf21e9acf403.ashx</a> .	Did not include pharmacological treatment
160. Joffres M, Jaramillo A, Dickinson J, <i>et al</i> . Recommendations on screening for depression in adults [published correction appears in CMAJ. 2013 Sep 3, 185(12):1067]. CMAJ. 2013;185(9):775-782. DOI:10.1503/cmaj.130403.	Did not include pharmacological treatment

Reference of the excluded CPG	Reasons for exclusion
161. Andersen, B. L., DeRubeis, R. J., Berman, B. S., Gruman, J., Champion, V. L., Massie, M. J., Holland, J. C., Partridge, A. H., Bak, K., Somerfield, M. R., Rowland, J. H., & American Society of Clinical Oncology (2014). Screening, assessment, and care of anxiety and depressive symptoms in adults with cancer: an American Society of Clinical Oncology guideline adaptation. <i>Journal of clinical oncology: official journal of the American Society of Clinical Oncology</i> , 32(15), 1605–1619. <a href="https://doi.org/10.1200/JCO.2013.52.4611">https://doi.org/10.1200/JCO.2013.52.4611</a>	Did not include pharmacological treatment
162. Implanted vagus nerve stimulation for treatment-resistant depression. Available from: <a href="https://www.nice.org.uk/guidance/ipg679">https://www.nice.org.uk/guidance/ipg679</a> .	Did not include pharmacological treatment
163. FROST, J. <i>et al.</i> Depression following acute coronary syndrome events: screening and treatment guidelines from the AAFP. <i>American Family Physician</i> , Kansas City, MO, v. 99, n. 12, online. Available from: <a href="https://www.aafp.org/afp/2019/0615/od2.html">https://www.aafp.org/afp/2019/0615/od2.html</a> .	Focused on specific populations
164. Noe KH, Locke DE, Sirven JI. Treatment of depression in patients with epilepsy. <i>Curr Treat Options Neurol</i> . 2011 Aug;13(4):371-9. DOI: 10.1007/s11940-011-0127-8. PMID: 21472437.	Focused on specific populations
165. Rayner L, Price A, Hotopf M, <i>et al.</i> The development of evidence-based European guidelines on the management of depression in palliative cancer care. <i>Eur J Cancer</i> 2011, 47(5):702-12. DOI:10.1016/j.ejca.2010.11.027.	Focused on specific populations
166. Williams D, Mierzwinski-Urban M. Adapted or Tailored Psychological Interventions for Treating Women with Mental Illness: A Review of Clinical Effectiveness and Guidelines [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2017 Oct 5. Available from: <a href="https://www.ncbi.nlm.nih.gov/books/NBK525628/">https://www.ncbi.nlm.nih.gov/books/NBK525628/</a> .	Focused on specific populations
167. WORLD HEALTH ORGANIZATION. <b>Pharmacological interventions (antidepressants) for 433 people with dementia who have associated depression</b> . Geneva: WHO, 2015. Available from: <a href="https://www.who.int/mental_health/mhgap/evidence/emmentia/q4/en/">https://www.who.int/mental_health/mhgap/evidence/emmentia/q4/en/</a> .	Focused on specific populations
168. Austin M-P, Highet N, The Expert Working Group. Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline. Melbourne, Australia: Centre of Perinatal Excellence, 2017.	Focused on specific populations
169. Colquhoun DM, Bunker SJ, Clarke DM, <i>et al.</i> Screening, referral and treatment for depression in patients with coronary heart disease. <i>Med J Aust</i> 2013, 198(9):483-4. DOI:10.5694/mja13.10153.	Focused on specific populations
170. Perinatal Depression Screening, Diagnosis, and treatment Guideline. Available from: <a href="https://wa.kaiserpermanente.org/static/pdf/public/guidelines/depression-perinatal.pdf">https://wa.kaiserpermanente.org/static/pdf/public/guidelines/depression-perinatal.pdf</a> .	Focused on specific populations

Reference of the excluded CPG	Reasons for exclusion
171. Andersen, B. L., DeRubeis, R. J., Berman, B. S., Gruman, J., Champion, V. L., Massie, M. J., Holland, J. C., Partridge, A. H., Bak, K., Somerfield, M. R., Rowland, J. H., & American Society of Clinical Oncology (2014). Screening, assessment, and care of anxiety and depressive symptoms in adults with cancer: an American Society of Clinical Oncology guideline adaptation. <i>Journal of clinical oncology: official journal of the American Society of Clinical Oncology</i> , 32(15), 1605–1619. <a href="https://doi.org/10.1200/JCO.2013.52.4611">https://doi.org/10.1200/JCO.2013.52.4611</a> .	Focused on specific populations
172. Li M, Kennedy EB, Byrne N, Gérin-Lajoie C, Katz MR, Keshavarz H, Sellick S, Green E. Management of Depression in Patients With Cancer: A Clinical Practice Guideline. <i>J Oncol Pract</i> . 2016 Aug. 12, (8):747-56. DOI: 10.1200/JOP.2016.011072. Epub 2016 Jul 5. Erratum in: <i>J Oncol Pract</i> . 2017 Feb;13(2):144. PMID: 27382000.	Focused on specific populations
173. NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE. Depression in adults with a chronic physical health problem: recognition and management. London: NICE, 2009. Available from: <a href="https://www.nice.org.uk/guidance/cg91">https://www.nice.org.uk/guidance/cg91</a> .	Focused on specific populations
174. MINISTERIO DE SALUD. Guía Clínica: Abordaje de la depresión en mujeres con cáncer de mama. Santiago: Minsal, 2017. Available from: <a href="https://diprece.minsal.cl/wrdprss_minsal/wp-content/uploads/2017/07/GPC_depresion_en_cancer_de_mama.pdf">https://diprece.minsal.cl/wrdprss_minsal/wp-content/uploads/2017/07/GPC_depresion_en_cancer_de_mama.pdf</a> .	Focused on specific populations
175. Concurrent Treatment for Substance Use Disorder and Trauma-Related Comorbidities: 2021 Update. Available from: <a href="https://www.cadth.ca/concurrent-treatment-substance-use-disorder-and-trauma-related-comorbidities-2021-update">https://www.cadth.ca/concurrent-treatment-substance-use-disorder-and-trauma-related-comorbidities-2021-update</a> .	Focused on specific populations
176. US Preventive Services Task Force. Interventions to prevent perinatal depression. <i>JAMA</i> , Chicago, v. 321, n. 6, p. 580-587. DOI <a href="http://doi.org/10.1001/jama.2019.0007">http://doi.org/10.1001/jama.2019.0007</a> .	Focused on specific populations
177. Galician Health Technology Assessment Agency. Clinical Practice Guideline on the Management of Major Depression in Adults. Santiago de Compostela, Spain: Galician Health Technology Assessment Agency, 2008.	Were outdated
178. National Collaborating Centre for Mental Health (UK). Depression. The Treatment and Management of Depression in Adults. Leicester, UK: British Psychological Society, 2010.	Were outdated
179. Agency for Healthcare Research and Quality. Detecting Depression in Older Adults with Dementia. Ann Arbor, MI: University of Michigan Health System; [s.d].	Focused on specific populations
180. Adult Depression in Primary Care. Bloomington, MN: ICSI, 2013.	Were outdated
181. Galletly C, Castle D, Dark F, <i>et al</i> . Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for the management of schizophrenia and related disorders. <i>Aust N Z J Psychiatry</i> 2016, 50(5):410-72. DOI:10.1177/000 4867416641195.	Subject matter



Reference of the excluded CPG	Reasons for exclusion
182. Malhi GS, Bassett D, Boyce P, <i>et al.</i> Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for mood disorders. Aust N Z J Psychiatry 2015, 49(12):1087-206. DOI:10.1177/0004867415617657.	Were outdated
183. Karyotaki E, Smit Y, Cuijpers P, <i>et al.</i> The Long-term Efficacy of Psychotherapy, Alone or in Combination with Antidepressants, in the Treatment of Adult Major Depression. Good Clinical Practice (GCP). Brussels, Belgium: Belgian psychotherapy Health Care Knowledge Centre (KCE), 2014. Available from: <a href="#">pression_Report_0.pdf</a> .	Focused on psychotherapy
184. MINISTERIO DE SALUD. Guía Clínica: Para el tratamiento de la Depresión en personas mayores de 15 años: Actualización en Psicoterapia. Santiago: MINSAL, (2017). Todos los derechos reservados. Este material puede ser reproducido total o parcialmente para fines de diseminación y capacitación. Prohibida su venta. ISBN: (número de registro) 1ª Edición: Santiago, 2017. Available from: <a href="https://diprece.minsal.cl/wrdprss_minsal/wp-content/uploads/2017/07/GPC_depresion_psicoterapia.pdf">https://diprece.minsal.cl/wrdprss_minsal/wp-content/uploads/2017/07/GPC_depresion_psicoterapia.pdf</a> .	Focused on psychotherapy
185. NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE. <b>Vortioxetine for treating major depressive episodes</b> . London: NICE, 2015. Available from: <a href="https://www.nice.org.uk/guidance/ta367/resources/vortioxetine-for-treating-major-depressive-episodes-pdf-82602733813189#:~:text=Vortioxetine%20(Brintellix%2C%20Lundbeck)%20is,major%20depressive%20episodes%20in%20adults">https://www.nice.org.uk/guidance/ta367/resources/vortioxetine-for-treating-major-depressive-episodes-pdf-82602733813189#:~:text=Vortioxetine%20(Brintellix%2C%20Lundbeck)%20is,major%20depressive%20episodes%20in%20adults</a> '.	Specific drugs
186. NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE. <b>Agomelatine for the treatment of major depressive episodes(terminated appraisal)</b> . London: NICE, 2011. Available from: <a href="https://www.nice.org.uk/guidance/ta231/resources/agomelatine-for-the-treatment-of-major-depressive-episodes-terminated-appraisal-pdf-82600_365554629">https://www.nice.org.uk/guidance/ta231/resources/agomelatine-for-the-treatment-of-major-depressive-episodes-terminated-appraisal-pdf-82600_365554629</a> .	Specific drugs
187. Sanacora G, Frye MA, McDonald W, Mathew SJ, Turner MS, Schatzberg AF, Summergrad P, Nemeroff CB; American Psychiatric Association (APA) Council of Research Task Force on Novel Biomarkers and Treatments. A Consensus Statement on the Use of Ketamine in the Treatment of Mood Disorders. JAMA Psychiatry. 2017 Apr 1, 74(4):399-405. DOI: 10.1001/jamapsychiatry.2017.0080. PMID: 28249076.	Specific drugs
188. Tran K, Argáez C. Quetiapine for Major Depressive Disorder: A Review of Clinical Effectiveness, Cost-Effectiveness, and Guidelines [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2020 Jan 30. PMID: 33074605.	Specific drugs
189. Esketamine for Treatment-Resistant Depression. Available from: <a href="https://www.hopkinsmedicine.org/health/treatment-tests-and-therapies/esketamine-for-treatment-resistant-depression">https://www.hopkinsmedicine.org/health/treatment-tests-and-therapies/esketamine-for-treatment-resistant-depression</a> ketamine for Treatment-Resistant Depression.	Specific drugs

Reference of the excluded CPG	Reasons for exclusion
190. Tran K, McGill SC, Horton J. Bupropion for Treatment-Resistant Depression [Internet]. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2021 Apr. Available from: <a href="https://www.ncbi.nlm.nih.gov/books/NBK571945/">https://www.ncbi.nlm.nih.gov/books/NBK571945/</a> .	Specific drugs
191. Antidepressants for Adults with Depression using Opioid Medications: Clinical Effectiveness and Guidelines. Ottawa: CADTH; 2018 Feb. (CADTH rapid response report: summary of abstracts). Available from: <a href="https://www.cadth.ca/sites/default/files/pdf/htis/2018/RB1196%20Antidepressants%20with%20opioid%20use%20Final.pdf">https://www.cadth.ca/sites/default/files/pdf/htis/2018/RB1196%20Antidepressants%20with%20opioid%20use%20Final.pdf</a>	Was not a CPG

## Supplemental Material 3 – Results

**Table 1. Appraisal of CPGs for Research & Evaluation Instrument AGREE II domain scores (%) and AGREE-REX domain scores (%).**

Country / Study	AGREE II domains' scores						AGREE-REX domains' scores		
	1. Scope and purpose	2. Stakeholder involvement	3. Rigour of development	4. Clarity of presentation	5. Applicability	6. Editorial independence	1. Clinical applicability	2. Values and preferences	3. Implementability
Chile, Ministerio de Salud [23]	83	76	89	94	57	17	33	29	41
Colombia, Ministerio de Salud [24]	100	85	86	100	96	92	72	57	67
United Kingdom, National Institute for Health and Care Excellence [25]	89	83	84	81	71	75	87	67	75
Germany, Härter et al. [26]	69	78	83	87	58	72	61	42	64
Peru EsSalud [27]	72	52	82	87	39	67	67	36	50
USA, Trangle et al. [28]	96	78	81	91	72	97	55	36	50
USA, American Psychological Association [29]	91	67	81	80	65	83	81	44	50
USA, Management of Major Depressive Disorder Working Group [30]	93	76	78	94	38	58	67	31	56
USA, Kaiser Permanente Care Management Institute [31]	83	63	76	93	46	58	54	25	33
Spain, García-Herrera Pérez Bryan et al. [32]	74	59	72	85	64	75	52	26	61
Spain, Working Group of the Clinical Practice Guideline on the Management of Depression in Adults, Ministry of Health, Social Services and Equality [33]	94	93	70	91	75	53	67	36	67
Canada, Registered Nurses' Association of Ontario [34]	72	74	69	80	76	86	57	42	67
USA, Qaseem et al. [35]	80	39	69	70	32	67	48	11	33
Mexico, Instituto Mexicano del Seguro Social [36]	87	46	69	83	14	67	48	11	42
Mexico, Secretaría del Salud [37]	81	43	69	80	32	31	43	11	50
Mexico, Secretaría del Salud [38]	94	56	63	81	42	64	43	15	42
Singapore, Chua et al. [39]	78	72	60	89	50	28	59	19	44
Australia, Malhi et al. [40]	74	63	58	78	24	67	50	22	44

Country / Study	AGREE II domains' scores						AGREE-REX domains' scores		
	1. Scope and purpose	2. Stakeholder involvement	3. Rigour of development	4. Clarity of presentation	5. Applicability	6. Editorial independence	1. Clinical applicability	2. Values and preferences	3. Implementability
Michigan Ambulatory Unipolar Depression Guideline [41]	65	43	57	80	33	53	41	28	31
France, Driot et al. [42]	69	30	56	72	11	83	35	4	33
Several countries, Bauer et al. [43]	61	54	54	83	32	75	39	11	22
Canada, Kennedy et al. [44]	63	48	54	89	26	53	44	17	36
Several countries, Dua et al. [45]	69	74	50	74	29	75	41	25	47
USA, McIntyre et al. [46]	87	56	48	83	32	69	39	21	33
Several countries, Bauer et al. [47]	69	48	47	61	28	75	48	15	19
Malaysia, Malaysian Health Technology Assessment Section [48]	81	50	47	70	54	78	59	33	67
USA, Gelenberg et al. [49]	48	43	46	83	44	42	59	32	42
The Netherlands, Nederland Depressie [50]	70	70	42	65	22	50	28	19	33
United Kingdom, Cleare et al. [51]	67	57	40	69	13	58	52	22	33
Korea, Won et al. [52]	57	28	38	67	21	44	35	14	42
Finland, Suomalainen Lääkäriseura Duodecim [53]	50	61	36	65	40	56	44	21	50
USA, Ruberto et al. [54]	43	11	35	39	1	72	15	4	11
Canada, CPGs and Protocols Advisory Committee, Ministry of Health, British Columbia [55]	85	37	35	85	39	42	37	21	44
USA, Giakoumatos and Osseir [56]	61	19	33	83	26	75	30	21	14
France, Bennabi et al. [57]	72	26	31	85	10	81	20	8	25
Saudi Arabia, Okasha et al. [58]	44	26	31	61	31	33	37	11	31
Several countries, Dodd S, Mitchell PB, Bauer et al. [59]	72	31	27	57	17	25	41	22	31
Several countries, Bauer et al. [60]	56	41	23	76	21	50	30	8	36
France, Bennabi et al. [61]	50	33	22	65	13	67	39	26	25
Pharmacological Management of Depression: Japanese Expert Consensus [62]	50	31	21	59	31	56	20	24	25
South Africa, Emsley [63]	50	48	19	67	13	19	26	12	33
USA, Connolly and Thase [64]	63	17	17	52	13	72	15	6	14



Country / Study	AGREE II domains' scores						AGREE-REX domains' scores		
	1. Scope and purpose	2. Stakeholder involvement	3. Rigour of development	4. Clarity of presentation	5. Applicability	6. Editorial independence	1. Clinical applicability	2. Values and preferences	3. Implementability
Korea, Wang et al. [65]	56	13	17	43	6	58	19	10	36
USA, Park and Zarate [66]	33	22	17	50	18	31	17	13	14
Canada, Voineskos et al. [67]	44	11	15	50	10	22	15	1	17
USA, Voytenko et al. [68]	54	39	15	65	8	42	17	8	22
Poland, Piotrowski et al. [69]	54	26	15	72	25	50	33	6	14
Australia, Bayes and Parker [70]	46	22	14	48	7	33	11	1	11
Australia, Malhi et al. [71]	44	20	13	63	17	39	17	7	25
France, Doumy et al. [72]	39	9	13	39	8	36	9	3	22
Canada, Mulsant et al. [73]	50	28	13	61	8	36	15	11	31
Depresja oporna na leczenie – zalecenia Konsultanta Krajowego w dziedzinie psychiatrii [74]	52	22	12	65	36	3	24	8	31
India, Avasthi and Grover [75]	70	24	12	80	36	0	22	7	31
Several countries, Möller et al. [76]	28	15	12	11	10	33	19	14	25
France, Charpeaud et al. [77]	33	13	10	46	10	36	13	0	19
USA, Busch and Sandberg [78]	46	11	10	65	15	17	19	6	25
USA, Mathys and Mitchell [79]	41	19	8	37	6	58	18	3	19
USA, Taylor [80]	41	7	8	57	8	33	22	7	33
Austria, Gartlehner et al. [81]	37	26	8	33	14	17	18	7	25
Spain, Pereira Sanchez and Santos [82]	54	24	6	61	8	33	15	7	22
India, Gautam et al. [83]	39	20	6	57	15	0	22	6	25
USA, Halaris [84]	41	17	6	31	13	11	18	5	30
Poland, Patejuk-Mazurek [85]	15	6	5	43	17	0	7	1	19

## Reference list

1. Ministério da Saúde (BR). *Diretrizes metodológicas: elaboração de diretrizes clínicas* [Internet]. 2. ed. rev., ampl., atual. Brasília, DF: 2020. Available from: <http://conitec.gov.br/images/Relatorios/2021/Diretrizes-Metodologicas->

- [Elaboracao-de-Diretrizes-Clinicas-2020.pdf](#). Accessed March 03, 2021.
2. Fervers B, Burgers JS, Haugh MC, et al. Predictors of high-quality clinical practice guidelines: examples in oncology. *Int J Qual Heal Care*. 2005;17(2):123-132.
  3. Guidelines International Network. *G-I-N PUBLIC Toolkit: Patient and Public Involvement in Guidelines*. Available from: <http://www.gin.net/activities/gin-public/toolkit>.
  4. Molino CGRC, Leite-Santos NC, Gabriel FC, et al. Factors associated with high-quality CPGs for the pharmacologic management of chronic diseases in primary care: a systematic review. *JAMA Intern Med*. 2019;179(4):553-560. doi:10.1001/jamainternmed.2018.7529
  5. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan: a web and mobile app for systematic reviews. *Syst Rev* 2016;5:210. doi:[10.1186/s13643-016-0384-4](https://doi.org/10.1186/s13643-016-0384-4).
  6. chronide.org [Internet]. [place unknown]: Chronide; c2019. Available from: <https://www.chronide.org/>. Accessed Jun 25, 2021.
  7. AGREE Next Steps Consortium. *The AGREE II Instrument* [Electronic version]. [place unknown]: AGREE; 2017. Available from: <http://www.agreetrust.org>. Accessed May 7, 2018.
  8. AGREE-REX Research Team. *The appraisal of CPGs research & evaluation-recommendation excellence (AGREE-REX)* [Internet]. [place unknown]: AGREE; 2019. Available from: <https://www.agreetrust.org/wp-content/uploads/2019/04/AGREE-REX-2019.pdf>. Accessed May 17, 2018.
  9. Institute of Medicine (US). *Clinical practice CPGs we can trust*. Washington, DC: National Academies Press; 2011. doi:[10.17226/13058](https://doi.org/10.17226/13058)
  10. Zafra-Tanaka JH, Goicochea-Lugo S, Villarreal-Zegarra D, Taype-Rondan A. Characteristics and quality of clinical practice CPGs for depression in adults: A scoping review. *BMC Psychiatry*. 2019;19(1):76. doi:10.1186/s12888-019-2057-z
  11. Gabriel FC, Stein AT, Melo DO, Fontes-Mota GCH, Santos IB, Rodrigues CDS, et al. Recommendations for the pharmacological treatment of treatment-resistant depression: A systematic review protocol. *PLoS One*. 2022; 17(4): e0267323. doi:[10.1371/journal.pone.0267323](https://doi.org/10.1371/journal.pone.0267323)

12. Ministério da Saúde (BR). *Portaria no. 1.083, de 2 de outubro de 2012. Aprova o protocolo clínico e diretrizes terapêuticas da dor crônica* [Internet]. Brasília, DF: Diário Oficial da União; 2012 [cited 2019 Jan 22]. Available from: <http://portal.arquivos2.saude.gov.br/images/pdf/2014/abril/02/pcdt-dor- cronica-2012.pdf>. Accessed June 22, 2019.
13. Ministério da Saúde (BR). *Portaria no. 1.266, de 14 de novembro de 2014. Aprova o protocolo clínico e diretrizes terapêuticas da doença de Gaucher*. Brasília, DF: Diário Oficial da União; 2014 [cited 2017 Apr 5]. Available from: <http://portal.arquivos.saude.gov.br/images/pdf/2014/dezembro/15/Doen--a-de-Gaucher.pdf>. Accessed April 5, 2017.
14. National Institute for Health and Care Excellence. *Obesity prevention: clinical guideline* [Internet]. [London]: NICE; 13 Dec 2006 [last updated 1 Mar 2015]. Available from: <https://www.nice.org.uk/guidance/cg43/resources/obesity-prevention-975445344709>. Accessed April 5, 2017.
15. Bahn RS, Burch HB, Cooper D, Garber JR, Greenlee MC, Klein I, et al. Hyperthyroidism and other causes of thyrotoxicosis: management guidelines of the American thyroid association and American association of clinical endocrinologists. *Thyroid* 2011;21(6):593-646. doi:[10.1089/thy.2010.0417](https://doi.org/10.1089/thy.2010.0417)
16. Urinary Tract Infection Guideline Team. *Urinary tract infection*. Ann Arbor (MI): University of Michigan Health System; 2011 Jun.
17. *agreetrust.org* [Internet]. [place unknown]: The AGREE Research Trust; c2010. Available from: <https://www.agreetrust.org>
18. Gabriel FC, Melo DO, Fráguas R, Leite-Santos NC, Mantovani da Silva RA, Ribeiro E. Pharmacological treatment of depression: a systematic review comparing clinical practice guideline recommendations. *PLoS One*. 2020;15(4):e0231700. doi:[10.1371/journal.pone.0231700](https://doi.org/10.1371/journal.pone.0231700)
19. The ADAPTE collaboration. *The ADAPTE process: resource toolkit for guideline adaptation*. [place unknown]: ADAPTE Collaboration, 2009. Available from: <https://www.g-i-n.net/document-store/working-groups-documents/adaptation/adapte-resource-toolkit- guideline-adaptation-2-0.pdf>. Accessed Apr 3, 2017.
20. Acuña-Izcaray A, Sánchez-Angarita E, Plaza V, Rodrigo G, Montes de Oca M, Gich I, et al. Quality assessment of asthma clinical practice guidelines: a systematic appraisal. *Chest*. 2013;144:390-7. doi:[10.1378/chest.12-2005](https://doi.org/10.1378/chest.12-2005).

21. Arevalo-Rodriguez I, Pedraza OL, Rodríguez A, Sánchez E, Gich I, Solà I, et al. Alzheimer's disease dementia guidelines for diagnostic testing: a systematic review. *Am J Alzheimers Dis Other Demen*. 2013;28:111-9. doi:10.1177/1533317512470209
22. Watine J, Friedberg B, Nagy E, Onody R, Oosterhuis W, Bunting PS, et al. Conflict between guideline methodologic quality and recommendation validity: a potential problem for practitioners. *Clin Chem*. 2006;52:65-72. doi:10.1177/1533317512470209
23. Ministerio de Salud (CL). *Depresión en personas de 15 años y más*. Santiago: Minsal; 2013.
24. Ministerio de Salud (COL). *Detección temprana y diagnóstico del episodio depresivo y trastorno depresivo recurrente en adultos: atención integral de los adultos con diagnóstico de episodio depresivo o trastorno depresivo recurrente: Guía de práctica clínica*. Bogotá: Minsalud; 2013. Available from: <http://www.iets.org.co/reportes-iets/Documentacin%20Reportes/Gu%C3%ADa.Completa.Depresion.2013.pdf>. Accessed June 30, 2017.
25. National Institute for Health and Care Excellence. *Depression in adults: recognition and management*. [London]: NICE; 2009 [cited 2017 Jun 30]. Available from: <https://www.nice.org.uk/guidance/cg90/evidenc>. Accessed June 30, 2017.
26. Härter M, Schorr S, Schneider F, eds. *S3-Leitlinie: nationale versorgungsleitlinie unipolare depression langfassung*. 2nd ed. Berlin: Springer; 2015. doi:10.1007/978-3-662-52906-5
27. Instituto de Evaluación de Tecnologías en Salud e Investigación (PE). *Guía de práctica clínica para el tamizaje y el manejo del episodio depresivo leve en el primer nivel de atención*. Lima: EsSalud; 2019. Available from: [http://www.essalud.gob.pe/ietsi/pdfs/tecnologias\\_sanitarias/GPC\\_Depresion\\_V\\_Extensa.pdf](http://www.essalud.gob.pe/ietsi/pdfs/tecnologias_sanitarias/GPC_Depresion_V_Extensa.pdf). Accessed June 30, 2017.
28. Trangle M, Gursky J, Haight R, Hinnenkamp T, Kessler D, Mack N, et al. *Depression in primary care*. Bloomington, MN: ICSI; updated March, 2016. Available from: [https://www.icsi.org/CPGs\\_more/catalog\\_CPGs\\_and\\_more/catalog\\_CPGs/catalog\\_behavioral\\_health\\_CPGs/depression](https://www.icsi.org/CPGs_more/catalog_CPGs_and_more/catalog_CPGs/catalog_behavioral_health_CPGs/depression). Accessed June 30, 2017.

29. American Psychological Association. *Clinical practice guideline for the treatment of depression across three age cohorts*. Washington, DC: APA; 2019 [cite 2017 Jun 30]. Available from: <https://www.apa.org/depression-guideline/guideline.pdf>. Accessed June 30, 2017.
30. Management of Major Depressive Disorder Working Group. *VA/DoD clinical practice guideline for the management of major depressive disorder*. 3rd ed. Washington, DC: US Department of Veterans Affairs, US Department of Defense; 2016.
31. Kaiser Permanente Care Management Institute. *Diagnosis and treatment of depression in adults: 2012 clinical practice guideline*. Oakland (CA): KPCMI; 2012.
32. García-Herrera Pérez Bryan JM, Nogueras Morilla EV, Muñoz Cobos M, Morales Asencio JM. *Guía de práctica clínica para el tratamiento de la depresión en atención primaria*. Málaga: Hospital Regional Universitario “Carlos Haya”; 2011.
33. Working Group of the Clinical Practice Guideline on the Management of Depression in Adults (ES). *Guideline on the management of depression in adults*. [Madrid]: Ministry of Health, Social Services and Equality; 2014.
34. Registered Nurses’ Association of Ontario. *Delirium, dementia, and depression in older adults: assessment and care*. 2nd ed. Toronto (ON): RNAO; 2016.
35. Qaseem A, Barry MJ, Kansagara D. Nonpharmacologic versus pharmacologic treatment of adult patients with major depressive disorder: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. 2016;164(5):350-359. doi:[10.7326/M15-2570](https://doi.org/10.7326/M15-2570)
36. Instituto Mexicano del Seguro Social. *Guía de práctica clínica gpc: diagnóstico y tratamiento de la depresión en el adulto mayor en el primer nivel de atención*. Ciudad de México: IMSS; 2011.
37. Secretaría de Salud (MX). *Diagnóstico y tratamiento del trastorno depresivo en el adulto*. Ciudad de México: Secretaría de Salud; 2015.
38. Secretaría de Salud (MX). *Diagnóstico y tratamiento del trastorno depresivo en el adulto: intervenciones de enfermería para la detección, atención y control de la depresión en el adulto mayor en los tres niveles de atención*. Ciudad de México: Secretaría de Salud; 2016.



39. Chua HC, Chan LL, Chee KS, Chen YH, Chin SA, Chua PLW, et al. *Depression*. Singapore: Ministry of Health; 2012. Available from: <http://www.smj.org.sg/sites/default/files/5302/5302cpg1.pdf>. Accessed June 30, 2017.
40. Malhi GS, Bassett D, Boyce P, et al. Royal Australian and New Zealand college of psychiatrists clinical practice CPGs for mood disorders. *Aust NZ J Psychiatry*. 2015;49(12):1087-1206.doi:[10.1177/0004867415617657](https://doi.org/10.1177/0004867415617657)
41. Unipolar Depression Guideline Team. *Ambulatory unipolar depression guideline*. Ann Arbor (MI): UMHS; c2021. Available from: <http://michmed-clinical.policystat.com/policy/8093108/>. . Accessed June 30, 2017.
42. Driot D, Bismuth M, Maurel A, et al. Management of first depression or generalized anxiety disorder episode in adults in primary care: a systematic metareview. *Presse Med*. 2017;46(12 Pt 1):1124-1138. doi:[10.1016/j.lpm.2017.10.010](https://doi.org/10.1016/j.lpm.2017.10.010)
43. Bauer M, Pfennig A, Severus E, et al. World federation of societies of biological psychiatry (WFSBP) CPGs for biological treatment of unipolar depressive disorders, part 1: update 2013 on the acute and continuation treatment of unipolar depressive disorders. *World J Biol Psychiatry*. 2013;14(5):334-385. doi:[10.3109/15622975.2013.804195](https://doi.org/10.3109/15622975.2013.804195)
44. Kennedy SH, Lam RW, McIntyre RS, et al. Canadian network for mood and anxiety treatments (CANMAT), 2016, clinical CPGs for the management of adults with major depressive disorder: section 3: pharmacological treatments. *Can J Psychiatry*. 2016;61(9):540-560. doi:[10.1177/0706743716659417](https://doi.org/10.1177/0706743716659417)
45. Dua T, Barbui C, Clark N, et al. Evidence-based CPGs for mental, neurological, and substance use disorders in low-and middle-income countries: summary of WHO recommendations. *PLoS Med*. 2011;8(11):e1001122. doi:[10.1371/journal.pmed.1001122](https://doi.org/10.1371/journal.pmed.1001122)
46. McIntyre RS, Suppes T, Tandon R, Ostacher M. Florida best practice psychotherapeutic medication CPGs for adults with major depressive disorder. *J Clin Psychiatry*. 2017;78(6):703-713.doi:[10.4088/JCP.16cs10885](https://doi.org/10.4088/JCP.16cs10885)
47. Bauer M, Severus E, Köhler S, Whybrow PC, Angst J, Möller, H-J. World federation of societies of biological psychiatry (WFSBP) CPGs for biological treatment of unipolar depressive disorders: part 2: maintenance treatment of

- major depressive disorder-update 2015. *World J Biol Psychiatry*. 2015;16(2):76-95. doi:[10.3109/15622975.2014.1001786](https://doi.org/10.3109/15622975.2014.1001786)
48. Malaysian Health Technology Assessment Section. *Management of major depressive disorder*. 2nd ed. Putrajaya: Ministry of Health; 2019.
49. Gelenberg AJ, Freeman MP, Markowitz JC, et al. Practice guideline for the treatment of patients with major depressive disorder. 3th ed. Washington, DC: American Psychological Association; 2010. Available from: [https://psychiatryonline.org/pb/assets/raw/sitewide/practice\\_CPGs/CPGs/mdd.pdf](https://psychiatryonline.org/pb/assets/raw/sitewide/practice_CPGs/CPGs/mdd.pdf). Accessed August 14, 2017.
50. NHG-werkgroep. *Depressie: version 3.1* [Under review]. Available from: <https://richtlijnen.nhg.org/standaarden/depressie>. Accessed March 18, 2021.
51. Cleare A, Pariante CM, Young AH, et al. Evidence-based CPGs for treating depressive disorders with antidepressants: a revision of the 2008 British association for psychopharmacology CPGs. *J Psychopharmacol*. 2015;29(5):459-525. doi:[10.1177/0269881115581093](https://doi.org/10.1177/0269881115581093)
52. Won E, Park S-C, Han K-M, et al. Evidence-based, pharmacological treatment guideline for depression in Korea, revised edition. *J Korean Med Sci*. 2014;29:468-484. <https://doi.org/10.3346/jkms.2014.29.4.468>
53. Suomalainen Lääkäriseura Duodecim. *Depressio*. Available from: <https://www.kaypahoito.fi/hoi50023>. Accessed September 18, 2021.
54. Ruberto VL, Jha MK, Murrough JW. Pharmacological treatments for patients with treatment-resistant depression. *Pharmaceuticals*. 2020;13(6):116. <https://doi.org/10.3390/ph13060116>
55. British Columbia. *Major depressive disorder in adults: diagnosis and management*. British Columbia: BC; 2013. Available FROM: <https://www2.gov.bc.ca/gov/content/health/practitioner-professional-resources/bc-guidelines/depression-in-adults>. Accessed January 24, 2017.
56. Giakoumatos CI, Osse D. The psychopharmacology algorithm project at the Harvard South Shore Program: an update on unipolar nonpsychotic depression. *Harv Rev Psychiatry*. 2019;27(1):33-52. doi:[10.1097/HRP.0000000000000197](https://doi.org/10.1097/HRP.0000000000000197)
57. Bennabi D, Charpeaud T, Yrondi A, et al. Clinical CPGs for the management of treatment-resistant depression: French recommendations from experts, the French association for biological psychiatry and neuropsychopharmacology and

- the foundation FondaMental. *BMC Psychiatry*. 2019;19(1):262. doi:[10.1186/s12888-019-2237-x](https://doi.org/10.1186/s12888-019-2237-x)
58. Okasha A, Alkdhari S, Al Sharqi A, et al. Arab treatment guidelines for the management of major depressive disorder. *Arab J Psychiatry*. 2017;28(2): 97-117. doi:[10.12816/0041709](https://doi.org/10.12816/0041709)
59. Dodd S, Mitchell PB, Bauer M, et al. Monitoring for antidepressant-associated adverse events in the treatment of patients with major depressive disorder: an international consensus statement. *World J Biol Psychiatry*. 2018;19(5):330-348. doi:[10.1080/15622975.2017.1379609](https://doi.org/10.1080/15622975.2017.1379609)
60. Bauer M, Severus E, Möller H-J, et al. Pharmacological treatment of unipolar depressive disorders: summary of WFSBP guidelines. *Int J Psychiatry Clin Pract*. 2017;21(3):166-176. doi:[10.1080/13651501.2017.1306082](https://doi.org/10.1080/13651501.2017.1306082)
61. Bennabi D, Yrondi A, Charpeaud T, et al. Clinical guidelines for the management of depression with specific comorbid psychiatric conditions French recommendations from experts: the French Association for Biological Psychiatry and Neuropsychopharmacology and the foundation FondaMental. *BMC Psychiatry*. 2019;19(1):50. doi:[10.1186/s12888-019-2025-7](https://doi.org/10.1186/s12888-019-2025-7)
62. Sakurai H, Uchida H, Kato M, Suzuki T, Baba H, Watanabe K, et al. Pharmacological management of depression: Japanese expert consensus. *J Affect Disord*. 2020;266:626-632. doi:[10.1016/j.jad.2020.01.149](https://doi.org/10.1016/j.jad.2020.01.149)
63. Emsley R, Flisher AJ, Grobler G, Seedat S, Szabo CP. The South African society of psychiatrists (SASOP) treatment CPGs for psychiatric disorders. *S Afr J Psychiatr*. 2013;19(3):2. doi:[10.4102/sajpspsychiatry.v19i3.942](https://doi.org/10.4102/sajpspsychiatry.v19i3.942)
64. Connolly KR, Thase ME. If at first you don't succeed: a review of the evidence for antidepressant augmentation, combination and switching strategies. *Drugs*. 2011;71(1):43-64. doi:[10.2165/11587620-000000000-00000](https://doi.org/10.2165/11587620-000000000-00000)
65. Wang HR, Bahk W-M, Park Y-M, et al. Korean medication algorithm for depressive disorder: comparisons with other treatment CPGs. *Psychiatry Investig*. 2014;11(1):1-11. doi:[10.4306/pi.2014.11.1.1](https://doi.org/10.4306/pi.2014.11.1.1)
66. Park LT, Zarate CA Jr. Depression in the primary care setting. *N Engl J Med*. 2019;380(6):559-568. doi:[10.1056/NEJMc1712493](https://doi.org/10.1056/NEJMc1712493)
67. Voineskos D, Daskalakis ZJ, Blumberger DM. Management of treatment-resistant depression: challenges and strategies. *Neuropsychiatr Dis Treat*. 2020;16:221-234. doi:[10.2147/NDT.S198774](https://doi.org/10.2147/NDT.S198774)

68. Voytenko V, Nykamp L, Achtyes E, et al. Evidence-based practice guideline for the treatment of adult patients with depressive disorders: part I: psychiatric management. *Psychiatr Psychol Klin*. 2018;18(3):234-241. doi:[10.15557/PiPK.2018.0029](https://doi.org/10.15557/PiPK.2018.0029)
69. Piotrowski P, Gondek TM, Rymaszewska J, et al. Guidelines of the Polish Psychiatric Association: wroclaw division, the Polish society of family medicine and the college of family physicians in Poland for diagnosis and treatment of depressive disorders in primary health care. *Fam Med Prim Care Rev*. 2017;19(3):335-346. doi:[10.5114/fmpcr.2017.69300](https://doi.org/10.5114/fmpcr.2017.69300)
70. Bayes A, Parker G. How to choose an antidepressant medication. *Acta Psychiatr Scand*. 2019;139(3):280-291. doi:[10.1111/acps.13001](https://doi.org/10.1111/acps.13001)
71. Malhi GS, Hitching R, Berk M, Boyce P, Porter R, Fritz K. Pharmacological management of unipolar depression. *Acta Psychiatr Scand Suppl*. 2013;127(suppl 443):6-23. doi:[10.1111/acps.12122](https://doi.org/10.1111/acps.12122)
72. Doumy O, Bennabi D, El-Hage W, et al. Dépression résistante: les stratégies de potentialisation. *Presse Med*. 2016;45(3):338-349. doi:[10.1016/j.lpm.2016.02.004](https://doi.org/10.1016/j.lpm.2016.02.004)
73. Mulsant BH, Blumberger DM, Ismail Z, Rabheru K, Rapoport MJ. A systematic approach to pharmacotherapy for geriatric major depression. *Clin Geriatr Med*. 2014;30(3):517-534. doi:[10.1016/j.cger.2014.05.002](https://doi.org/10.1016/j.cger.2014.05.002)
74. Gałęcki P, Bliźniewska-Kowalska K. Treatment-resistant depression – recommendations of the National Consultant in the field of psychiatry. *Psychiatr Pol*. 2021;55(1):7–21. doi:[10.12740/PP/OnlineFirst/115208](https://doi.org/10.12740/PP/OnlineFirst/115208)
75. Avasthi A, Grover S. Clinical practice CPGs for management of depression in elderly. *Indian J Psychiatry*. 2018;60(suppl 3):S341-S362. doi:[10.4103/0019-5545.224474](https://doi.org/10.4103/0019-5545.224474)
76. Möller HJ, Bitter I, Bobes J, et al. Position statement of the European psychiatric association (EPA) on the value of antidepressants in the treatment of unipolar depression. *Eur Psychiatry*. 2012;27(2):114-128. doi:[10.1016/j.eurpsy.2011.08.002](https://doi.org/10.1016/j.eurpsy.2011.08.002)
77. Charpeaud T, Molierè F, Bubrowszky M, et al. Dépression résistante: les stratégies de changement et d'association de médicaments antidépresseurs. *Presse Med*. 2016;45(3):329-337. doi:[10.1016/j.lpm.2016.02.003](https://doi.org/10.1016/j.lpm.2016.02.003)

78. Busch FN, Sandberg LS. Combined treatment of depression. *Psychiatr Clin North Am.* 2012;35(1):165-179. doi:[10.1016/j.psc.2011.10.002](https://doi.org/10.1016/j.psc.2011.10.002)
79. Mathys M, Mitchell BG. Targeting treatment-resistant depression. *J Pharm Pract.* 2011;24(6):520-533. doi:[10.1177/0897190011426972](https://doi.org/10.1177/0897190011426972)
80. Taylor WD. Depression in the elderly. *N Engl J Med.* 2014;371(13):1228-1236. doi:[10.1056/NEJMc1402180](https://doi.org/10.1056/NEJMc1402180)
81. Gartlehner G, Thaler K, Hill S, Hansen RA. How should primary care doctors select which antidepressants to administer?. *Curr Psychiatry Rep.* 2012;14(4):360-369. doi:[10.1007/s11920-012-0283-x](https://doi.org/10.1007/s11920-012-0283-x)
82. Pereira Sanchez V, Molero Santos P. Protocolo diagnóstico y terapéutico de la depresión. *Medicine.* 2019;12(86):5070-5074. doi:[10.1016/j.med.2019.09.015](https://doi.org/10.1016/j.med.2019.09.015)
83. Gautam S, Jain A, Gautam M, Vahia VN, Grover S. Clinical practice CPGs for the management of depression. *Indian J Psychiatry.* 2017;59(suppl 1):S34-S50. doi:[10.4103/0019-5545.196973](https://doi.org/10.4103/0019-5545.196973)
84. Halaris A. A primary care focus on the diagnosis and treatment of major depressive disorder in adults. *J Psychiatr Pract.* 2011;17(5):340-350. doi:[10.1097/01.pra.0000405364.90043.bd](https://doi.org/10.1097/01.pra.0000405364.90043.bd)
85. Patejuk-Mazurek I. Selected antidepressants and anxiolytics: practical guidelines and case studies. *Psychiatria.* 2017;14(3):135-142. Polish. Available from: <https://journals.viamedica.pl>. Accessed January 24, 2017.