



Randomized Controlled trial evaluating the Efficacy of Wrap Therapy for Wound Healing Acceleration in NPUAP Stage II and III Pressure Ulcer Patients

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2011-000371
Article Type:	Research
Date Submitted by the Author:	01-Sep-2011
Complete List of Authors:	<p>Bito, Seiji; National Hospital Organization Tokyo Medical Center, Division of Clinical Epidemiology Mizuhara, Akihiro; Sanwa Medical Corporation, Higashi-Washinomiya Hospita Onishi, Sandai; Nursing Health Services Facilities for the Elderly Hatta Takeuchi, Kensuke; Sanwa Medical Corporation, Higashi-Washinomiya Hospita Suzuki, Masatsune; Tsukuba Medical Center Hospital, General Internal Medicine Akiyama, Kazuhiro; Tokatsu Clinic Hospital Kobayashi, Kazuyo; Wakaba Ostomy Center Matsunaga, Kayoko; Fujita Health University School of Medicine, Department of Dermatology</p>
Primary Subject Heading:	Dermatology
Keywords:	WOUND MANAGEMENT, Geriatric dermatology < DERMATOLOGY, HEALTH ECONOMICS, PRIMARY CARE

SCHOLARONE™
Manuscripts

Randomized Controlled trial evaluating the Efficacy of Wrap Therapy for Wound Healing Acceleration in NPUAP Stage II and III Pressure Ulcer Patients

Seiji Bito¹⁾, Akihiro Mizuhara²⁾, Sandai Onishi³⁾, Kensuke Takeuchi²⁾, Masatsune Suzuki⁴⁾, Kazuhiro Akiyama⁵⁾, Kazuyo Kobayashi⁶⁾, Kayoko Matsunaga⁷⁾

¹⁾ Division of Clinical Epidemiology, National Hospital Organization Tokyo Medical Center

²⁾ Sanwa Medical Corporation Higashi-Washinomiya Hospital

³⁾ Nursing Health Services Facilities for the Elderly Hatta

⁴⁾ Department of General Internal Medicine, Tsukuba Medical Center Hospital

⁵⁾ Tokatsu Clinic Hospital

⁶⁾ Wakaba Ostomy Center

⁷⁾ Department of Dermatology, Fujita Health University School of Medicine

Corresponding Author:

Seiji Bito

Division of Clinical Epidemiology, National Hospital Organization Tokyo Medical Center,

2-5-1 Higashigaoka, Meguro-ku, Tokyo

152-0021 Japan

e-mail: bitoseiji@kankakuki.go.jp

phone: 81-3-3411-0111 fax: 81-3-3411-3130

Abstract: 253 words

Text: words: 3,165 words

1
2
3
4 Abstract
5
6

7 **Objectives:** To evaluate if 'wrap therapy' using food wraps, which is widely used in Japanese
8 clinical sites, is not inferior when compared to guideline adherence treatments.
9

10 **Design:** Multi-centre, prospective, randomized, open blinded endpoint clinical trial.
11

12 **Setting:** Fifteen hospitals in Japan
13

14 **Patients:** Sixty-six elderly patients with new NPUAP Stage II or III pressure ulcers.
15

16 **Interventions:** Out of these 66 patients, 31 were divided into the conventional treatment
17 guidelines group and 35 into the wrap therapy group.
18

19 **Main outcome measures:** The primary end point was the period until the pressure ulcers were
20 cured. The secondary end point was a comparison of the speed of change in the Pressure Ulcer
21 Scale for Healing (PUSH) score.
22

23 **Results:** Sixty-four of the 66 patients were analysed. The estimated mean period until healing
24 was 57.5 days [95% confidence interval (CI) 45.2–69.8] in the control group as opposed to 59.8
25 days (95% CI 49.7–69.9) in the wrap therapy group. By the extent of pressure ulcer
26 infiltration, the mean period until healing was 16.0 days in the control group (95% CI
27 8.1–23.9) as opposed to 18.8 days in the wrap therapy group (95% CI 10.3–27.2) with NPUAP
28 Stage II ulcers, and 71.8 days (95% CI 61.4–82.3) as opposed to 63.2 days (95% CI 53.0–73.4),
29 respectively, with Stage III ulcers. There is no statistical significance in difference in PUSH
30 scores.
31

32 **Conclusions:** It might be possible to recommend wrap therapy as an effective treatment in
33 primary care settings as an extremely simple and inexpensive dressing care.
34

35 **Trial registration:** UMIN Clinical Trials Registry UMIN000002658
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Article Summary

Article focus

- ‘Wrap therapy’ is a method for localized care of pressure ulcers using polyethylene sheets, such as in food wraps, to the pressure ulcers. There are numerous case reports promoting efficiency of this therapy in Japan.
- We hypothesized non-inferiority of the efficacy of wrap therapy on adult patients with pressure ulcers classified as National Pressure Ulcer Advisory Panel (NPUAP) Stage II or III comparing with that of guideline adhesion treatment to Japanese patients.

Key messages

- The survival curves showed the inferiority of wrap therapy comparing with guideline adhesion treatment on the period until the pressure ulcers healed.
- Focusing on NPUAP Stage III pressure ulcers, wrap therapy showed tendency of faster healing than guideline adhesion treatment.

Strengths and limitations of this study

- This study is the first study revealing the efficacy of wrap therapy, that is inexpensive and simple for home care, using RCT design.
- Relatively weak statistical power and impossibility of blindness of the treatment.

INTRODUCTION

Among the health problems specific to the frail elderly, pressure ulcers¹⁻³ has been a major health problem, and the establishment and spread of an effective treatment method for it has been a pressing issue. After guideline publication by Agency for Health Care Policy and Research⁴, there have been few standard policy announcements regarding localized treatments with high evidence levels^{5,6}. The Japanese Society of Pressure Ulcers (JSPU) published its 'Evidence-Based Localized Pressure Ulcer Treatment Guidelines' in 2005, and treatments based on these have become the conventional treatments in Japan⁷. According to the guidelines, overall complex treatments aiming to debride necrotic tissue and reduce the ulcers are recommended for deep pressure ulcers. However, the recommendation grades regarding individual ointments and dressings, physical treatments, and surgical treatments are low level. Several intervention researches have been performed on specific drugs in localized treatments, but few clinical breakthrough methods have been discovered in the effects of these interventions⁸⁻¹⁴.

'Wrap therapy' is a method for localized care of pressure ulcers through which healing can be expected based on the natural healing effects by applying polyethylene sheets, such as in food wraps, to the pressure ulcers. This method has spread following its proposal by Toriyabe Shunichi¹⁵. On the effectiveness of wrap therapy in treating pressure ulcers, and of their simplicity and low cost, there are numerous case reports and case series research supporting no-inferiority of wrap therapy in Japan¹⁶⁻²⁰. On March 2010, JSPU approved wrap therapy as one of the first treatment choices in limited situations.

Our purpose is to evaluate the efficacy of wrap therapy on adult patients with pressure ulcers classified as National Pressure Ulcer Advisory Panel (NPUAP) Stage II or III on their backs using the current conventional treatment in Japan, described in the guideline published by JSPU, as a control. If this research verifies that wrap therapy is not harmful and has equivalent or better efficacy compared to conventional treatments, a low-cost treatment method can be well applied to the care of the elderly.

METHODS

Study Design and Setting

The study evaluated outcomes and analysis of the prospective randomized open blinded endpoint trial at numerous facilities. For this study, we recruited facilities by appealing to them for participation via mailing lists related to JSPU and those regarding pressure ulcer diagnosis and treatment. To assure implementation of the research work, we set the following as suitable standards for the facilities sharing the work: facilities that were able to use body pressure diffusion mats for pressure ulcer patients, facilities with experience of wrap therapy on some patients, the existence of care systems for pressure ulcer care and environments that were able to perform pressure ulcer treatments during hospitalization. Fifteen hospitals finally were identified to be eligible facilities.

Patients and randomization

For patient registration, we set the following inclusion and exclusion criteria: patients aged 50 or over with one or more NPUAP Stage II or III pressure ulcers on either their torso or trochanter, body temperature of 35.5°C min. to 37.5°C max., 600kcal or over daily intake, no critical nutritional impairment, renal failure, cirrhosis, immunosuppression, uncontrollable diabetes, or malignant tumours according to an examination performed within past four weeks. End-of-life patients whose estimated alive period were less than three months were excluded. When the patient did not possess the ability to make the autonomous decision to participate in the study, the outline and methods of the research were explained to their representing immediate family member, from whom written consent was obtained. After obtaining written consent, then uncoordinated random allocations were performed at the registration centre in the research office within 24 hours.

Interventions

Wrap therapy was defined as a method of treatment that uses food wraps and perforated polyethylene as a wound dressing. In many cases, wrap therapy follows the procedure of

1
2
3
4 thoroughly irrigating the pressure ulcers (Fig. 1-1) before directly covering the pressure
5
6 ulcers with food wrap if the exudation is small, or with commercially available perforated
7
8 polyethylene sheets and diapers, or with a cover sheet combined with sanitary towels if the
9
10 exudation is great (Figs. 1-2 and 1-3). Irrigation and the covering process are performed every
11
12 day²¹.

13
14 To standardize the categories and treatment procedures of wrap therapy as much as possible,
15
16 a meeting for instruction of interventions was held beforehand. A DVD of the lecture was
17
18 distributed to the physicians that were unable to participate in the instruction meeting.

19
20 Patients allocated to the control group were treated using methods conforming the 'Evidence-
21
22 Based Localized Pressure Ulcer Treatment Guidelines' issued by JSPU⁷. These guidelines are
23
24 diagnosis and treatment guidelines created by JSPU based on current diagnosis and
25
26 treatment evidence.

27 28 Outcomes evaluated

29
30
31 The primary endpoint was defined as the period from the start of registration until the
32
33 pressure ulcers that had been the subject of observation for three months were healed. After
34
35 the patient had been discharged from the hospital, continuous observation was performed as
36
37 far as possible through diagnosis and treatment at home. Every wound heal was confirmed
38
39 by supervising physicians. As the secondary endpoint, we measured the PUSH score for the
40
41 localized status of the pressure ulcers as defined by NPUAP at two, four, six, eight, ten and 12
42
43 weeks from the start of registration²².

44
45 To avoid the occurrence of observation bias, we established an 'outcome evaluation centre'.
46
47 Digital camera images were sent from every facility to the outcome evaluation centre, and all
48
49 outcomes were evaluated there. The outcome evaluation centre was blinded so as not to know
50
51 the treatment group to which the subject belonged. Because it was impossible to evaluate the
52
53 exudates amount using images, records were made by the observer directly.

54
55 We also recorded the total amount of ointment used in localized treatments during the
56
57 observation period, and performed rough calculations with regards to cost.

58
59 Adverse Events and Role of the Patient Safety Monitoring Board
60

1
2
3
4 A safety evaluation committee was established comprising several specialists and
5 non-specialists. The members were independent from the patient registration, research
6 offices and the outcome evaluation centre. In all cases of death, worsening of the pressure
7 ulcers during the research period, systemic deterioration or sepsis occurring within 30 days
8 from the day of the protocol treatment, and when the connection to the pressure ulcers was
9 undeniable, researchers sharing the research at the core facilities reported to the committee.
10
11
12
13
14

15 16 17 Statistical Analysis

18
19
20 We aimed to clarify our statistical hypothesis that wrap therapy was not inferior compared to
21 conventional treatments conforming to the guidelines. The mean period until healing for the
22 localized treatment of Stage II pressure ulcers was set as 21 days based on previous
23 literature⁹⁻¹⁴. The non-inferior threshold was set at seven days according to clinical judgment.
24 When set to a tolerable threshold difference of 7 days when one side has a 5% significance
25 level and a test power of 90%, the required number of cases in the two cohorts was 80.
26 Similarly, the required number of cases for Stage III pressure ulcers was 60. For the total
27 number of target cases, 140 was set as the number of target registered patients. The analysts
28 were blinded about which group was the wrap therapy or the conventional treatment until
29 the analysis was finished. Intention to treat analysis was performed.
30
31
32
33
34
35
36
37
38

39 For the main endpoint comparisons, a Kaplan-Meier plots were created and the estimated
40 mean value until the endpoint occurrence and its 95% confidence interval (CI) were calculated.
41 The differences in the PUSH scores were calculated from two weeks immediately after the
42 start of observations, between two to four weeks, four to six weeks, six to eight weeks, eight to
43 ten weeks and ten to 12 weeks, and described the speed of pressure ulcer healing over time for
44 both groups. We used PASW Statistics 18 (SPSS, Inc) for the statistical analysis.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

The study flow diagram is shown in Fig. 2. During the research period, a total of 66 patients participated. Of these, two could not be followed up due to early discharge or transfer to another hospital. A total of 64 participants were analysed: 29 in the conventional treatment group and 35 in the wrap therapy group. The percentage of patients whose endpoints could be pursued four weeks after registration was 95%, eight weeks after registration was 77% and 12 weeks after registration was 64%. The characteristics distribution of the patients analysed was 33 females and 29 males, and the locations of the target pressure ulcers were most common on the sacrum (56%), followed by the trochanter (13%), gluteus (6%) and the coccyx (5%).

Table 1 shows a comparison of the patients' characteristics distribution, their health at registration, the status of their pressure ulcers and the treatment method for the two groups. The mean values in the conventional treatment and wrap therapy cohorts were 12.7 versus 12.7 for the Braden Scale at registration²³. The percentages using hydrocolloids, hydrogels or polyurethane foam as pressure ulcer dressings were 76% in the control group and 3% in the intervention cohort.

Time until the Pressure Ulcers Healed

The percentage of patients in whom healing of the pressure ulcers was verified from among the samples that could be followed up at four weeks after registration were 26% of the conventional group as opposed to 21% in the wrap therapy group; at eight weeks after registration the percentages were 46% in the conventional group as opposed to 52% in the wrap therapy group (Fig. 2).

Fig. 3-1 shows the survival curves of both groups. The mean of the estimated value until healing was 57.5 days (95% CI 45.2 days; 69.8 days) for the conventional group as opposed to 59.8 days (95% CI 49.7 days; 69.9 days) for the wrap therapy group. Fig. 3-2 and Fig. 3-3 describes the survival curve for the patients with NPUAP Stage II and III pressure ulcers, respectively. For Stage II pressure ulcers, the estimated mean value until healing were 16.0 days (95% CI 8.1 days; 23.9 days) for the conventional group compared to 18.8 days (95% CI

1
2
3
4 10.3 days; 27.2 days) for the wrap therapy group. Meanwhile, for Stage III pressure ulcers,
5 the estimated mean value until healing were 71.8 days (95% CI 61.4 days; 82.3 days) for the
6 conventional group compared to 63.2 days (95% CI 53.0 days; 73.4 days) for the wrap therapy
7 group.
8
9

10 11 12 13 Speed of Pressure Ulcer Healing Using PUSH Scores 14 15

16
17 The mean values of the difference in the PUSH scores at registration and upon either healing
18 or the final observation divided by the treatment period were 1.1 ± 2.1 points in the
19 conventional group and 0.9 ± 1.3 points in the wrap therapy group ($p = 0.73$ Student's t-test).
20 No significant difference was identified in the mean PUSH score reduction values in either
21 cohort from immediately after the start of observations to the second week, from the second to
22 the fourth week, from the fourth to the sixth week, from the sixth to the eighth week, from the
23 eighth to the tenth week and from the tenth to the twelfth week (Table 2).
24
25
26
27
28
29
30

31 Adverse Events 32 33

34
35 During the total observation period, there was systemic worsening such as pneumonia,
36 occurring immediately after the start of observation in three cases in the conventional group,
37 and four cases in the wrap therapy group. Two cases in the conventional group died of
38 pneumonia and one died from heart failure for a total of three deaths. In the wrap therapy
39 group, one patient died of pneumonia and one of senescence for a total of two deaths.
40 Regarding localized adverse events, there were problems with the covered skin (eczema,
41 maceration, rash, etc.) in the conventional group and six cases in the wrap therapy group.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSIONS

Statement of principal findings

'Wrap therapy' has already been introduced nationwide in Japanese clinical facilities. The main reasons for its spread in clinical settings are both that wrap therapy is a rational treatment method compared to basic wound-healing therapies, in that it is possible to maintain a lubricated environment without applying stress to the wound^{24,25}, and because compared to complex pressure ulcer treatments that combine various ointments and dressings, it has frequently been recognized empirically in actual clinical settings as having equal or better effects in treating pressure ulcers.

Looking only at the main outcome survival curves, and taking all the patients into consideration, almost identical results were obtained in both the wrap therapy and conventional treatment groups for the period until the pressure ulcers healed. Meanwhile, when the two treatment groups are compared using the NPUAP classification, although no clear difference between the survival curves of the two treatments were identified with shallow Stage II pressure ulcers, in the comparison between the Stage III pressure ulcer cohorts, the survival curve of the wrap therapy cohort showed that healing tended to be faster than in the conventional treatment cohort. No statistical significance, however, was presented due to insufficient statistical power. It might be possible to present results in which wrap therapy is sufficiently effective as an alternative treatment to the current conventional treatments for comparatively deep pressure ulcers such as Stage III pressure ulcers with bigger sample study. Nor was any significant difference identified between the wrap therapy and conventional treatment cohorts regarding the extent of PUSH score reduction either, and this result can also be stated to support the fact that wrap therapy is not inferior to conventional treatments. Nevertheless, initially we estimated 80 cases and 60 cases in the NPUAP classification subgroups as the non-inferior estimate, but on this occasion, the total effective sample, at 64 cases, lacked sufficient statistical power.

During the comparison of adverse events, there were concerns that skin problems caused by food wraps and perforated polyethylene used in the wrap therapy might occur, but in our research, no significant difference was identified with the results of the conventional treatments. Rashes due to adhesive plasters and tape were identified in four cases in the

1
2
3 conventional cohort, but none were observed in the wrap therapy cohort. This is thought to be
4 because it is basically possible to implement wrap therapy merely by making the patient wear
5 paper diapers without affixing tape after applying the dressing.
6
7
8
9

10 11 Strengths and weakness of the study 12 13

14
15 This research is the world's first randomized controlled trials that comparing conventional
16 treatments and wrap therapy. The conclusions obtained will greatly influence pressure ulcer
17 care methods in the future. Theoretical superiority of wrap therapy to conventional
18 treatments conforming to guidelines is the simplicity and cost effectiveness of the
19 treatment^{20,26}. Conventional treatments require the use of multiple ointments and dressings
20 of suitable size and type, and the treatment requires a certain level of specialist skills. The
21 simplicity of treatment protocols using wrap therapy might better promote the spread of care
22 skills. Regarding cost effectiveness, the various ointments and dressings such as hydrocolloid
23 used in pressure ulcer treatments are often expensive, but the perforated polyethylene and
24 diaper sheets used in wrap therapy can be supplied at less than ten cents per treatment. If
25 the accelerated pressure ulcer healing effects of wrap therapy are the same as those of
26 conventional treatments, the efficiency of wrap therapy is presumably extremely great when
27 considering cost effectiveness. Larger scale additional tests strengthening the verified
28 backing of these research results could offer new, simple and effective methods of pressure
29 ulcer care and noticeably improve the benefits to both patients and society.
30
31
32
33
34
35
36
37
38
39
40
41
42

43 This research, on the other hand, has some limitations. The greatest concern was
44 impossibility of blindness of the treatment. Without blindness, health staff may have biases
45 on the treatments themselves and wound evaluations. To minimize observation bias, we
46 attempted to eliminate bias occurring during the evaluations by combining all the outcomes
47 in a single outcome evaluation centre as far as possible. The second limitation resulted in bias
48 among the participating facilities. Wrap therapy is currently viewed as an alternative
49 treatment of choice. The facilities that participated in this research currently use wrap
50 therapy clinically under this present condition, and the possibility of wrap therapy effects
51 being overvalued beforehand cannot be denied.
52
53
54
55
56
57
58
59
60

Meaning of the study

The results we found could be called the high quality evidence to date from among the evidence that supports the safety and efficacy of wrap therapy, which have been reported many times in Japanese academic conferences concerned with wound healing and treatment of the elderly. In future, as the rapid aging of the population progresses worldwide, the problem of caring for the frail elderly in Japan, which has the highest rate of population aging in the world, will become a highly compelling problem in the societies of Europe and North America. Wrap therapy may be an epoch-making method of treatment and the verification of its usefulness by clinical experiment is thought to be a major advance for the future elderly care.

Acknowledgments

The authors would like to thank the following volunteering facilities for registering patients for this study: Tokatsu-Clinic Hospital, Wakaba Ostomy Centre, Sozokai Heiwadai Hospital, Tokushima Kensei Hospital, Ryumondo Ono Hospital, Kojinkai Hoshigaura Hospital, Ibi Kosei Hospital, Niimi Central Hospital, Oyama Municipal Hospital, Hakuai Kaita Hospital, Yatsu Hoken Hospital, Kikai Tokushukai Hospital, Naruto Yamakami Hospital, Higashi Washinomiya Hospital, and Kaneda Hospital. Shigeru Ichioka and Yuichiro Kume of the Orthopedic Surgery Department, Saitama Medical University for being in charge of the outcome evaluation centre and evaluating the images of pressure ulcers. Toshiko Yamana of the Human Care Studies Department at Tohto College of Health Sciences, Shingo Okada at Kitamihara Clinic, and Yoshiaki Ito at Kiyosu Clinic for supervising the Safety Evaluation Committee, and Megumi Matsubara and Miki Hirano at Higashi Washinomiya Hospital.

Trial registration: UMIN Clinical Trials Registry UMIN000002658. Summary protocol is available on

<https://upload.umin.ac.jp/cgi-bin/ctr/ctr.cgi?function=brows&action=brows&type=detail&recptno=R000003235&admin=0&language=J>

Contributors: SB conducted literature review, developed the initial protocol, examined statistical analysis and drafted the manuscript. AM wrote grant application, recruited collaboration facilities, developed intervention protocol and planned research meeting and startup meeting. SO contributed to research design, reviewed and amended the research protocol including intervention protocol and contributed to enrolment of patients. KT also contributed to research design, reviewed and amended the research protocol and contributed to enrolment of patients. MS contributed to research design, reviewed and amended the research protocol. KA contributed to enrolment of patients. KK contributed to research design and enrolment of patients. KM managed an outcome evaluation centre.

Funding: This study was supported by Division of the Health for the Elderly at Japanese Ministry of Health, Labor, and Welfare. Grant name 'Examination and Research Work into New Pressure Ulcer Treatments for the Care of the Elderly'. Funding agency did not interfere with any process of study content. All researchers are independent from the founder. No funding was received from any specific commercial company.

Competing interests: All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organisation for the submitted work, no financial relationships with any organisations that might have an interest in the submitted work in the previous three years, and no other relationships or activities that could appear to have influenced the submitted work.

Ethical approval: The Japanese Academy of Family Medicine Ethics Review Board examined and approved our research plan and informed consent process in October, 2009.

Data sharing: no additional data available

Table 1: Patient Characteristics and Health Status, and Pressure Ulcer Status at Registration
(n=64)

	Conventional Treatment Cohort (n=29)	Wrap Therapy Cohort (n=35)	P value
Mean Age (years)	82±10	81±12	0.60
Sex (Female Percentage)	48	54	0.28
<u>Nutrition Absorption Status</u> (Each %)			0.27
Oral Nutrition	0	9	
Enteral Alimentation: Nasally	38	40	
Enteral Alimentation: Gastrostoma	10	3	
Central Venous Nutrition			
Use of Pressure-Resistant Diffusion Mattress (Percentage 'Yes')	93	100	0.20
Use of Diapers (Percentage 'Yes')	90	91	0.97
<u>Depth of Target Pressure Ulcer</u>			0.09
NPUAP Class II (%)	28	11	
NPUAP Class III (%)	72	89	
Pressure Ulcer Pockets (% Present)	38	34	0.28
<u>Calories Absorbed (Each %)</u>			0.58
Over 1200kcal	45	49	
800kcal to 1200kcal	45	37	
600kcal to 800kcal	10	9	
Serum Albumin (Mean ± SD)	2.8±0.5	2.9±0.5	0.61
Serum Creatinine (Mean ± SD)	0.66±0.3	0.64±0.3	0.77

1 2 3 4 5 6 7	Braden Scale at Registration (Mean ± SD)	12.8±3.5	12.7±2.8	0.89
8 9 10 11	PUSH Score at Registration (Mean ± SD)	10.8±2.6	10.7±2.7	0.91
12 13 14 15	Pressure Ulcer Surface Area at Registration (Mean ± SD)	14±21	15±25	0.79
16 17 18 19 20 21 22	Use of ointments or sprays including pharmaceuticals with tissue regeneration accelerant actions	21	14	0.006
23 24 25 26 27 28 29	Percentage using hydrocolloids, hydrogels, or polyurethane foam as a pressure ulcer dressing	76	3	<0.0001

Table 2: Comparison of the mean reduction in PUSH scores for both cohorts for two weeks immediately after the start of observations, between two to four weeks, four to six weeks, six to eight weeks, eight to ten weeks and ten to 12 weeks, respectively. See separate appendix.

	0 to 2W	2W to 4W	4W to 6W	6W to 8W	8W to 10W	10W to 12W
n	54	45	34	25	18	11
Control: Mean [95% CI]	1.8 [1.0-2.6]	0.4 [0.0-0.9]	0.7 [0.1-1.3]	0.8 [0.1-1.5]	0.3 [-0.1-0.7]	1.0 [0.1-1.9]
Wrapp therapy: Mean [95% CI]	1.8 [1.0-2.6]	0.4 [-0.2-1.0]	0.1 [-0.4-0.6]	0.3 [-0.2-0.8]	0.4 [-0.2-1.0]	0.5 [-0.1-1.1]
P value*	0.77	0.44	0.24	0.43	0.79	0.54

* tested by Mann-Whitney U test

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 1-1




Fig. 1-2



Fig. 1-3




review only



view only

Fig. 3

Fig. 3-1: Comparison of Survival Curves Using the Period until Pressure Ulcers Healing as the Endpoint - All Cases



Only

Fig. 3-2: Comparison of Survival Curves Using the Period until Pressure Ulcers Healing as the Endpoint - NPUAP Class II Pressure Ulcers Cohort





Fig. 3-3: Comparison of Survival Curves Using the Period until Pressure Ulcers Healing as the Endpoint - NPUAP Class III Pressure Ulcers Cohort



References

1. Brem H, Lyder C. Protocol for the successful treatment of pressure ulcers. *Am J Surg*. 2004; **188**(1A Suppl): 9-17.
2. Orlando PL. Pressure ulcer management in the geriatric patient. *Ann Pharmacother*. 1998; **32**(11): 1221-7.
3. Smith DM. Pressure ulcers in the nursing home. *Ann Intern Med*. 1995; **123**(6): 433-42.
4. Agency for Health Care Policy and Research. *Pressure ulcer treatment: Clinical Practice Guideline Quick Reference Guide*. 1994; (15): 1-25.
5. Cullum N, Nelson EA, Flemming K, et al. Systematic reviews of wound care management: (5) beds; (6) compression; (7) laser therapy, therapeutic ultrasound, electrotherapy and electromagnetic therapy. *Health Technol Assess*. 2001; **5**(9): 1-221.
6. McInnes E, Bell-Syer SE, Dumville JC, et al. Support surfaces for pressure ulcer prevention. *Cochrane Database Syst Rev*. 2008; (4): CD001735.
7. Japanese Society of Pressure Ulcers. *Evidence-Based Localized Pressure Ulcer Treatment Guidelines*. Tokyo: Shorinsya Co.; 2005.
8. Adunsky A, Ohry A. Decubitus direct current treatment (DDCT) of pressure ulcers: results of a randomized double-blinded placebo controlled study. *Arch Gerontol Geriatr*. 2005; **41**(3): 261-9.
9. Hollisaz MT, Khedmat H, Yari F. A randomized clinical trial comparing hydrocolloid, phenytoin and simple dressings for the treatment of pressure ulcers [ISRCTN33429693]. *BMC Dermatol*. 2004; **4**(1): 18.
10. Landi F, Aloe L, Russo A, et al. Topical treatment of pressure ulcers with nerve growth factor: a randomized clinical trial. *Ann Intern Med*. 2003; **139**(8): 635-41.
11. Graumlich JF, Blough LS, McLaughlin RG, et al. Healing pressure ulcers with collagen or hydrocolloid: a randomized, controlled trial. *J Am Geriatr Soc*. 2003; **51**(2): 147-54.
12. Belmin J, Meaume S, Rabus MT, et al. Sequential treatment with calcium alginate dressings and hydrocolloid dressings accelerates pressure ulcer healing in older subjects: a multicenter randomized trial of sequential versus nonsequential treatment with hydrocolloid dressings alone. *J Am Geriatr Soc*. 2002; **50**(2): 269-74.

13. Seeley J, Jensen JL, Hutcherson J. A randomized clinical study comparing a hydrocellular dressing to a hydrocolloid dressing in the management of pressure ulcers. *Ostomy Wound Manage.* 1999; **45**(6): 39-44, 6-7.
14. Heyneman A, Beele H, Vanderwee K, et al. A systematic review of the use of hydrocolloids in the treatment of pressure ulcers. *J Clin Nurs.* 2008; **17**(9): 1164-73.
15. Toriyabe S, Saito H, Sakurai K. Use of a food wrap as a dressing material. *Advances in Wound Care.* 1999; **12**(8): 405-6.
16. Mizuhara A. Treatment results and problem of Wrap therapy. *Japanese Journal of Pressure Ulcers.* 2009; **11**(3): 286-7.
17. Okamura S, Hisatomi M, Iwasaki K, et al. *Japanese Journal of Pressure Ulcers.* 2002; **4**(3): 427-30.
18. Toriyabe S, Suemaru S. A study of food-wrap therapy for Stage III and Stage IV pressure ulcers. *Nippon Ishikai Zasshi.* 2000; **123**(10): 1605-11.
19. Bailey KA. Wrap for leg ulcers. *Am J Nurs.* 1996; **96**(6): 17.
20. Ohnishi S, Koike T, Shiogama K, et al. Use of food wrap films for treating pressure ulcers. *Japanese J of Med and Pharmaceutical Science.* 2006; **55**(4): 561-7.
21. Takahashi J, Yokota O, Fujisawa Y, et al. An evaluation of polyvinylidene film dressing for treatment of pressure ulcers in older people. *J Wound Care.* 2006; **15**(10): 449-50, 52-4.
22. Woodbury MG, Houghton PE, Campbell KE, et al. Pressure ulcer assessment instruments: a critical appraisal. *Ostomy Wound Manage.* 1999; **45**(5): 42-5, 8-50, 3-5.
23. Lewicki LJ, Mion L, Splane KG, et al. Patient risk factors for pressure ulcers during cardiac surgery. *AORN J.* 1997; **65**(5): 933-42.
24. Ozturk F, Ermertcan AT. Wound healing: a new approach to the topical wound care. *Cutan Ocul Toxicol.* 2011; **30**(2):92-9.
25. Bishop SM, Walker M, Rogers AA, et al. Importance of moisture balance at the wound-dressing interface. *J Wound Care.* 2003; **12**(4): 125-8.
26. Schuurman JP, Schoonhoven L, Defloor T, et al. Economic evaluation of pressure ulcer care: a cost minimization analysis of preventive strategies. *Nurs Econ.* 2009; **27**(6): 390-400, 15.



CONSORT 2010 checklist of information to include when reporting a randomised trial*

Section/Topic	Item No	Checklist item	Reported on page No
Title and abstract			
	1a	Identification as a randomised trial in the title	1
	1b	Structured summary of trial design, methods, results, and conclusions (for specific guidance see CONSORT for abstracts)	2
Introduction			
Background and objectives	2a	Scientific background and explanation of rationale	3
	2b	Specific objectives or hypotheses	3
Methods			
Trial design	3a	Description of trial design (such as parallel, factorial) including allocation ratio	4
	3b	Important changes to methods after trial commencement (such as eligibility criteria), with reasons	18
Participants	4a	Eligibility criteria for participants	4
	4b	Settings and locations where the data were collected	4
Interventions	5	The interventions for each group with sufficient details to allow replication, including how and when they were actually administered	4-5
Outcomes	6a	Completely defined pre-specified primary and secondary outcome measures, including how and when they were assessed	5-6
	6b	Any changes to trial outcomes after the trial commenced, with reasons	NA
Sample size	7a	How sample size was determined	6
	7b	When applicable, explanation of any interim analyses and stopping guidelines	6
Randomisation:			
Sequence generation	8a	Method used to generate the random allocation sequence	4
	8b	Type of randomisation; details of any restriction (such as blocking and block size)	4
Allocation concealment mechanism	9	Mechanism used to implement the random allocation sequence (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned	NA
Implementation	10	Who generated the random allocation sequence, who enrolled participants, and who assigned participants to	4

1		interventions	
2	Blinding	11a	If done, who was blinded after assignment to interventions (for example, participants, care providers, those assessing outcomes) and how
3			5
4		11b	If relevant, description of the similarity of interventions
5			NA
6	Statistical methods	12a	Statistical methods used to compare groups for primary and secondary outcomes
7		12b	Methods for additional analyses, such as subgroup analyses and adjusted analyses
8			6
9	Results		
10	Participant flow (a diagram is strongly recommended)	13a	For each group, the numbers of participants who were randomly assigned, received intended treatment, and were analysed for the primary outcome
11			7, 13-14
12		13b	For each group, losses and exclusions after randomisation, together with reasons
13			7, 18
14	Recruitment	14a	Dates defining the periods of recruitment and follow-up
15		14b	Why the trial ended or was stopped
16			NA
17	Baseline data	15	A table showing baseline demographic and clinical characteristics for each group
18			13-14
19	Numbers analysed	16	For each group, number of participants (denominator) included in each analysis and whether the analysis was by original assigned groups
20			18
21	Outcomes and estimation	17a	For each primary and secondary outcome, results for each group, and the estimated effect size and its precision (such as 95% confidence interval)
22			7-8
23		17b	For binary outcomes, presentation of both absolute and relative effect sizes is recommended
24			8
25	Ancillary analyses	18	Results of any other analyses performed, including subgroup analyses and adjusted analyses, distinguishing pre-specified from exploratory
26			7-8
27	Harms	19	All important harms or unintended effects in each group (for specific guidance see CONSORT for harms)
28			8
29	Discussion		
30	Limitations	20	Trial limitations, addressing sources of potential bias, imprecision, and, if relevant, multiplicity of analyses
31			10
32	Generalisability	21	Generalisability (external validity, applicability) of the trial findings
33			11
34	Interpretation	22	Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence
35			9-10
36	Other information		
37	Registration	23	Registration number and name of trial registry
38			11
39	Protocol	24	Where the full trial protocol can be accessed, if available
40			11
41	Funding	25	Sources of funding and other support (such as supply of drugs), role of funders
42			12

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

*We strongly recommend reading this statement in conjunction with the CONSORT 2010 Explanation and Elaboration for important clarifications on all the items. If relevant, we also recommend reading CONSORT extensions for cluster randomised trials, non-inferiority and equivalence trials, non-pharmacological treatments, herbal interventions, and pragmatic trials. Additional extensions are forthcoming: for those and for up to date references relevant to this checklist, see www.consort-statement.org.

For peer review only



Randomized Controlled trial evaluating the Efficacy of Wrap Therapy for Wound Healing Acceleration in NPUAP Stage II and III Pressure Ulcer Patients

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2011-000371.R1
Article Type:	Research
Date Submitted by the Author:	20-Nov-2011
Complete List of Authors:	Bito, Seiji; National Hospital Organization Tokyo Medical Center, Division of Clinical Epidemiology Mizuhara, Akihiro; Sanwa Medical Corporation, Higashi-Washinomiya Hospita, Onishi, Sandai; Nursing Health Services Facilities for the Elderly Hatta, Takeuchi, Kensuke; Sanwa Medical Corporation, Higashi-Washinomiya Hospita, Suzuki, Masatsune; Tsukuba Medical Center Hospital, General Internal Medicine Akiyama, Kazuhiro; Tokatsu Clinic Hospital, Kobayashi, Kazuyo; Wakaba Ostomy Center, Matsunaga, Kayoko; Fujita Health University School of Medicine, Department of Dermatology
Primary Subject Heading:	Dermatology
Secondary Subject Heading:	Geriatric medicine, Pharmacology and therapeutics
Keywords:	WOUND MANAGEMENT, Geriatric dermatology < DERMATOLOGY, HEALTH ECONOMICS, PRIMARY CARE

SCHOLARONE™
Manuscripts

Randomized Controlled trial evaluating the Efficacy of Wrap Therapy for Wound Healing Acceleration in NPUAP Stage II and III Pressure Ulcer Patients

Seiji Bito¹⁾, Akihiro Mizuhara²⁾, Sandai Onishi³⁾, Kensuke Takeuchi²⁾, Masatsune Suzuki⁴⁾,
Kazuhiro Akiyama⁵⁾, Kazuyo Kobayashi⁶⁾, Kayoko Matsunaga⁷⁾

¹⁾ Division of Clinical Epidemiology, National Hospital Organization Tokyo Medical Center

²⁾ Sanwa Medical Corporation Higashi-Washinomiya Hospital

³⁾ Nursing Health Services Facilities for the Elderly Hatta

⁴⁾ Department of General Internal Medicine, Tsukuba Medical Center Hospital

⁵⁾ Tokatsu Clinic Hospital

⁶⁾ Wakaba Ostomy Center

⁷⁾ Department of Dermatology, Fujita Health University School of Medicine

Corresponding Author:

Seiji Bito

Division of Clinical Epidemiology, National Hospital Organization Tokyo Medical Center,

2-5-1 Higashigaoka, Meguro-ku, Tokyo

152-0021 Japan

e-mail: bitoseiji@kankakuki.go.jp

phone: 81-3-3411-0111 fax: 81-3-3411-3130

Abstract: 252 words

Text: words: 3,451 words

1
2
3 Abstract
4
5
6

7 **Objectives:** To evaluate if ‘wrap therapy’ using food wraps, which is widely used in Japanese
8 clinical sites, is not inferior when compared to guideline adherence treatments.
9

10 **Design:** Multi-centre, prospective, randomized, open blinded endpoint clinical trial.
11

12 **Setting:** Fifteen hospitals in Japan
13

14 **Patients:** Sixty-six elderly patients with new NPUAP Stage II or III pressure ulcers.
15

16 **Interventions:** Out of these 66 patients, 31 were divided into the conventional treatment
17 guidelines group and 35 into the wrap therapy group.
18

19 **Main outcome measures:** The primary end point was the period until the pressure ulcers were
20 cured. The secondary end point was a comparison of the speed of change in the Pressure Ulcer
21 Scale for Healing (PUSH) score.
22
23

24 **Results:** Sixty-four of the 66 patients were analysed. The estimated mean period until healing
25 was 57.5 days [95% confidence interval (CI) 45.2–69.8] in the control group as opposed to 59.8
26 days (95% CI 49.7–69.9) in the wrap therapy group. By the extent of pressure ulcer
27 infiltration, the mean period until healing was 16.0 days in the control group (95% CI
28 8.1–23.9) as opposed to 18.8 days in the wrap therapy group (95% CI 10.3–27.2) with NPUAP
29 Stage II ulcers, and 71.8 days (95% CI 61.4–82.3) as opposed to 63.2 days (95% CI 53.0–73.4),
30 respectively, with Stage III ulcers. There is no statistical significance in difference in PUSH
31 scores.
32
33
34
35
36
37

38 **Conclusions:** It might be possible to consider wrap therapy as an alternative choice in primary
39 care settings as a simple and inexpensive dressing care.
40
41

42 **Trial registration:** UMIN Clinical Trials Registry UMIN000002658
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Article Summary

Article focus

- ‘Wrap therapy’ is a method for localized care of pressure ulcers using polyethylene sheets, such as in food wraps, to the pressure ulcers. There are numerous case reports promoting efficiency of this therapy in Japan.
- We hypothesized non-inferiority of the efficacy of wrap therapy on adult patients with pressure ulcers classified as National Pressure Ulcer Advisory Panel (NPUAP) Stage II or III comparing with that of guideline adhesion treatment to Japanese patients.

Key messages

- The survival curves **did not** showed the inferiority of wrap therapy comparing with guideline adhesion treatment on the period until the pressure ulcers healed.
- **The result of this study imply clinical utility of wrap therapy which have been reported many times in Japanese academic conferences concerned with wound healing and treatment of the elderly. Bigger sample trials are necessary to confirm this implication as rigorous clinical evidence.**

Strengths and limitations of this study

- This study is the first study revealing the efficacy of wrap therapy, that is inexpensive and simple for home care, using RCT design.
- Relatively weak statistical power and impossibility of blindness of the treatment.

INTRODUCTION

Among the health problems specific to the frail elderly, pressure ulcers¹⁻³ has been a major health problem, and the establishment and spread of an effective treatment method for it has been a pressing issue. After guideline publication by Agency for Health Care Policy and Research⁴, there have been few standard policy announcements regarding localized treatments with high evidence levels^{5,6}. The Japanese Society of Pressure Ulcers (JSPU) published its 'Evidence-Based Localized Pressure Ulcer Treatment Guidelines' in 2005, and treatments based on these have become the conventional treatments in Japan⁷. According to the guidelines, overall complex treatments aiming to debride necrotic tissue and reduce the ulcers are recommended for deep pressure ulcers. However, the recommendation grades regarding individual ointments and dressings, physical treatments, and surgical treatments are low level. Several intervention researches have been performed on specific drugs in localized treatments, but few clinical breakthrough methods have been discovered in the effects of these interventions⁸⁻¹⁴.

'Wrap therapy' is a method for localized care of pressure ulcers through which healing can be expected based on the natural healing effects by applying polyethylene sheets, such as in food wraps, to the pressure ulcers. This method has spread following its proposal by Toriyabe Shunichi¹⁵. On the effectiveness of wrap therapy in treating pressure ulcers, and of their simplicity and low cost, there are numerous case reports and case series research supporting no-inferiority of wrap therapy in Japan¹⁶⁻²⁰. On March 2010, JSPU approved wrap therapy as one of the first treatment choices in limited situations.

Our purpose is to evaluate the efficacy of wrap therapy on adult patients with pressure ulcers classified as National Pressure Ulcer Advisory Panel (NPUAP) Stage II or III on their backs using the current conventional treatment in Japan, described in the guideline published by JSPU, as a control. If this research verifies that wrap therapy is not harmful and has equivalent or better efficacy compared to conventional treatments, a low-cost treatment method can be well applied to the care of the elderly.

METHODS

Study Design and Setting

The study evaluated outcomes and analysis of the prospective randomized open blinded endpoint trial at numerous facilities. For this study, we recruited facilities by appealing to them for participation via mailing lists related to JSPU and those regarding pressure ulcer diagnosis and treatment. To assure implementation of the research work, we set the following as suitable standards for the facilities sharing the work: facilities that were able to use body pressure diffusion mats for pressure ulcer patients, facilities with experience of wrap therapy on some patients, the existence of care systems for pressure ulcer care and environments that were able to perform pressure ulcer treatments during hospitalization. Fifteen hospitals finally were identified to be eligible facilities.

Patients and randomization

For patient registration, we set the following inclusion and exclusion criteria: patients aged 50 or over with one or more NPUAP Stage II or III pressure ulcers on either their torso or trochanter, body temperature of 35.5°C min. to 37.5°C max., 600kcal or over daily intake, no critical nutritional impairment, renal failure, cirrhosis, immunosuppression, uncontrollable diabetes, or malignant tumours according to an examination performed within past four weeks. End-of-life patients whose estimated alive period were less than three months were excluded. When the patient did not possess the ability to make the autonomous decision to participate in the study, the outline and methods of the research were explained to their representing immediate family member, from whom written consent was obtained. After obtaining written consent, then uncoordinated random allocations were performed at the registration centre in the research office within 24 hours.

Prospective, Randomized, Open, Blinded-Endpoint design was applied for this study because blinding two different interventions was impossible in clinical setting. We adopted non-stratified pure random allocation for randomization of the enrolled patients. If health staff obtained informed consent from eligible patients, then they send fax to the allocation centre located at Higashi-Washinomiya Hospital with basic information including location and stage of pressure ulcers. The allocation centre finally decided eligibility and inform

1
2
3 registration confirmation and allocation results to the facilities by fax within 48 hours of the
4
5 reception.
6
7

8 Interventions

9
10
11
12 Wrap therapy was defined as a method of treatment that uses food wraps and perforated
13 polyethylene as a wound dressing. In many cases, wrap therapy follows the procedure of
14 thoroughly irrigating the pressure ulcers (Fig. 1-1) before directly covering the pressure
15 ulcers with food wrap if the exudation is small, or with commercially available perforated
16 polyethylene sheets and diapers, or with a cover sheet combined with sanitary towels if the
17 exudation is great (Figs. 1-2 and 1-3). Irrigation and the covering process are performed every
18
19 day²¹.
20
21
22
23

24 To standardize the categories and treatment procedures of wrap therapy as much as possible,
25 a meeting for instruction of interventions was held beforehand. A DVD of the lecture was
26 distributed to the physicians that were unable to participate in the instruction meeting.
27

28 Patients allocated to the control group were treated using methods conforming the 'Evidence-
29 Based Localized Pressure Ulcer Treatment Guidelines' issued by JSPU⁷. These guidelines are
30 diagnosis and treatment guidelines created by JSPU based on current diagnosis and
31 treatment evidence.
32
33
34
35
36
37

38 Outcomes evaluated

39
40
41
42 The primary endpoint was defined as the period from the start of registration until the
43 pressure ulcers that had been the subject of observation for three months were healed. After
44 the patient had been discharged from the hospital, continuous observation was performed as
45 far as possible through diagnosis and treatment at home. Every wound heal was confirmed
46
47 by supervising physicians. As the secondary endpoint, we measured the PUSH score for the
48 localized status of the pressure ulcers as defined by NPUAP at two, four, six, eight, ten and 12
49 weeks from the start of registration²².
50
51
52
53
54
55
56
57
58
59
60

Method for blinded end point evaluation

To avoid the occurrence of observation bias, we established an 'outcome evaluation centre'. Digital camera images were sent from every facility to the outcome evaluation centre, and all outcomes were evaluated there. The outcome evaluation centre was blinded so as not to know the treatment group to which the subject belonged. Because it was impossible to evaluate the exudates amount using images, records were made by the observer directly. **In case the outcome evaluation centre found some problems that excuse the leakages of blindness, then that was reported to the administration office immediately.**

We also recorded the total amount of ointment used in localized treatments during the observation period, and performed rough calculations with regards to cost.

Adverse Events and Role of the Patient Safety Monitoring Board

A safety evaluation committee was established comprising several specialists and non-specialists. The members were independent from the patient registration, research offices and the outcome evaluation centre. In all cases of death, worsening of the pressure ulcers during the research period, systemic deterioration or sepsis occurring within 30 days from the day of the protocol treatment, and when the connection to the pressure ulcers was undeniable, researchers sharing the research at the core facilities reported to the committee.

Statistical Analysis

We aimed to clarify our statistical hypothesis that wrap therapy was not inferior compared to conventional treatments conforming to the guidelines. The mean period until healing for the localized treatment of Stage II pressure ulcers was set as 21 days based on previous literature⁹⁻¹⁴. The non-inferior threshold was set at seven days according to clinical judgment. When set to a tolerable threshold difference of 7 days when one side has a 5% significance level and a test power of 90%, the required number of cases in the two cohorts was 80. Similarly, the required number of cases for Stage III pressure ulcers was 60. For the total number of target cases, 140 was set as the number of target registered patients. The analysts were blinded about which group was the wrap therapy or the conventional treatment until

1
2
3 the analysis was finished. Intention to treat analysis was performed.

4
5 For the main endpoint comparisons, a Kaplan-Meier plots were created and the estimated
6 mean value until the endpoint occurrence and its 95% confidence interval (CI) were calculated.
7
8 The differences in the PUSH scores were calculated from two weeks immediately after the
9 start of observations, between two to four weeks, four to six weeks, six to eight weeks, eight to
10 ten weeks and ten to 12 weeks, and described the speed of pressure ulcer healing over time for
11 both groups. We used PASW Statistics 18 (SPSS, Inc) for the statistical analysis.
12
13
14
15
16
17
18

19 RESULTS

20
21
22 **We started recruitments in October, 2009 and followed until May, 2010.** The study flow
23 diagram is shown in Fig. 2. During the research period, a total of 66 patients participated. Of
24 these, two could not be followed up due to early discharge or transfer to another hospital. A
25 total of 64 participants were analysed: 29 in the conventional treatment group and 35 in the
26 wrap therapy group. The percentage of patients whose endpoints could be pursued four weeks
27 after registration was 95%, eight weeks after registration was 77% and 12 weeks after
28 registration was 64%. The characteristics distribution of the patients analysed was 33 females
29 and 29 males, and the locations of the target pressure ulcers were most common on the
30 sacrum (56%), followed by the trochanter (13%), gluteus (6%) and the coccyx (5%).
31
32
33
34
35
36
37

38 Table 1 shows a comparison of the patients' characteristics distribution, their health at
39 registration, the status of their pressure ulcers and the treatment method for the two groups.
40 The mean values in the conventional treatment and wrap therapy cohorts were 12.7 versus
41 12.7 for the Braden Scale at registration²³. The percentages using hydrocolloids, hydrogels or
42 polyurethane foam as pressure ulcer dressings were 76% in the control group and 3% in the
43 intervention cohort.
44
45
46
47
48
49

50 Time until the Pressure Ulcers Healed

51
52
53
54 The percentage of patients in whom healing of the pressure ulcers was verified from among
55 the samples that could be followed up at four weeks after registration were 26% of the
56 conventional group as opposed to 21% in the wrap therapy group; at eight weeks after
57
58
59
60

1
2
3 registration the percentages were 46% in the conventional group as opposed to 52% in the
4 wrap therapy group.

5
6 Fig. 3-1 shows the survival curves of both groups. The mean of the estimated value until
7 healing was 57.5 days (95% CI 45.2 days; 69.8 days) for the conventional group as opposed to
8 59.8 days (95% CI 49.7 days; 69.9 days) for the wrap therapy group. Fig. 3-2 and Fig. 3-3
9 describes the survival curve for the patients with NPUAP Stage II and III pressure ulcers,
10 respectively. For Stage II pressure ulcers, the estimated mean value until healing were 16.0
11 days (95% CI 8.1 days; 23.9 days) for the conventional group compared to 18.8 days (95% CI
12 10.3 days; 27.2 days) for the wrap therapy group. Meanwhile, for Stage III pressure ulcers,
13 the estimated mean value until healing were 71.8 days (95% CI 61.4 days; 82.3 days) for the
14 conventional group compared to 63.2 days (95% CI 53.0 days; 73.4 days) for the wrap therapy
15 group.
16
17
18
19
20
21
22
23

24 25 26 Speed of Pressure Ulcer Healing Using PUSH Scores

27
28
29 The mean values of the difference in the PUSH scores at registration and upon either healing
30 or the final observation divided by the treatment period were 1.1 ± 2.1 points in the
31 conventional group and 0.9 ± 1.3 points in the wrap therapy group ($p = 0.73$ Student's t-test).
32 No significant difference was identified in the mean PUSH score reduction values in either
33 cohort from immediately after the start of observations to the second week, from the second to
34 the fourth week, from the fourth to the sixth week, from the sixth to the eighth week, from the
35 eighth to the tenth week and from the tenth to the twelfth week (Table 2).
36
37
38
39
40
41
42
43

44 45 46 Adverse Events

47 During the total observation period, there was systemic worsening such as pneumonia,
48 occurring immediately after the start of observation in three cases in the conventional group,
49 and four cases in the wrap therapy group. Two cases in the conventional group died of
50 pneumonia and one died from heart failure for a total of three deaths. In the wrap therapy
51 group, one patient died of pneumonia and one of senescence for a total of two deaths.
52 Regarding localized adverse events, there were problems with the covered skin (eczema,
53 maceration, rash, etc.) in the conventional group and six cases in the wrap therapy group.
54
55
56
57
58
59
60

DISCUSSIONS

Statement of principal findings

'Wrap therapy' has already been introduced nationwide in Japanese clinical facilities. The main reasons for its spread in clinical settings are both that wrap therapy is a rational treatment method compared to basic wound-healing therapies, in that it is possible to maintain a lubricated environment without applying stress to the wound^{24,25}, and because compared to complex pressure ulcer treatments that combine various ointments and dressings, it has frequently been recognized empirically in actual clinical settings as having equal or better effects in treating pressure ulcers.

Looking only at the main outcome survival curves, and taking all the patients into consideration, almost identical results were obtained in both the wrap therapy and conventional treatment groups for the period until the pressure ulcers healed. Meanwhile, when the two treatment groups are compared using the NPUAP classification, although no clear difference between the survival curves of the two treatments were identified with shallow Stage II pressure ulcers, in the comparison between the Stage III pressure ulcer cohorts, the survival curve of the wrap therapy cohort showed that healing tended to be faster than in the conventional treatment cohort. No statistical significance, however, was presented in this study. We can't state so far that wrap therapy should be chosen as the first recommended therapy to shorten the healing time of pressure ulcer. Further study with bigger sample is needed to ensure these visual differences in survival curves. Nor was any significant difference identified between the wrap therapy and conventional treatment cohorts regarding the extent of PUSH score reduction either, and this result can also be stated to support the fact that wrap therapy is not inferior to conventional treatments. Nevertheless, initially we estimated 80 cases and 60 cases in the NPUAP classification subgroups as the non-inferior estimate, but on this occasion, the total effective sample, at 64 cases, lacked sufficient statistical power.

During the comparison of adverse events, there were concerns that skin problems caused by food wraps and perforated polyethylene used in the wrap therapy might occur, but in our research, no significant difference was identified with the results of the conventional treatments. Rashes due to adhesive plasters and tape were identified in four cases in the

1
2
3 conventional cohort, but none were observed in the wrap therapy cohort. This is thought to be
4 because it is basically possible to implement wrap therapy merely by making the patient wear
5 paper diapers without affixing tape after applying the dressing.
6
7
8

9 10 Strengths and weakness of the study 11

12
13
14 This research is the world's first randomized controlled trials that comparing conventional
15 treatments and wrap therapy. Theoretical superiority of wrap therapy to conventional
16 treatments conforming to guidelines is the simplicity and cost effectiveness of the
17 treatment^{20,26}. Conventional treatments require the use of multiple ointments and dressings
18 of suitable size and type, and the treatment requires a certain level of specialist skills. The
19 simplicity of treatment protocols using wrap therapy might better promote the spread of care
20 skills. Regarding cost effectiveness, the various ointments and dressings such as hydrocolloid
21 used in pressure ulcer treatments are often expensive, but the perforated polyethylene and
22 diaper sheets used in wrap therapy can be supplied at less than ten cents per treatment. If
23 the accelerated pressure ulcer healing effects of wrap therapy are the same as those of
24 conventional treatments, the efficiency of wrap therapy is presumably extremely great when
25 considering cost effectiveness. Larger scale additional tests strengthening the verified
26 backing of these research results could offer new, simple and effective methods of pressure
27 ulcer care and noticeably improve the benefits to both patients and society.
28
29

30
31 This research, on the other hand, has some limitations. The greatest concern was
32 impossibility of blindness of the treatment. Without blindness, health staff may have biases
33 on the treatments themselves and wound evaluations. To minimize observation bias, we
34 attempted to eliminate bias occurring during the evaluations by combining all the outcomes
35 in a single outcome evaluation centre as far as possible. The second limitation resulted in bias
36 among the participating facilities. Wrap therapy is currently viewed as an alternative
37 treatment of choice. The facilities that participated in this research currently use wrap
38 therapy clinically under this present condition, and the possibility of wrap therapy effects
39 being overvalued beforehand cannot be denied.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Practical applications

When we consider clinical application of wrap therapy, we should pay some attentions. First, because wrap therapy is simple and inexpensive method, it may be accepted for practical application in home care and primary care setting. Many primary care physicians and nurse practitioners are not familiar with complex protocols using new-coming modern dressings, which have additional advantages for quality of life for patients, and wrap therapy may be a useful application as an initial intervention for pressure ulcers. If wrap therapy is not effective, then using other modern dressings should be considered. Second, this attractive method has ethical concerns. Wrap therapy is not be approved as a formal medical intervention which is covered social health insurance in Japan. Because this therapy uses food wrapping sheet without sterilization, approval as an insured health intervention by Japanese Ministry of Health, Labour and Welfare is difficult. All devises for wrap therapy are so far prepared by health providers as voluntary actions. Even this therapy is useful for home-care setting, it should be examined under supervision of well-trained health professionals.

Meaning of the study

Even though our conclusions failed empirical statement of effectiveness of wrap therapy directly, the results would not deny actual situations in current Japanese clinical settings. In future, as the rapid aging of the population progresses worldwide, the problem of caring for the frail elderly in Japan, which has the highest rate of population aging in the world, will become a highly compelling problem in the societies of Europe and North America. Wrap therapy may be an epoch-making method of treatment and the verification of its usefulness by clinical experiment is thought to be a major advance for the future elderly care.

Acknowledgments

The authors would like to thank the following volunteering facilities for registering patients for this study: Tokatsu-Clinic Hospital, Wakaba Ostomy Centre, Sozokai Heiwadai Hospital, Tokushima Kensei Hospital, Ryumondo Ono Hospital, Kojinkai Hoshigaura Hospital, Ibi Kosei Hospital, Niimi Central Hospital, Oyama Municipal Hospital, Hakuai Kaita Hospital, Yatsu Hoken Hospital, Kikai Tokushukai Hospital, Naruto Yamakami Hospital, Higashi Washinomiya Hospital, and Kaneda Hospital. Shigeru Ichioka and Yuichiro Kume of the Orthopedic Surgery Department, Saitama Medical University for being in charge of the outcome evaluation centre and evaluating the images of pressure ulcers. Toshiko Yamana of the Human Care Studies Department at Tohto College of Health Sciences, Shingo Okada at Kitamihara Clinic, and Yoshiaki Ito at Kiyosu Clinic for supervising the Safety Evaluation Committee, and Megumi Matsubara and Miki Hirano at Higashi Washinomiya Hospital.

Trial registration: UMIN Clinical Trials Registry UMIN000002658. Summary protocol is available on

<https://upload.umin.ac.jp/cgi-bin/ctr/ctr.cgi?function=brows&action=brows&type=detail&recptno=R000003235&admin=0&language=J>

Contributors: SB conducted literature review, developed the initial protocol, examined statistical analysis and drafted the manuscript. AM wrote grant application, recruited collaboration facilities, developed intervention protocol and planned research meeting and startup meeting. SO contributed to research design, reviewed and amended the research protocol including intervention protocol and contributed to enrolment of patients. KT also contributed to research design, reviewed and amended the research protocol and contributed to enrolment of patients. MS contributed to **basic concept of** research design, reviewed and amended the research protocol. KA contributed to enrolment of patients and **acquisition of data**. KK contributed to research design and enrolment of patients. KM managed an outcome evaluation centre. **All authors meet ICMJE authorship criteria for inclusion.**

Funding: This study was supported by Division of the Health for the Elderly at Japanese Ministry of Health, Labor, and Welfare. Grant name 'Examination and Research Work into

1
2
3 New Pressure Ulcer Treatments for the Care of the Elderly'. Funding agency did not interfere
4 with any process of study content. All researchers are independent from the founder. No
5 funding was received from any specific commercial company.
6
7
8

9
10 Competing interests: All authors have completed the Unified Competing Interest form at
11 www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and
12 declare: no support from any organisation for the submitted work, no financial relationships
13 with any organisations that might have an interest in the submitted work in the previous
14 three years, and no other relationships or activities that could appear to have influenced the
15 submitted work.
16
17
18
19

20
21 Ethical approval: The Japanese Academy of Family Medicine Ethics Review Board examined
22 and approved our research plan and informed consent process in October, 2009.
23
24
25
26

27
28 Data sharing: no additional data available
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Patient Characteristics and Health Status, and Pressure Ulcer Status at Registration
(n=64)

	Conventional Treatment Cohort (n=29)	Wrap Therapy Cohort (n=35)	P value
Mean Age (years)	82±10	81±12	0.60
Sex (Female Percentage)	48	54	0.28
<u>Nutrition Absorption Status</u> (Each %)	52	49	0.27
Oral Nutrition	0	9	
Enteral Alimentation: Nasally	38	40	
Enteral Alimentation: Gastrostoma	10	3	
Central Venous Nutrition			
Use of Pressure-Resistant Diffusion Mattress (Percentage 'Yes')	93	100	0.20
Use of Diapers (Percentage 'Yes')	90	91	0.97
<u>Depth of Target Pressure Ulcer</u>			
NPUAP Class II (%)	28	11	0.09
NPUAP Class III (%)	72	89	
Pressure Ulcer Pockets (% Present)	38	34	0.28
<u>Calories Absorbed (Each %)</u>			
Over 1200kcal	45	49	0.58
800kcal to 1200kcal	45	37	
600kcal to 800kcal	10	9	
Serum Albumin (Mean ± SD)	2.8±0.5	2.9±0.5	0.61
Serum Creatinine (Mean ± SD)	0.66±0.3	0.64±0.3	0.77
Braden Scale at Registration	12.8±3.5	12.7±2.8	0.89


(Mean ± SD)			
PUSH Score at Registration	10.8±2.6	10.7±2.7	0.91
(Mean ± SD)			
Pressure Ulcer Surface Area at Registration (Mean ± SD)	14±21	15±25	0.79
Use of ointments or sprays including pharmaceuticals with tissue regeneration accelerant actions	21	14	0.006
Percentage using hydrocolloids, hydrogels, or polyurethane foam as a pressure ulcer dressing	76	3	<0.0001

We used Student t-test for comparison of mean values of the two groups. Chi-square test was adopted for comparisons of frequency.

Table 2: Comparison of the mean reduction in PUSH scores for both cohorts for two weeks immediately after the start of observations, between two to four weeks, four to six weeks, six to eight weeks, eight to ten weeks and ten to 12 weeks, respectively. See separate appendix.

	0 to 2W	2W to 4W	4W to 6W	6W to 8W	8W to 10W	10W to 12W
n	54	45	34	25	18	11
Control: Mean [95% CI]	1.8 [1.0-2.6]	0.4 [0.0-0.9]	0.7 [0.1-1.3]	0.8 [0.1-1.5]	0.3 [-0.1-0.7]	1.0 [0.1-1.9]
Wrapp therapy: Mean [95% CI]	1.8 [1.0-2.6]	0.4 [-0.2-1.0]	0.1 [-0.4-0.6]	0.3 [-0.2-0.8]	0.4 [-0.2-1.0]	0.5 [-0.1-1.1]
P value*	0.77	0.44	0.24	0.43	0.79	0.54

* tested by Mann-Whitney U test




Review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


Fig. 3

Fig. 3-1: Comparison of Survival Curves Using the Period until Pressure Ulcers Healing as the Endpoint - All Cases




Y axis means the proportion of patients that has not been confirmed healing of pressure ulcers. The distribution curves represent the results of an intention-to-treat survival analysis involving all patients in Fig. 3-1, NPUAP stage II patients in Fig. 3-2 and NPUAP stage III patients in Fig. 3-3.

Fig. 3-2: Comparison of Survival Curves Using the Period until Pressure Ulcers Healing as the Endpoint - NPUAP Class II Pressure Ulcers Cohort



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 3-3: Comparison of Survival Curves Using the Period until Pressure Ulcers Healing as the Endpoint - NPUAP Class III Pressure Ulcers Cohort



Review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Brem H, Lyder C. Protocol for the successful treatment of pressure ulcers. *Am J Surg*. 2004; **188**(1A Suppl): 9-17.
2. Orlando PL. Pressure ulcer management in the geriatric patient. *Ann Pharmacother*. 1998; **32**(11): 1221-7.
3. Smith DM. Pressure ulcers in the nursing home. *Ann Intern Med*. 1995; **123**(6): 433-42.
4. Agency for Health Care Policy and Research. *Pressure ulcer treatment: Clinical Practice Guideline Quick Reference Guide*. 1994; (15): 1-25.
5. Cullum N, Nelson EA, Flemming K, et al. Systematic reviews of wound care management: (5) beds; (6) compression; (7) laser therapy, therapeutic ultrasound, electrotherapy and electromagnetic therapy. *Health Technol Assess*. 2001; **5**(9): 1-221.
6. McInnes E, Bell-Syer SE, Dumville JC, et al. Support surfaces for pressure ulcer prevention. *Cochrane Database Syst Rev*. 2008; (4): CD001735.
7. Japanese Society of Pressure Ulcers. *Evidence-Based Localized Pressure Ulcer Treatment Guidelines*. Tokyo: Shorinsya Co.; 2005.
8. Adunsky A, Ohry A. Decubitus direct current treatment (DDCT) of pressure ulcers: results of a randomized double-blinded placebo controlled study. *Arch Gerontol Geriatr*. 2005; **41**(3): 261-9.
9. Hollisaz MT, Khedmat H, Yari F. A randomized clinical trial comparing hydrocolloid, phenytoin and simple dressings for the treatment of pressure ulcers [ISRCTN33429693]. *BMC Dermatol*. 2004; **4**(1): 18.
10. Landi F, Aloe L, Russo A, et al. Topical treatment of pressure ulcers with nerve growth factor: a randomized clinical trial. *Ann Intern Med*. 2003; **139**(8): 635-41.
11. Graumlich JF, Blough LS, McLaughlin RG, et al. Healing pressure ulcers with collagen or hydrocolloid: a randomized, controlled trial. *J Am Geriatr Soc*. 2003; **51**(2): 147-54.
12. Belmin J, Meaume S, Rabus MT, et al. Sequential treatment with calcium alginate dressings and hydrocolloid dressings accelerates pressure ulcer healing in older subjects: a multicenter randomized trial of sequential versus nonsequential treatment with hydrocolloid dressings alone. *J Am Geriatr Soc*. 2002; **50**(2): 269-74.
13. Seeley J, Jensen JL, Hutcherson J. A randomized clinical study comparing a

- hydrocellular dressing to a hydrocolloid dressing in the management of pressure ulcers. *Ostomy Wound Manage.* 1999; **45**(6): 39-44, 6-7.
14. Heyneman A, Beele H, Vanderwee K, et al. A systematic review of the use of hydrocolloids in the treatment of pressure ulcers. *J Clin Nurs.* 2008; **17**(9): 1164-73.
15. Toriyabe S, Saito H, Sakurai K. Use of a food wrap as a dressing material. *Advances in Wound Care.* 1999; **12**(8): 405-6.
16. Mizuhara A. Treatment results and problem of Wrap therapy. *Japanese Journal of Pressure Ulcers.* 2009; **11**(3): 286-7.
17. Okamura S, Hisatomi M, Iwasaki K, et al. *Japanese Journal of Pressure Ulcers.* 2002; **4**(3): 427-30.
18. Toriyabe S, Suemaru S. A study of food-wrap therapy for Stage III and Stage IV pressure ulcers. *Nippon Ishikai Zasshi.* 2000; **123**(10): 1605-11.
19. Bailey KA. Wrap for leg ulcers. *Am J Nurs.* 1996; **96**(6): 17.
20. Ohnishi S, Koike T, Shiogama K, et al. Use of food wrap films for treating pressure ulcers. *Japanese J of Med and Pharmaceutical Science.* 2006; **55**(4): 561-7.
21. Takahashi J, Yokota O, Fujisawa Y, et al. An evaluation of polyvinylidene film dressing for treatment of pressure ulcers in older people. *J Wound Care.* 2006; **15**(10): 449-50, 52-4.
22. Woodbury MG, Houghton PE, Campbell KE, et al. Pressure ulcer assessment instruments: a critical appraisal. *Ostomy Wound Manage.* 1999; **45**(5): 42-5, 8-50, 3-5.
23. Lewicki LJ, Mion L, Splane KG, et al. Patient risk factors for pressure ulcers during cardiac surgery. *AORN J.* 1997; **65**(5): 933-42.
24. Ozturk F, Ermertcan AT. Wound healing: a new approach to the topical wound care. *Cutan Ocul Toxicol.* 2011; **30**(2):92-9.
25. Bishop SM, Walker M, Rogers AA, et al. Importance of moisture balance at the wound-dressing interface. *J Wound Care.* 2003; **12**(4): 125-8.
26. Schuurman JP, Schoonhoven L, Defloor T, et al. Economic evaluation of pressure ulcer care: a cost minimization analysis of preventive strategies. *Nurs Econ.* 2009; **27**(6): 390-400, 15.

CONSORT Statement 2006 - Checklist for Non-inferiority and Equivalence Trials



Items to include when reporting a non-inferiority or equivalence randomized trial

PAPER SECTION And topic	Item	Descriptor	Reported on Page #
TITLE & ABSTRACT	1	<u>How participants were allocated to interventions</u> (e.g., "random allocation", "randomized", or "randomly assigned"), specifying that the trial is a non-inferiority or equivalence trial.	1-2
INTRODUCTION Background	2	<u>Scientific background and explanation of rationale</u> , including the rationale for using a non-inferiority or equivalence design.	4
METHODS Participants	3	<u>Eligibility criteria for participants</u> (detailing whether participants in the non-inferiority or equivalence trial are similar to those in any trial(s) that established efficacy of the reference treatment) and the <u>settings and locations where the data were collected</u> .	5
Interventions	4	<u>Precise details of the interventions intended for each group</u> detailing whether the reference treatment in the non-inferiority or equivalence trial is identical (or very similar) to that in any trial(s) that established efficacy, <u>and how and when they were actually administered</u> .	5-6
Objectives	5	<u>Specific objectives and hypotheses</u> , including the hypothesis concerning non-inferiority or equivalence.	4
Outcomes	6	<u>Clearly defined primary and secondary outcome measures</u> detailing whether the outcomes in the non-inferiority or equivalence trial are identical (or very similar) to those in any trial(s) that established efficacy of the reference treatment and, when applicable, any <u>methods used to enhance the quality of measurements</u> (e.g., multiple observations, training of assessors).	6
Sample size	7	<u>How sample size was determined</u> detailing whether it was calculated using a non-inferiority or equivalence criterion and specifying the margin of equivalence with the rationale for its choice. When applicable, <u>explanation of any interim analyses and stopping rules</u> (and whether related to a non-inferiority or equivalence hypothesis).	7
Randomization -- Sequence generation	8	<u>Method used to generate the random allocation sequence</u> , including details of any restrictions (e.g., blocking, stratification)	5
Randomization -- Allocation concealment	9	<u>Method used to implement the random allocation sequence</u> (e.g., numbered containers or central telephone), clarifying whether the sequence was concealed until interventions were assigned.	5
Randomization -- Implementation	10	<u>Who generated the allocation sequence, who enrolled participants, and who assigned participants to their groups</u> .	5
Blinding (masking)	11	<u>Whether or not participants, those administering the interventions, and those assessing the outcomes were blinded to group assignment</u> . If done, <u>how the success of blinding was evaluated</u> .	7
Statistical methods	12	<u>Statistical methods used to compare groups for primary outcome(s)</u> , specifying whether a one or two-sided confidence interval approach was used. <u>Methods for additional analyses</u> , such as subgroup analyses and adjusted analyses.	7-8
RESULTS Participant flow	13	<u>Flow of participants through each stage</u> (a diagram is strongly recommended). Specifically, for each group report the numbers of participants randomly assigned, receiving intended treatment, completing the study protocol, and analyzed for the primary outcome. <u>Describe protocol deviations from study as planned, together with reasons</u> .	20
Recruitment	14	<u>Dates defining the periods of recruitment and follow-up</u> .	8
Baseline data	15	<u>Baseline demographic and clinical characteristics of each group</u> .	8
Numbers analyzed	16	<u>Number of participants (denominator) in each group included in each analysis and whether the analysis was "intention-to-treat" and/or alternative analyses were conducted</u> . State the results in absolute numbers when feasible (e.g., 10/20, not 50%).	8

Outcomes and estimation	17	<u>For each primary and secondary outcome, a summary of results for each group, and the estimated effect size and its precision (e.g., 95% confidence interval). For the outcome(s) for which non-inferiority or equivalence is hypothesized, a figure showing confidence intervals and margins of equivalence may be useful.</u>	8,9,15-17,21
Ancillary analyses	18	<u>Address multiplicity by reporting any other analyses performed, including subgroup analyses and adjusted analyses, indicating those pre-specified and those exploratory.</u>	9,22,23
Adverse events	19	<u>All important adverse events or side effects in each intervention group.</u>	9
<i>DISCUSSION</i> Interpretation	20	<u>Interpretation of the results</u> , taking into account the <i>non-inferiority or equivalence hypothesis and any other study hypotheses</i> , sources of potential bias or imprecision and the dangers associated with multiplicity of analyses and outcomes.	10-11
Generalizability	21	<u>Generalizability (external validity) of the trial findings.</u>	12
Overall evidence	22	<u>General interpretation of the results in the context of current evidence.</u>	12

www.consort-statement.org

Fig. 1-1



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Fig. 1-2



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Fig. 1-3



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49