

Quality of medical care and excess mortality in psychiatric patients—a nationwide register-based study in Sweden

Emma Björkenstam,^{1,2} Rickard Ljung,^{1,2,3} Bo Burström,¹ Ellenor Mittendorfer-Rutz,⁴ Johan Hallqvist,^{5,6} Gunilla Ringbäck Weitoff²

To cite: Björkenstam E, Ljung R, Burström B, *et al.* Quality of medical care and excess mortality in psychiatric patients—a nationwide register-based study in Sweden. *BMJ Open* 2012;**2**: e000778. doi:10.1136/bmjopen-2011-000778

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2011-000778>).

Received 7 January 2012
Accepted 27 January 2012

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

ABSTRACT

Objective: To assess overall and cause-specific mortality and the quality of somatic care among psychiatric patients.

Design: A register-based cohort study.

Setting: All individuals aged 20–79 years in Sweden in 2005.

Participants: In total 6 294 339 individuals.

Primary outcome measure: The individuals were followed for mortality in 2006 and 2007, generating 72 187 deaths. Psychiatric patients were grouped according to their diagnosis in the National Patient Register. Mortality risk of psychiatric patients was compared with that of non-psychiatric patients. Estimates of RR of mortality were calculated as incidence rate ratios (IRRs) with 95% CIs using Poisson regression analysis. Psychiatric patients were compared with non-psychiatric patients for three healthcare quality indicators: the proportion of avoidable hospitalisations, case death rate after myocardial infarction and statin use among diabetic patients.

Results: Compared with individuals with no episodes of treatment for mental disorder, psychiatric patients had a substantially increased risk of all studied causes of death as well as death from conditions considered amenable to intervention by the health service, that is, avoidable mortality. The highest mortality was found among those with another mental disorder, predominantly substance abuse (for women, an IRR of 4.7 (95% CI 4.3 to 5.0) and for men, an IRR of 4.8 (95% CI 4.6 to 5.0)). The analysis of quality of somatic care revealed lower levels of healthcare quality for psychiatric patients, signalling failures in public health and medical care.

Conclusion: This study shows a marked increase in excess mortality, suggesting a lower quality of somatic healthcare in psychiatric patients.

ARTICLE SUMMARY

Article focus

- To investigate whether psychiatric patients have higher overall and cause-specific mortality.
- To study whether the quality of somatic care for psychiatric patients is different from the care provided to patients with no concurrent mental disorder.

Key messages

- This study shows a marked increase in excess mortality in mentally ill men and women.
- The findings suggest a lower quality of health-care in the treatment of somatic disorders in psychiatric patients regarding the indicators studied.
- Careful medical examination of psychiatric patients together with efforts to promote a healthier life style may be of great importance in order to prevent, detect and treat somatic disease.

Strengths and limitations of this study

- The strengths of the study include the population-based design, using national registers with high completeness.
- We were unable to study the effects of health-related behaviours such as excessive alcohol consumption or tobacco smoking, which is a limitation.

INTRODUCTION

Psychiatric patients have been reported to have higher mortality rates and shorter life expectancy compared with the general population.^{1–6} This applies especially to patients with severe mental disorders, such as schizophrenia^{7–10} and bipolar disorder.^{11–12} The highest risks have been found among patients with substance abuse.^{3 5 13 14}

Psychiatric patients have a particularly pronounced risk of suicide.^{15–18} Cardiovascular disease is also a major cause of excess

For numbered affiliations see end of article.

Correspondence to

Emma Björkenstam;
emma.bjorkenstam@ki.se

death, partly due to a higher prevalence of smoking, obesity and hypertension.^{19–21} There are conflicting data on cancer incidence and mortality in psychiatric patients, although most studies suggest that, while cancer mortality is higher, incidence is no different from that of the general population.^{22–23} Other potential explanations for the excess mortality are differences in access, provision and quality of somatic healthcare.²⁴ For instance, it has been shown that people with a diagnosis of a mental disorder have less access to primary healthcare.²⁵ Lower quality of diabetes care²⁶ and cardiovascular procedures^{19–27–28} have also been documented. In Sweden, different indicators are often used as measures reflecting healthcare quality.²⁹

The concept of avoidable mortality was introduced in 1976 as an indicator of quality of healthcare.³⁰ Avoidable mortality measures deaths that could be argued to have been avoidable either by the healthcare system or by public health interventions and has been applied in comparisons between regions, populations and over time. Psychiatric patients have been shown to have higher avoidable mortality rates than the general population,^{13–31} and questions have been raised as to whether the medical care of physical disorders provided to psychiatric patients is less adequate than for the population in general.³²

The aim of this study was to investigate whether psychiatric patients have higher overall and cause-specific mortality and whether the quality of somatic care for psychiatric patients is different from the care provided to patients with no concurrent mental disorder.

Results are reported with regard to age, sex, mental disorder and comorbid substance abuse disorders.

METHODS

Study population

The study population comprised individuals alive and registered in Sweden in 2004 and 2005, aged 20–79 years, that is, those born between 1926 and 1985. After excluding individuals diagnosed with mental retardation (as defined by the International Classification of Diseases, tenth revision (ICD-10): F70–F79), the study population comprised 6 294 339 individuals. The cohort was followed from 1 January 2006 until 31 December 2007. We used the unique personal identity number assigned to each Swedish citizen or permanent resident to link information from four population-based registers.³³

Types of mental disorder

We identified all individuals recorded with a principal psychiatric diagnostic code as defined by ICD-10 F04–F99 (ie, dementia excluded, besides mental retardation) in inpatient care or specialised outpatient care in 2004 or 2005 in the National Patient Register. The National Patient Register covers all inpatient care in Sweden since 1987 (psychiatric inpatient care since 1973, where both alcohol and drugs services are

included) and all specialised outpatient care since 2001. If a patient had more than one discharge or outpatient visit, the principal diagnoses from all discharges and outpatient visits were kept to further categorise the patients. Five mutually exclusive exposure groups were created according to a hierarchy of diagnoses: (1) schizophrenia and other non-affective psychoses (diagnosis codes F20–F29), (2) affective disorder (F30–F39), (3) anxiety disorder (F40–F48), (4) other psychiatric diagnoses (F04–F99 except for diagnoses above) and (5) persons without inpatient and outpatient care due to mental disorder served as the reference group.

Psychiatric in- or outpatient care due to substance abuse disorder was also studied in a sub-analysis and was divided into categories according to whether the substance abuse was attributed to alcohol (F10), narcotic-related drugs (F11–F19) or both.

As certain mental disorders, for example, bipolar disorder and schizophrenia, are highly comorbid with alcohol and other substance abuse,³⁴ schizophrenia/other non-affective psychoses, affective or anxiety disorder were analysed with and without a coexisting substance abuse disorder (attributed to either alcohol or narcotics-related drugs). In these analyses, persons without inpatient and outpatient care due to mental disorder served as the reference group.

Causes of death

The underlying causes of death, coded according to ICD-10, were obtained from the Causes of Death Register. This register contains information on all deceased Swedish residents since 1952 and has a very high coverage. Since 1997, all deceased have been included, though for 0.5%, there is a lack of medical information.

Causes of death were grouped as follows: ischaemic heart disease (IHD) (diagnosis codes I20–I25), cancer (C00–C97), suicide and death with undetermined intent (X60–X84 and Y10–Y34) and external causes of death (suicide excluded) (V01–Y89 (except for X60–X84, Y10–Y34)). The indicators of avoidable mortality were divided into causes of death reflecting the outcome of medical care and causes reflecting the effect of the national health policy. The following diseases were included as medical care indicators: typhoid (A01.0), tetanus (A35), tuberculosis (A15–A19 and B90), malignant neoplasm of cervix uteri (C53), Hodgkin's disease (C81), diabetes (E10–E14), chronic rheumatic heart disease (I05–I09), hypertensive disease (I10–I15), stroke (I60–I69), asthma (J45–J46), appendicitis (K35–K38), abdominal hernia (K40–K45), cholelithiasis and cholecystitis (K80–K81 and K83.0), maternal deaths (O00–O99) and osteomyelitis (M86–M87). Malignant neoplasms of the oesophagus (C15); malignant neoplasms of the trachea, bronchus or lung (C34); cirrhosis of the liver (K70 and K73–K74) and motor vehicle accidents (V00–V99) were included as national health policy indicators.

Healthcare quality indicators

In analyses of indicators presumably reflecting aspects of healthcare quality, the age span was restricted to individuals aged 40–79 years because the studied outcomes are rare in younger individuals. The indicator regarding avoidable hospitalisation is based on the assumption that unnecessary hospitalisation can be avoided if patients with selected conditions receive proper outpatient care. This indicator includes hospitalisations for some chronic conditions (anaemia, asthma, diabetic complications, heart failure, hypertension, chronic obstructive lung disease and angina pectoris) and hospitalisations for some acute conditions (bleeding ulcers, diarrhoea, epileptic seizures, inflammatory diseases of the female pelvic organs, pyelitis and ear, nose and throat infection). The indicator regarding 28-day case death rate after myocardial infarction is an internationally established indicator of how well the healthcare system handles acute care after myocardial infarction. As a third indicator, the proportion on treatment with lipid-lowering drugs was measured among patients receiving diabetes drugs. This treatment ought to be especially important for diabetes patients, due to the many risk factors in this population. All three indicators have been described elsewhere.²⁹

Statistical analysis

Poisson regression analyses, adjusted for age, were used to evaluate the association between mental disorder, mortality and indicators of quality of healthcare. As a measure of the relative occurrence of death, we used the incidence rate ratio (IRR). We assessed person-years at risk by adding up the years the individuals were alive and living in Sweden during the follow-up period. SAS Genmod procedure was used to calculate IRRs with 95% CIs. SAS Enterprise Guide 4.2 (SAS Institute Inc.) was used. Age-standardised percentages were calculated when analysing differences regarding healthcare quality.

RESULTS

Cohort characteristics for the 6 294 339 individuals (3 141 454 women and 3 152 885 men) are presented in [table 1](#). In total, 101 500 women and 90 946 men were treated for mental disorder in 2004 or 2005. About 11% of mentally ill women had a diagnosis of schizophrenia and other non-affective psychoses, 37% affective disorders, 32% anxiety disorder and 20% other psychiatric diagnoses and for men 13%, 26%, 23% and 37%, respectively. Almost three-quarters of the men (73%) included in the group for other mental disorders were treated for substance abuse.

Among all psychiatric patients, 5498 (2.9% in all, 2% in women and 4% in men, respectively) died during the follow-up period compared with 66 689 (1.1% in all, 0.9% in women and 1.3% in men) among persons without episodes of treatment for mental disorder.

The IRRs for different causes of death and by mental disorder for both sexes are shown in [table 2](#). When

compared with the general population, mortality was considerably higher among all categories of mental disorder and for all causes of death. The highest excess mortality was found among individuals in the category comprised of having another mental disorder, predominantly substance abuse (for women, an IRR of 4.7 (95% CI 4.3 to 5.0) and for men, an IRR of 4.8 (95% CI 4.6 to 5.0)).

The IRRs for different causes of death presented in [table 3](#) show that, regardless of mental disorder, those with comorbid substance abuse disorders had the highest mortality risk.

Women treated for both alcohol-related and narcotic-related substance abuse had the highest risk for premature death, except for cancer and avoidable mortality ([table 4](#)). Except for cancer, men treated for both alcohol-related and narcotic-related substance abuse had the highest risk for premature death in all studied causes of death.

Separate analyses of different age groups 20–44, 45–64 and 65–79 showed that the younger the patient, the more pronounced increased risk for premature death (data not shown). This was true for all categories of mental disorder and for all studied causes of death, except for suicide.

The majority of patients (78% of women and 77% of men) with affective and anxiety disorders had been treated in outpatient care only. For the other disorders, around half of the patients had been treated in outpatient care only. Mortality rates were higher among those hospitalised than those treated in outpatient care. This was true for all categories of mental disorders and for all causes of death (data not shown).

The different healthcare quality indicators are presented in [table 5](#). The percentage of avoidable admissions was higher among people treated for mental disorder, for women spanning from 2.4% to 4.1% (compared with 1.1% for those without episodes of treatment for mental disorder) and for men spanning from 3.4% to 5.4% (compared with 1.6%). Psychiatric patients had a higher 28-day case death rate for myocardial infarction, especially those treated for schizophrenia and other non-affective psychoses (women 53.5% (42.0–65.0) and men 51.4% (40.4–62.3)) compared with 26.5% of women (25.7–27.3) and 28.1% (27.5–28.6) of men not treated for mental disorder. The indicator on lipid-lowering drug therapy shows that psychiatric patients with diabetes were given lipid-lowering drugs to a lesser extent than diabetic patients without a concurrent mental disorder; this was particularly evident among both women and men with schizophrenia and other non-affective psychoses.

DISCUSSION

Our study of more than 6 million women and men shows that psychiatric patients had a substantially increased risk of death compared with individuals with no episodes of

Table 1 Cohort characteristics

	Women				Men			
	Schizophrenia and other non-affective psychoses				Schizophrenia and other non-affective psychoses			
	Affective disorder	Anxiety disorder	Other mental disorder	No mental disorder	Affective disorder	Anxiety disorder	Other mental disorder	No mental disorder
N	11 165	32 690	20 400	3 039 954	11 993	21 209	33 982	3 061 939
%	0.2	0.5	0.3	48.3	0.2	0.3	0.5	48.6
Mean age (SD)	50.1 (13.6)	42.4 (14.3)	43.8 (16.4)	48.1 (16.0)	45.8 (13.0)	42.0 (13.4)	47.2 (15.2)	47.2 (15.6)
Person-years	21 923	64 788	40 039	6 017 464	23 463	41 790	65 925	6 042 510
Causes of death								
All-cause mortality	344	669	638	27 421	459	389	1822	39 268
Avoidable mortality:	34	68	54	3 125	30	28	140	3 904
medical care indicators								
Avoidable mortality:	22	57	89	2859	34	38	226	4 062
national health policy indicators								
Ischaemic heart disease	42	68	72	3 279	79	68	280	7 878
Cancer (all types)	83	163	111	12 369	65	71	247	13 636
Suicide	42	117	73	439	79	69	174	1 259
Deaths from external causes (suicide excluded)	15	44	47	652	39	48	232	1 867
Healthcare quality indicators*	8413	22 833	11 171	1 961 968	7864	11 126	22 762	1 931 860
1. Avoidable inpatient medical care								
Number treated in 2006	234	521	409	22 433	235	341	1 189	29 159
2. Acute Myocardial Infarction (AMI) Myocardial infarction—28-day case death								
AMI in 2006 or 2007	77	164	134	10 993	105	140	410	23 786
Dead within 28 days	36	62	45	3 108	50	42	171	6 467
3. Lipid-lowering drug therapy								
Number receiving diabetes drug therapy in 2006	722	1 200	726	80 453	744	680	1 744	116 270
Of these, number who were given lipid-lowering drugs in 2006	292	618	385	45 193	318	367	743	65 440
								114 267

*Individuals aged 40–79 years.

Table 2 Associations of mental disorder in relation to mortality (incidence rate ratios* with 95 % CIs)

Mental disorder group	Avoidable mortality:				Suicide	Death from external causes (suicide excluded)
	All-cause mortality	Ischaemic heart disease	Cancer (all types)	medical care indicators		
Women						
No mental disorder	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)
Schizophrenia and other non-affective psychoses	3.8 (3.4–4.2)	4.1 (3.1–5.6)	1.9 (1.5–2.4)	3.5 (2.5–4.9)	2.1 (1.4–3.2)	24.9 (18.1–34.2) 6.7 (4.0–11.1)
Affective disorder	2.4 (2.2–2.6)	2.1 (1.7–2.7)	1.3 (1.1–1.5)	2.2 (1.7–2.8)	1.9 (1.5–2.5)	22.3 (18.1–27.3) 6.1 (4.5–8.3)
Anxiety disorder	2.6 (2.3–2.8)	2.9 (2.1–3.8)	1.7 (1.4–2.0)	2.2 (1.6–3.0)	2.4 (1.8–3.2)	16.0 (12.5–20.6) 5.3 (3.6–7.8)
Other mental disorder (including substance abuse disorder)	4.7 (4.3–5.0)	4.6 (3.6–5.8)	1.8 (1.5–2.2)	3.5 (2.7–4.6)	6.2 (5.0–7.6)	26.7 (20.8–34.3) 12.6 (9.4–17.0)
Men						
No mental disorder	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)
Schizophrenia and other non-affective psychoses	4.6 (4.2–5.0)	4.3 (3.4–5.3)	1.9 (1.5–2.5)	3.4 (2.4–4.9)	3.0 (2.1–4.1)	16.1 (12.9–20.3) 6.2 (4.5–8.6)
Affective disorder	2.8 (2.7–3.1)	2.2 (1.9–2.7)	1.5 (1.2–1.7)	3.0 (2.4–3.8)	1.4 (1.1–2.0)	17.2 (14.6–20.2) 4.9 (3.8–6.2)
Anxiety disorder	2.8 (2.5–3.0)	2.7 (2.1–3.5)	1.5 (1.2–2.0)	2.3 (1.6–3.4)	2.3 (1.7–3.2)	8.3 (6.5–10.6) 4.5 (3.4–6.1)
Other mental disorder (including substance abuse disorder)	4.8 (4.6–5.0)	3.7 (3.3–4.2)	1.9 (1.6–2.1)	3.9 (3.3–4.6)	5.5 (4.8–6.3)	12.6 (10.8–14.8) 11.8 (10.3–13.6)

*Adjusted for age.

treatment for mental disorder, regardless of cause of death. Overall, people with schizophrenia and other non-affective psychoses and people treated for substance abuse disorder had the highest risk for premature death. Psychiatric patients also had higher rates of potentially avoidable hospitalisations, higher case death rate after myocardial infarction and lower use of lipid-lowering therapy in diabetes.

The strengths of the study include the population-based design, using national registers with high completeness. The validity in the Causes of Death Register is very high, where information of cause of death is lacking in only 0.5% of the deceased.³⁵ The National Patient Register has close to complete coverage of all inpatient care. However in 2005, around 40% of psychiatric outpatient visits were not included, hence some patients who were treated in outpatient care are classified as unexposed, as there is no record of their outpatient visit. Also, as the National Patient Register does not cover primary care, we were not able to study all levels of psychiatric care. Hence, patients treated for mental health problems solely in primary care are classified as unexposed. Assuming a similar effect for patients in primary care as for psychiatric patients in our study selected by a more strict definition of mental health problems, that is, patients who have seen a psychiatrist, this misclassification will dilute our results towards the null. We were unable to study the effects of health-related behaviours such as excessive alcohol consumption or tobacco smoking, which is a limitation.

We found that psychiatric patients were at higher risk of premature death from IHD, particularly among those treated for schizophrenia and other non-affective psychoses. This has previously been shown in other settings,^{10 11 19 20} and recently a study showed that the excess cardiovascular mortality among schizophrenia patients has increased over the past 25 years.⁷ We also found an elevated risk for cancer mortality, although to a lesser extent, in particular in patients treated for schizophrenia and other non-affective psychoses and substance abuse disorders. Studies evaluating cancer incidence and mortality at the same time have shown that, while individuals with mental disorder did not show any increased incidence rate for many types of cancer, they did have higher cancer mortality, which might be explained by delays in detection,^{23 36} disparities in screening for cancer²² and treatment compliance.³⁷

Psychiatric patients had an elevated risk of premature death from conditions considered amenable to intervention by the health service, that is, avoidable mortality, compared with those not treated in psychiatric care. In particular, those treated for schizophrenia with comorbid substance abuse disorder had a higher risk, which has been shown by others.^{13 31}

In line with previous studies, we found psychiatric patients to have significantly higher suicide risk, in particular those treated for schizophrenia and other non-affective psychosis, and affective disorder and

Table 3 Associations of mental disorder and substance abuse in relation to mortality (incidence rate ratios* with 95% CIs)

Mental disorder group	N	All-cause mortality	Ischaemic heart disease	Cancer (all types)	Avoidable mortality: medical care indicators	Avoidable mortality: national health policy indicators	Suicide	Death from external causes (suicide excluded)
Women								
No mental disorder	3039954	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)
Schizophrenia and other non-affective psychoses with substance abuse disorder	502	10.8 (7.1–16.6)	5.7 (0.8–40.4)	5.2 (2.2–12.6)	–	21.5 (8.9–51.8)	44.6 (14.3–138.8)	43.1 (13.8–134.1)
Schizophrenia and other non-affective psychoses without substance abuse disorder	10663	3.6 (3.2–4.0)	4.1 (3.0–5.6)	1.8 (1.5–2.3)	3.5 (2.5–5.0)	1.7 (1.0–2.7)	24.2 (17.4–33.6)	5.5 (3.1–9.8)
Affective disorder with substance abuse disorder	1919	6.4 (4.9–8.3)	3.5 (1.1–10.9)	2.7 (1.5–4.8)	3.4 (1.1–10.6)	6.5 (3.1–13.6)	61.1 (37.1–100.6)	24.1 (11.4–50.8)
Affective disorder without substance abuse disorder	35326	2.3 (2.1–2.5)	2.1 (1.6–2.7)	1.2 (1.1–1.4)	2.2 (1.7–2.8)	1.8 (1.3–2.3)	20.3 (16.3–25.2)	5.4 (3.9–7.5)
Anxiety disorder with substance abuse disorder	1551	7.3 (5.4–9.8)	10.9 (4.9–24.3)	2.0 (0.9–4.4)	1.7 (0.2–12.2)	6.8 (2.8–16.4)	63.1 (36.3–109.7)	22.9 (9.5–55.4)
Anxiety disorder without substance abuse disorder	31139	2.4 (2.2–2.7)	2.6 (1.9–3.5)	1.7 (1.4–2.0)	2.2 (1.6–3.0)	2.2 (1.6–3.0)	13.8 (10.5–18.2)	4.5 (3.0–6.9)
Other mental disorder with substance abuse disorder	387	17.0 (10.4–27.7)	–	–	11.3 (1.6–80.4)	–	171.6 (85.1–346.0)	21.6 (3.0–153.5)
Other mental disorder without substance abuse disorder	10773	2.6 (2.3–3.0)	2.9 (2.0–4.1)	1.4 (1.1–1.8)	2.6 (1.8–3.8)	0.9 (0.4–1.8)	10.0 (5.9–17.0)	1.8 (0.7–4.9)
Substance abuse disorder without a psychiatric diagnosis	9240	7.9 (7.1–8.7)	8.4 (6.2–11.4)	2.4 (1.9–3.1)	5.3 (3.6–7.8)	13.4 (10.7–16.7)	39.8 (29.8–53.2)	28.1 (20.5–38.5)
Men								
No mental disorder	3061939	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)
Schizophrenia and other non-affective psychoses with substance abuse disorder	1088	12.5 (9.8–15.9)	4.9 (1.9–13.2)	3.8 (1.7–8.4)	5.3 (1.3–21.1)	11 (5.2–23.1)	41.3 (25.6–66.7)	22.3 (12.3–40.3)
Schizophrenia and other non-affective psychoses without substance abuse disorder	10905	4.2 (3.8–4.6)	4.3 (3.4–5.3)	1.8 (1.4–2.4)	3.4 (2.3–4.9)	2.5 (1.7–3.6)	13.9 (10.7–17.9)	4.9 (3.4–7.1)
Affective disorder with substance abuse disorder	2237	6.7 (5.6–8.1)	5.4 (3.4–8.6)	1.6 (0.9–3.0)	5.8 (3.0–11.2)	5.3 (2.9–9.5)	43.7 (31.8–60.2)	11.3 (6.5–19.4)
Affective disorder without substance abuse disorder	21525	2.6 (2.4–2.8)	2.0 (1.6–2.4)	1.5 (1.2–1.7)	2.8 (2.2–3.6)	1.1 (0.8–1.6)	14.5 (12.1–17.4)	4.3 (3.3–5.6)
Anxiety disorder with substance abuse disorder	1900	9.7 (7.9–11.8)	7.7 (4.5–13.2)	2.5 (1.2–4.9)	2.5 (0.6–10.2)	8.9 (4.9–16.1)	27.7 (17.8–43.1)	24.8 (16.3–37.8)

Continued

Table 3 Continued

Mental disorder group	N	All-cause mortality	Ischaemic heart disease	Cancer (all types)	Avoidable mortality: medical care indicators	Avoidable mortality: national health policy indicators	Suicide	Death from external causes (suicide excluded)
Anxiety disorder without substance abuse disorder	19 309	2.2 (2.0–2.5)	2.4 (1.8–3.1)	1.5 (1.2–1.9)	2.3 (1.6–3.4)	1.8 (1.2–2.6)	6.5 (4.9–8.6)	2.7 (1.8–4.0)
Other mental disorder with substance abuse disorder	380	9.8 (6.7–14.4)	21.0 (11.3–39.0)	2.3 (0.6–9.1)	8.7 (2.2–35.0)	–	47.6 (22.6–100.1)	10.6 (2.7–42.5)
Other mental disorder without substance abuse disorder	9298	2.5 (2.2–2.8)	1.7 (1.3–2.2)	1.4 (1.1–1.8)	3.5 (2.7–4.7)	1.2 (0.7–2.0)	5.4 (3.5–8.4)	2.3 (1.4–3.9)
Substance abuse disorder without a psychiatric diagnosis	24 304	6.1 (5.8–6.4)	4.8 (4.2–5.5)	2.1 (1.8–2.4)	4.0 (3.3–5.0)	7.7 (6.7–8.8)	14.8 (12.5–17.6)	16.1 (14.0–18.6)

*Adjusted for age.

especially those treated with a comorbid substance abuse disorder.¹⁷

Psychiatric patients also had substantially increased risk of deaths from external causes when suicide was excluded. Studies have shown that people with mental disorder are at greater risk of accidental and violent death, particularly when alcohol and narcotic-related drug comorbidities are involved.³⁸ As mentioned earlier, our results showed that mentally ill individuals with a coexisting substance abuse disorder have significantly higher risk for premature death. When we excluded those with a coexisting substance abuse disorder, the increased mortality risk remained for all diagnosis groups except for schizophrenia and other non-affective psychosis, but with lower estimates (data not shown).

Consistent with previous research, substance abuse disorders strongly contributed to premature death among both female and male psychiatric patients, especially for those with both alcohol and narcotic-related drug abuse.

When studying a set of indicators aimed to reflect the outcome of healthcare quality, we found that patients with a mental disorder had higher rates of potentially avoidable hospitalisations. Under some circumstances, hospitalisation is necessary and well motivated, but for most studied diseases, it is a failure. A recent study on avoidable hospitalisation demonstrated that the quality of physical healthcare received by patients with schizophrenia was poorer than that of the general population, signalling failures in public health and medical care.³⁴

The indicator of 28-day case death rate after myocardial infarction showed worse outcome for those with a mental disorder. In a Danish study,¹⁹ less somatic hospitalisation than needed and less use of invasive heart disease procedures among persons with severe mental disorder than among the general population were suggested as additional reasons for their excess mortality in IHD.

Another finding was the low percentage of lipid-lowering drugs given to diabetic patients with a concurrent mental disorder. As drug therapy against lipid disorders is particularly important in diabetic patients,³⁹ this is a clear indication of inequality in healthcare.

Disparities in access to and utilisation of healthcare, as well as the quality of healthcare provision, may contribute to inequalities in health.⁴⁰ One reason for these disparities could be that a person with mental disorder may not be able to effectively communicate and express concerns because of cognitive disturbance. Also, psychiatrists and other mental health providers may prioritise psychiatric issues and neglect physical problems.⁴¹ There is also some evidence of inequalities in case recognition and quality of medical care for psychiatric patients, where general practitioners might minimise the clinical significance of physical complaints when assessing and treating patients with a mental disorder, a process known as diagnostic overshadowing.⁴² As there is excess mortality across a wide range of mental disorders, somatic care should be

Table 4 Associations of substance abuse in relation to mortality (incidence rate ratios* with 95% CIs)

	N	All-cause mortality	Ischaemic heart disease	Cancer (all types)	Avoidable mortality: medical care indicators	Avoidable mortality: national health policy indicators	Suicide	Death from external causes (suicide excluded)
Women								
No substance abuse disorder	3127	855 1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)
Substance abuse disorder, alcohol	8185	7.7 (6.9–8.5)	4.1 (0.6–29.3)	2.3 (1.8–3.0)	4.8 (3.2–7.2)	13.8 (11.1–17.2)	29.5 (22.1–39.4)	22.6 (16.0–32.0)
Substance abuse disorder, narcotic-related drug abuse	4589	6.6 (5.5–8.0)	4.7 (2.3–9.4)	2.7 (1.8–4.1)	3.9 (1.9–8.2)	6.1 (3.5–10.8)	36.9 (25.8–52.8)	22.9 (13.9–37.6)
Substance abuse disorder, both alcohol and narcotic-related drug abuse	825	11.2 (8.0–15.7)	8.7 (6.4–11.8)	2.6 (1.0–6.9)	3.7 (0.5–26.5)	7.8 (2.5–24.1)	54.4 (28.1–105.1)	65.8 (32.7–132.4)
Men								
No substance abuse disorder	3122	976 1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)	1 (REF)
Substance abuse disorder, alcohol	20429	5.5 (5.3–5.9)	4.8 (4.2–5.5)	2.0 (1.7–2.4)	3.8 (3.1–4.7)	7.5 (6.6–8.7)	12.5 (10.4–14.9)	12.8 (10.8–15.0)
Substance abuse disorder, narcotic-related drug abuse	7753	9.4 (8.4–10.6)	5.8 (3.9–8.7)	2.7 (1.7–4.1)	4.9 (2.6–9.1)	4.1 (2.4–7.0)	22.4 (17.7–28.4)	22.5 (18.0–28.3)
Substance abuse disorder, both alcohol and narcotic-related drug abuse	1727	14.5 (12.2–17.2)	6.8 (3.7–12.7)	2.5 (1.2–5.2)	8.7 (3.9–19.4)	16.0 (10.0–25.4)	35.2 (24.2–51.2)	29.3 (19.6–43.9)

*Adjusted for age.

Table 5 Age-standardised percentage (with 95% CIs) as measures of healthcare quality indicators for individuals aged 40–79 years, 2006

Mental disorder group	N	Avoidable inpatient medical care	Myocardial infarction—28-day case death	Lipid-lowering drug therapy
Women				
No mental disorder	1 961 968	1.1 (1.1 to 1.1)	26.5 (25.7 to 27.3)	56.1 (55.7 to 56.4)
Schizophrenia and other non-affective psychoses	8413	2.9 (2.6 to 3.3)	53.5 (42.0 to 65.0)	40.9 (36.8 to 45.1)
Affective disorder	22 833	2.4 (2.2 to 2.6)	34.8 (26.5 to 43.2)	51.9 (49.0 to 54.8)
Anxiety disorder	17 078	3.0 (2.7 to 3.3)	36.1 (28.8 to 43.4)	54.9 (50.9 to 58.8)
Other mental disorder (including substance abuse disorder)	11 171	4.1 (3.7 to 4.5)	45.0 (35.8 to 54.3)	49.0 (44.8 to 53.1)
Men				
No mental disorder	1 931 860	1.6 (1.5 to 1.6)	28.1 (27.5 to 28.6)	56.2 (55.9 to 56.5)
Schizophrenia and other non-affective psychoses	7864	3.8 (3.3 to 4.4)	51.4 (40.4 to 62.3)	41.9 (37.2 to 46.5)
Affective disorder	15 502	3.4 (3.1 to 3.7)	39.2 (32.6 to 45.7)	53.8 (50.8 to 56.8)
Anxiety disorder	11 126	4.4 (3.9 to 4.9)	32.5 (23.1 to 42.0)	56.4 (51.9 to 60.8)
Other mental disorder (including substance abuse disorder)	22 762	5.4 (5.1 to 5.7)	43.9 (38.6 to 49.1)	44.2 (41.7 to 46.7)

improved for all psychiatric patients, regardless of the severity of the mental disorder.

In addition to inequalities in utilisation and quality of healthcare, there are many factors that contribute to poor physical health in people with a mental disorder, including health-related behaviours and medication side-effects. The increased morbidity and mortality seen in this population are largely due to a higher prevalence of modifiable risk factors, many of which are related to health-related behaviours. The use of alcohol and illicit drugs is more common among psychiatric patients. It is also known that psychiatric patients smoke to a larger extent,^{3 43} which could be an explanation for the increased mortality in IHD and cancer. Regardless of the kind of mental disorder, we found psychiatric patients to have an increased risk of dying in smoking-related cancer (data not shown). An unhealthy diet may also elevate the risk of various somatic diseases and thus increase the risk of death. The effects of these health-related behaviours have all been proposed as possible causative factors for excess mortality.⁴⁴ Also, medications used to treat severe mental disorder may increase the risk of diabetes and cardiovascular disease, as most mood stabilisers are associated with weight gain.^{12 45} However, these factors make careful monitoring of the physical health status of patients with mental disorders even more important.

CONCLUSIONS

This study shows a marked increase in excess mortality in mentally ill women and men. This was especially evident in individuals with a comorbid substance abuse disorder. The findings also suggest a lower quality of healthcare in the treatment of somatic disorders in psychiatric patients. Careful medical examination of psychiatric patients together with efforts to promote a healthier life

style may be of great importance in order to prevent, detect and treat somatic disease.

Author affiliations

¹Department of Public Health Sciences, Division of Social Medicine, Karolinska Institute, Stockholm, Sweden

²Department of Statistics, Monitoring and Evaluation, National Board of Health and Welfare, Stockholm, Sweden

³Department of Molecular Medicine and Surgery, Upper Gastrointestinal Research, Karolinska Institute, Stockholm, Sweden

⁴Department of Clinical Neuroscience, Division of Insurance Medicine, Karolinska Institute, Stockholm, Sweden

⁵Department of Public Health Sciences, Division of Public Health Epidemiology, Karolinska Institute, Stockholm, Sweden

⁶Department of Public Health and Caring Sciences, Uppsala University, Uppsala, Sweden

Acknowledgements We thank Dr Robert Bodén and Dr Herman Holm for excellent advice.

Contributors EB, GRW and RL originated the idea. EB analysed the data and wrote the manuscript draft. All authors contributed to the interpretation of the results and to the writing of the final manuscript.

Funding EMR is the recipient of the Assistant Professor grant from the Swedish Research Council (A0268301).

Competing interests None.

Ethics approval The ethics approval was provided by Central Ethical Review Board in Sweden (dnr 2011/295-31/4).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data available.

REFERENCES

1. D'Avanzo B, La Vecchia C, Negri E. Mortality in long-stay patients from psychiatric hospitals in Italy—results from the Qualyop Project. *Soc Psychiatry Psychiatr Epidemiol* 2003;38: 385–9.
2. Grigoletti L, Perini G, Rossi A, *et al*. Mortality and cause of death among psychiatric patients: a 20-year case-register study in an area with a community-based system of care. *Psychol Med* 2009;39:1875–84.
3. Harris EC, Barraclough B. Excess mortality of mental disorder. *Br J Psychiatry* 1998;173:11–53.

4. Kisely S, Smith M, Lawrence D, *et al.* Mortality in individuals who have had psychiatric treatment: population-based study in Nova Scotia. *Br J Psychiatry* 2005;187:552–8.
5. Meloni D, Miccinesi G, Bencini A, *et al.* Mortality among discharged psychiatric patients in Florence, Italy. *Psychiatr Serv* 2006;57:1474–81.
6. Miller BJ, Paschall CB 3rd, Svendsen DP. Mortality and medical comorbidity among patients with serious mental illness. *Psychiatr Serv* 2006;57:1482–7.
7. Brown S, Kim M, Mitchell C, *et al.* Twenty-five year mortality of a community cohort with schizophrenia. *Br J Psychiatry* 2010;196:116–21.
8. McGrath J, Saha S, Chant D, *et al.* Schizophrenia: a concise overview of incidence, prevalence, and mortality. *Epidemiol Rev* 2008;30:67–76.
9. Saha S, Chant D, McGrath J. A systematic review of mortality in schizophrenia: is the differential mortality gap worsening over time? *Arch Gen Psychiatry* 2007;64:1123–31.
10. Osby U, Correia N, Brandt L, *et al.* Mortality and causes of death in schizophrenia in Stockholm county, Sweden. *Schizophr Res* 2000;45:21–8.
11. Osby U, Brandt L, Correia N, *et al.* Excess mortality in bipolar and unipolar disorder in Sweden. *Arch Gen Psychiatry* 2001;58:844–50.
12. Roshanaei-Moghaddam B, Katon W. Premature mortality from general medical illnesses among persons with bipolar disorder: a review. *Psychiatr Serv* 2009;60:147–56.
13. Ringback Weitof G, Gullberg A, Rosén M, *et al.* Avoidable mortality among psychiatric patients. *Soc Psychiatry Psychiatr Epidemiol* 1998;33:430–7.
14. Rosen CS, Kuhn E, Greenbaum MA, *et al.* Substance abuse-related mortality among middle-aged male VA psychiatric patients. *Psychiatr Serv* 2008;59:290–6.
15. Carlborg A, Winnerbäck K, Jönsson EG, *et al.* Suicide in schizophrenia. *Expert Rev Neurother* 2010;10:1153–64.
16. Lester D. Sex differences in completed suicide by schizophrenic patients: a meta-analysis. *Suicide Life Threat Behav* 2006;36:50–6.
17. Limosin F, Loze JY, Philippe A, *et al.* Ten-year prospective follow-up study of the mortality by suicide in schizophrenic patients. *Schizophr Res* 2007;94:23–8.
18. Palmer BA, Pankratz VS, Bostwick JM. The lifetime risk of suicide in schizophrenia: a reexamination. *Arch Gen Psychiatry* 2005;62:247–53.
19. Laursen TM, Munk-Olsen T, Agerbo E, *et al.* Somatic hospital contacts, invasive cardiac procedures, and mortality from heart disease in patients with severe mental disorder. *Arch Gen Psychiatry* 2009;66:713–20.
20. Lawrence D, Kisely S, Pais J. The epidemiology of excess mortality in people with mental illness. *Can J Psychiatry* 2010;55:752–60.
21. Lawrence DM, Holman CD, Jablensky AV, *et al.* Death rate from ischaemic heart disease in Western Australian psychiatric patients 1980–1998. *Br J Psychiatry* 2003;182:31–6.
22. Howard LM, Barley EA, Davies E, *et al.* Cancer diagnosis in people with severe mental illness: practical and ethical issues. *Lancet Oncol* 2010;11:797–804.
23. Kisely S, Sadek J, MacKenzie A, *et al.* Excess cancer mortality in psychiatric patients. *Can J Psychiatry* 2008;53:753–61.
24. Druss BG, Bradford WD, Rosenheck RA, *et al.* Quality of medical care and excess mortality in older patients with mental disorders. *Arch Gen Psychiatry* 2001;58:565–72.
25. Levinson Miller C, Druss BG, Dombrowski EA, *et al.* Barriers to primary medical care among patients at a community mental health center. *Psychiatr Serv* 2003;54:1158–60.
26. Frayne SM, Halanych JH, Miller DR, *et al.* Disparities in diabetes care: impact of mental illness. *Arch Intern Med* 2005;165:2631–8.
27. Druss BG. Cardiovascular procedures in patients with mental disorders. *JAMA* 2000;283:3198–9.
28. Kisely S, Smith M, Lawrence D, *et al.* Inequitable access for mentally ill patients to some medically necessary procedures. *CMAJ* 2007;176:779–84.
29. National Board of Health and Welfare, Swedish association of Local Authorities and regions. *Quality and Efficiency Swedish Health Care*. Stockholm, Sweden: Regional Comparisons 2010, 2011.
30. Rutstein DD, Berenberg W, Chalmers TC, *et al.* Measuring the quality of medical care. *N Engl J Med* 1976;294:582–8.
31. Amadeo F, Barbui C, Perini G, *et al.* Avoidable mortality of psychiatric patients in an area with a community-based system of mental health care. *Acta Psychiatr Scand* 2007;115:320–5.
32. Sokal J, Messias E, Dickerson FB, *et al.* Comorbidity of medical illnesses among adults with serious mental illness who are receiving community psychiatric services. *J Nerv Ment Dis* 2004;192:421–7.
33. Ludvigsson JF, Otterblad-Olausson P, Pettersson BU, *et al.* The Swedish personal identity number: possibilities and pitfalls in healthcare and medical research. *Eur J Epidemiol* 2009;24:659–67.
34. Thornicroft G. *Shunned: Discrimination Against People with Mental Illness*. New York, United States: Oxford University Press, 2006.
35. National Board of Health and Welfare. *Causes of Death 2010*. Stockholm, Sweden: National Board of Health and Welfare, 2011.
36. Lawrence D, Holman CD, Jablensky AV, *et al.* Excess cancer mortality in Western Australian psychiatric patients due to higher case fatality rates. *Acta Psychiatr Scand* 2000;101:382–8.
37. Kunkel EJ, Woods CM, Rodgers C, *et al.* Consultations for 'maladaptive denial of illness' in patients with cancer: psychiatric disorders that result in noncompliance. *Psychooncology* 1997;6:139–49.
38. Gau SS, Cheng AT. Mental illness and accidental death. Case-control psychological autopsy study. *Br J Psychiatry* 2004;185:422–8.
39. Colhoun HM, Betteridge DJ. Treatment of lipid disorders in patients with diabetes. *Curr Treat Options Cardiovasc Med* 2006;8:37–45.
40. Lawrence D, Kisely S. Inequalities in healthcare provision for people with severe mental illness. *J Psychopharmacol* 2010;24(Suppl 4):61–8.
41. Kopp M, Fleischhacker WW, Stürz K, *et al.* Poor health behaviour and reduced quality of life of people treated with psychotropic drugs. *Hum Psychopharmacol* 2011;26:161–7.
42. Jones S, Howard L, Thornicroft G. 'Diagnostic overshadowing': worse physical health care for people with mental illness. *Acta Psychiatr Scand* 2008;118:169–71.
43. Brown S, Birtwistle J, Roe L, *et al.* The unhealthy lifestyle of people with schizophrenia. *Psychol Med* 1999;29:697–701.
44. Brown S, Inskip H, Barraclough B. Causes of the excess mortality of schizophrenia. *Br J Psychiatry* 2000;177:212–17.
45. De Hert M, Correll CU, Bobes J, *et al.* Physical illness in patients with severe mental disorders. I. Prevalence, impact of medications and disparities in health care. *World Psychiatry* 2011;10:52–77.

bmjopen-2011-000778 - Quality of medical care and excess mortality in psychiatric patients - a nationwide register-based study in Sweden
STROBE Statement—Checklist of items that should be included in reports of *cohort studies*

	Item No	Recommendation
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract Page 1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found Page 3
Introduction		
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported Page 4
Objectives	3	State specific objectives, including any prespecified hypotheses Page 5
Methods		
Study design	4	Present key elements of study design early in the paper Page 6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection Page 6
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up Page 6
		(b) For matched studies, give matching criteria and number of exposed and unexposed N/A
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable Page 6 - 8
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group Page 6 - 8
Bias	9	Describe any efforts to address potential sources of bias N/A
Study size	10	Explain how the study size was arrived at Page 6
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why N/A
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding Page 8 - 9
		(b) Describe any methods used to examine subgroups and interactions N/A
		(c) Explain how missing data were addressed N/A
		(d) If applicable, explain how loss to follow-up was addressed N/A

Results		
Participants	13*	<p>(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed</p> <p>Page 10 and page 20</p> <p>(b) Give reasons for non-participation at each stage</p> <p>N/A</p> <p>(c) Consider use of a flow diagram</p> <p>N/A</p>
Descriptive data	14*	<p>(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders</p> <p>Page 20</p> <p>(b) Indicate number of participants with missing data for each variable of interest</p> <p>N/A</p> <p>(c) Summarise follow-up time (eg, average and total amount)</p> <p>Page 20</p>
Outcome data	15*	<p>Report numbers of outcome events or summary measures over time</p> <p>Page 10 and page 20</p>
Main results	16	<p>(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included</p> <p>Page 10-11 and page 21-23</p> <p>(b) Report category boundaries when continuous variables were categorized</p> <p>N/A</p> <p>(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period</p> <p>N/A</p>
Other analyses	17	<p>Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses</p> <p>Page 11 and page 24</p>
Discussion		
Key results	18	<p>Summarise key results with reference to study objectives</p> <p>Page 2 and page 12 – 16</p>
Limitations	19	<p>Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias</p> <p>Page 2 and page 12</p>
Interpretation	20	<p>Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence</p> <p>Page 12 – 16</p>
Generalisability	21	<p>Discuss the generalisability (external validity) of the study results</p> <p>Page 15</p>
Other information		
Funding	22	<p>Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based</p> <p>Page 16</p>

*Give information separately for exposed and unexposed groups.