

## Correction

Kausto J, Solovieva S, Virta LJ, *et al.* Partial sick leave associated with disability pension: propensity score approach in register-based cohort study. *BMJ Open* 2012;2:e001752. Incorrect figures were published in table 5. The corrected table is published below.

*BMJ Open* 2012;2:e001752corr1. doi:10.1136/bmjopen-2012-001752corr1

**Table 5** Absolute and relative risk reduction (a risk difference between partial and full sick leave groups, ARR\* and RRR\*) of the use of partial and full disability pension with 95% CIs in the matched subsample

	Full disability pension				Partial disability pension					
	ARR (%)	95% CI	RRR (%)	95% CI	NNT	ARR (%)	95% CI	RRR (%)	95% CI	NNT
Total	6	3 to 9	41	24 to 55	16	-8	-10 to (-5)	-159	-264 to (-84)	-14
Men	10	4 to 16	47	20 to 65	10	-5	-10 to (-0)	-102	-302 to (-2)	-21
Women	4	0.5 to 7	32	5 to 52	26	-9	-12 to (-6)	-180	-315 to (-89)	-12
Mental disorders	8	3 to 13	42	17 to 59	12	-10	-14 to (-7)	-361	-795 to (-138)	-10
Musculoskeletal disorders	6	2 to 11	51	19 to 70	15	-7	-12 to (-2)	-94	-202 to (-24)	-15
Traumas and tumours	2	-4 to 8	17	-54 to 56	53	0	-8 to 2	-95	-409 to 25	-31

\*Negative values indicate an increase in risk.

\*NNT, number needed to treat to prevent full disability pension.