

# BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [editorial.bmjopen@bmj.com](mailto:editorial.bmjopen@bmj.com)

# BMJ Open

**The PTED study: design of a non-inferiority, randomised controlled trial to compare the effectiveness and cost-effectiveness of percutaneous transforaminal endoscopic discectomy (PTED) vs. open microdiscectomy for patients with a symptomatic lumbar disc herniation.**

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-018230
Article Type:	Protocol
Date Submitted by the Author:	21-Jun-2017
Complete List of Authors:	Seiger, Ankie; Vrije Universiteit Amsterdam, Health Sciences Gadjradj, Pravesh; Erasmusmc, Neurosurgery Harhangi, Sanjay; Erasmus MC, Neurosurgery van Susante, Job; Ziekenhuis Rijnstate, Orthopedics Peul, Wilco; Leiden University Medical Center, Neurosurgery Tulder, Maurits; University of Amsterdam, Health Sciences De Boer, Michiel; Vrije Universiteit Amsterdam, Health sciences Rubinstein, Sidney; Vrije Universiteit Amsterdam, Health Science
<b>Primary Subject Heading</b>:	Surgery
Secondary Subject Heading:	Health economics, Epidemiology, Neurology
Keywords:	HEALTH ECONOMICS, NEUROSURGERY, ORTHOPAEDIC & TRAUMA SURGERY

SCHOLARONE™  
Manuscripts

1  
2  
3 **The PTED study: design of a non-inferiority, randomised controlled trial to compare the effectiveness and**  
4 **cost-effectiveness of percutaneous transforaminal endoscopic discectomy (PTED) vs. open microdiscectomy**  
5 **for patients with a symptomatic lumbar disc herniation.**  
6  
7  
8

9 Seiger A<sup>1</sup>, Gadgradj P.S.<sup>2,3</sup>, Harhangi B.S.<sup>2</sup>, van Susante J.L.C.<sup>4</sup>, Peul W.C.<sup>3</sup>, van Tulder M.W.<sup>1</sup>, de Boer M.R.<sup>1</sup>,  
10 Rubinstein S.M.<sup>1</sup>  
11

12  
13 <sup>1</sup> Department of Health Sciences, Faculty of Science, Amsterdam Public Health research institute, Vrije Universiteit  
14 Amsterdam, The Netherlands  
15

16  
17 <sup>2</sup> Department of Neurosurgery, Erasmus MC: University Medical Centre Rotterdam, The Netherlands  
18

19  
20 <sup>3</sup> Department of Neurosurgery, Leiden University Medical Centre, LUMC, The Netherlands  
21

22  
23 <sup>4</sup> Department of Orthopedics, Rijnstate Hospital, The Netherlands  
24

25 **Corresponding author:**

26 Ankie Seiger, MSc

27 Vrije Universiteit Amsterdam, Faculty of Science, Department of Health Sciences

28 De Boelelaan 1085

29 1081 HV Amsterdam

30 a.seiger@vu.nl

31 +31 20 59 82745  
32  
33

34  
35 (Word count: 4729)  
36

37  
38 **Keywords:** Surgery, lumbar disc herniation, open microdiscectomy, percutaneous transforaminal endoscopic  
39 discectomy, effectiveness, economic evaluation.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## ABSTRACT

**INTRODUCTION:** Lumbosacral radicular syndrome is often caused by a disc herniation. The standard surgical technique to remove a disc herniation is open microdiscectomy. An alternative technique is Percutaneous Transforaminal Endoscopic Discectomy (PTED), which is less invasive. In The Netherlands, PTED is not currently considered standard care and therefore, not reimbursed within public health insurance. A pragmatic, multi-centre, non-inferiority, randomised controlled trial has been designed to determine the effectiveness and cost-effectiveness of PTED versus open microdiscectomy for the treatment of a lumbar disc herniation.

**METHOD AND ANALYSIS:** In total, 682 patients between 18-70 years of age with > 10 weeks of radiating pain or with > 6 weeks of excessive radiating pain are to be recruited from participating centres. Patients must have an indication for surgery based upon an MRI demonstrating compression of the nerve root from a lumbar disc herniation. Patients are to be randomised to PTED or open microdiscectomy. The primary outcome is self-reported leg pain measured by the 0-100 mm Visual Analogue Scale. Secondary outcomes include self-reported health and functional status; back pain; self-perceived recovery; and a physical examination. Outcomes will be measured the day following treatment and at 2, 4, and 6 weeks, and 3, 6, 9, 12 and 24 months. Physical examination will be performed at 6 weeks, and 3 and 12 months. An economic evaluation will be performed from a societal perspective and cost-questionnaires will be used (e.g. EQ-5D-5L). The data will be analysed longitudinally; the non-inferiority margin for the primary outcome is 5. Bootstrapping techniques will be used for the economic evaluation.

**ETHICS AND DISSEMINATION:** This study has received approval of the Medical Ethical Committee of the VU Medical Centre Amsterdam: NL50951.029.14. The results will be published in an international peer reviewed scientific journal.

**REGISTRATION:** This trial is registered at ClinicalTrials.gov: NCT02602093.

### Strengths and Limitations

- Large, multi-centre, pragmatic, randomised controlled trial
- Use of standardized and validated outcomes instruments
- Data are to be analysed longitudinally and multi-level
- Inclusion of an economic evaluation
- Potential performance bias due to lack of blinding of patients and providers

## INTRODUCTION

Lumbosacral radicular syndrome is a common health problem with a lifetime prevalence that varies from 12.2% to 43% and has a point prevalence ranging from 1.6% to 13.4%.<sup>1-6</sup> Lumbosacral radicular syndrome is often caused by a lumbar herniated disc<sup>7</sup> and is associated with a greater incidence of sickness benefit,<sup>8</sup> increased pain and disability, and poorer quality of life<sup>9</sup> than those with non-specific low-back pain. In cases of a disc herniation, lumbosacral radicular syndrome can be treated either conservatively or surgically.<sup>7</sup>

To remove the disc herniation, the standard surgical technique is open microdiscectomy. A more recently developed technique is percutaneous transforaminal endoscopic discectomy (PTED). In short, open microdiscectomy is performed under general anaesthesia and surgeons operate with a direct vision on the herniated disc, while PTED is conducted transforaminally and these patients undergo local anaesthesia and

1  
2  
3 surgeons operate through a working cannula with an indirect vision via an endoscope. Based upon the current  
4 literature, PTED is a safe method for the removal of a lumbar disc herniation.<sup>10</sup> Possible benefits of PTED versus  
5 open microdiscectomy are: 1) Decreased medical costs because patients are treated on an outpatient basis; 2) It  
6 is easier to remove intra- and extra-foraminal herniated discs; 3) There is less chance of scar formation; and 4)  
7 The technique is potentially more effective for obese patients. However, too few, large prospective studies have  
8 examined this in detail, therefore, the benefits may be speculated.<sup>11-13</sup>

9  
10  
11  
12 Despite that PTED is becoming more commonly used, there are still questions regarding its effect and the  
13 associated costs.<sup>11, 13, 14</sup> A recent systematic review<sup>11</sup> identified three randomised controlled trials (RCTs), which  
14 examined the effect of PTED compared to open microdiscectomy.<sup>15-17</sup> Their results suggest that there is low to  
15 very low quality evidence that PTED is no more effective than open microdiscectomy for self-reported back pain,  
16 leg pain, functional status, recovery, return-to-work, and satisfaction with surgery. Importantly, all three studies  
17 were of poor methodological quality and examined relatively few patients (i.e. ranging from 40 to 60  
18 individuals). A more recent study concluded that PTED shows similar results compared to open  
19 microdiscectomy.<sup>13</sup> However, this was a single-centre study; it was conducted by a surgeon with a keen interest  
20 in the results of PTED; it included patients over a long period of time (i.e. the study started in 2006 and was  
21 published in 2017) suggesting possible selection of patients included in the trial; and the inclusion criteria  
22 published in the original protocol were different from those in the final publication. Additionally, the economic  
23 evaluation of this study has not yet been published.<sup>18</sup> This makes it difficult to assess the cost-effectiveness.  
24 Therefore, discussion regarding the effectiveness and cost-effectiveness of PTED remains.

25  
26  
27  
28  
29  
30  
31 In The Netherlands, the effectiveness of PTED has been heatedly debated. According to the Dutch Health Care  
32 Institute, a new surgical technique must meet certain requirements in order to be reimbursed by the public  
33 health system. The Health Care Institute promotes the quality of Dutch Health Care and advises the Ministry of  
34 Health, Welfare and Sport on the content of the public health insurance. Based on a review,<sup>11</sup> the Health Care  
35 Institute claimed there is insufficient evidence for PTED to be included for reimbursement from the public health  
36 insurance package and as a result, patients are forced to pay the costs of treatment out-of-pocket. In order to  
37 deal with this issue and to answer the remaining questions about PTED, this large, pragmatic, methodologically  
38 rigorous multi-centre study has been designed.

39  
40  
41  
42  
43 This study is expected to have a major societal impact because it will determine if PTED should be included in  
44 the Dutch health insurance package. Furthermore, this study will provide more insight in PTED internationally,  
45 resulting in improved care for patients with a lumbar disc herniation. The primary hypothesis of this study is that  
46 PTED is not less effective and not less cost-effective compared to standard care (i.e. open microdiscectomy) for  
47 patients with symptomatic, lumbosacral radicular syndrome as a result of a lumbar disc herniation. Therefore, a  
48 non-inferiority design will be used.

## 52 **METHOD AND ANALYSIS**

### 54 **Study design**

55 A pragmatic, multi-centre non-inferiority randomised controlled trial (RCT) will be used. Following the baseline  
56 measurements, wherein clinical and socio-demographic measurements will be collected, patients are to be  
57  
58  
59  
60

1  
2  
3 randomised to one of two groups: the control group will receive standard open microdiscectomy and the  
4 intervention group will receive PTED. Patients will be followed for two years, but the primary analysis will be  
5 conducted on the one-year data.  
6  
7

8 Important protocol modifications will be registered at ClinicalTrials.gov and communicated to all relevant parties  
9 involved in this study (Medical Ethical Committee, ZonMw, included patients, participating surgeons, and  
10 members from the advisory board (listed in the acknowledgments)).  
11  
12

### 13 **Study population**

14 In total, 682 patients with a magnetic resonance imaging (MRI) confirmed lumbosacral radicular syndrome, due  
15 to a lumbar disc herniation are to be recruited. Patients will be recruited from five hospitals and one private  
16 health clinic located in Arnhem/Zevenaar, Leiderdorp, Tilburg and Rotterdam (The Netherlands). Each patient is  
17 required to sign a written informed consent prior to participation. If patient recruitment is slower than planned,  
18 (one or two) additional hospitals may be recruited.  
19

20 In order to be eligible to participate and in accordance with the Dutch Guideline Lumbosacral Radicular  
21 Syndrome,<sup>19</sup> a subject must meet all of the following conditions:  
22  
23

#### 24 **Inclusion criteria**

- 25 • 18-70 years of age;
- 26 • > 10 weeks of radiating pain with or without motor or sensory loss in the leg, or with > 6 weeks of  
27 excessive radiating pain and no tendency for any clinical improvement;
- 28 • Indication for surgery;
- 29 • MRI demonstrating lumbar disc herniation with nerve compression with or without concomitant  
30 spinal or lateral recess stenosis or sequestration;
- 31 • Sufficient knowledge of the Dutch language in order to complete forms and follow instructions  
32 independently.  
33

#### 34 **Exclusion criteria**

- 35 • Previous surgery on the same or adjacent disc level;
- 36 • Cauda equina syndrome;
- 37 • Spondylitic- or degenerative spondylolisthesis;
- 38 • Pregnancy;
- 39 • Severe comorbid medical or psychiatric disorder (American Society of Anaesthesiologists >2);
- 40 • Severe caudal or cranial sequestration;
- 41 • Contra-indication for surgery;
- 42 • Moving abroad at short notice.  
43

#### 44 **Study procedures**

45 Participating surgeons will screen all eligible patients with a lumbar disc herniation during the consultation  
46 (Table 1). If eligible for inclusion, patients will receive information relevant to the study by means of a letter.  
47 Dutch law requires that patients are given at least two days to consider participation. Following this initial  
48  
49  
50  
51  
52

1  
2  
3 screening, the patient is to be examined by a trained research nurse and the informed consent is obtained;  
4 baseline measurements are performed; and patients are randomised.  
5  
6

### 7 **Randomisation**

8 Patients will be randomised in a 1:1 ratio to PTED or open microdiscectomy. An experienced statistician  
9 prepares computer-generated, random number tables. Treatment allocation will be concealed. The key will be  
10 withheld from all participants and researchers involved in this study. Variable block sizes of 4, 6 and 8 are used  
11 and stratified by treatment centre. The random-number tables are to be entered into a computer system by an  
12 independent software company, and allocation will be performed by the computer system once baseline data  
13 and physical examination are obtained from an independent research nurse responsible for the treatment  
14 allocation.  
15  
16

### 17 **Blinding**

18 No attempt will be made to blind the patients. Blinding is considered impossible, because the procedures are  
19 fundamentally different. Furthermore, outcomes assessors cannot be blinded, given that the primary outcomes  
20 are all self-reported. The analysis will be performed blinded for treatment allocation.  
21  
22  
23  
24

### 25 **Treatment**

#### 26 *Intervention: Percutaneous transforaminal endoscopic discectomy (PTED)*

27 PTED is to be conducted as follows:<sup>20</sup> local anaesthesia is to be administered and consists of light sedation with  
28 dexmedetomidine or a combination of propofol and remifentanyl for the convenience of the patient. The  
29 amount of administered sedation should still allow the patient to respond to nerve root manipulation.  
30 Verification of the site is to be performed by an image intensifier using fluoroscopy (anteroposterior and lateral  
31 view) and is depending upon the patient's posture. An incision just above the dorsolateral side of the pelvis is  
32 conducted, where a needle is to be set from the incision to the superior articular process of the lower involved  
33 vertebrae of the herniated disc. Position will be checked again under fluoroscopy. After the needle has reached  
34 the superior articular process, a guide wire is to be inserted. Following that, a series of conical rods are to be  
35 introduced, subsequently a drill/reamer is to be introduced through the cannula and rods. After drilling through  
36 the superior articular process is conducted in order to enlarge the neuroforamen, the instruments are to be  
37 removed, but the guidewire is left in place and the endoscope with the working channels are to be introduced  
38 via an 8 mm cannula. The image intensifier ensures that the position of the cannula is maintained. Following  
39 removal of the disc herniation with a rongeur, the cannula and endoscope are removed. The patient is to be  
40 treated on an outpatient basis.  
41  
42  
43  
44  
45  
46  
47

#### 48 *Comparison: Open Microdiscectomy*

49 Open microdiscectomy is to be conducted as follows: general or spinal anaesthesia is to be administered.  
50 Verification is to be performed using a fluoroscopy and the patient is to be positioned prone or in the salaam  
51 position. Loupe or microscope magnification may be used according to the surgeon's preference. A paramedian  
52 incision is to be performed and the level is to be indicated. Following the identification of the lamina, the yellow  
53 ligament will be removed to identify the nerve root and disc herniation. Laminotomy as well as foraminotomy is  
54 to be performed, if necessary. The amount of degenerative disc material shall be removed at the discretion of  
55  
56  
57  
58  
59  
60



1  
2  
3 the attending surgeon. Post-operative policy will be followed and it is expected that the duration of recovery in  
4 the hospital may vary from 1-2 days, but the patient will be discharged as soon as medically responsible.  
5  
6

### 7 *Co-interventions*

8 Pain medication will be offered to patients, should this be necessary. In addition, use of co-interventions will be  
9 monitored by self-reported cost questionnaires, in which medication usage and any health care utilization is  
10 recorded throughout the follow-up period.  
11  
12

### 13 **Learning curve**

14 It will be necessary to train surgeons in the use of PTED because prior to the start of this study only two  
15 surgeons in The Netherlands were proficient in this technique. One of these surgeons is participating in this  
16 study (BSH). This experienced surgeon will provide the training to the other surgeons, all of who have more than  
17 ten years of surgical experience. The initial training will be first conducted on cadavers, and only once the  
18 surgeons are comfortable with the use of the procedure, will they perform this technique on patients under the  
19 tutelage of the PTED-experienced surgeon. It is expected that 50 patients per surgeon will be necessary to  
20 become proficient in PTED (defined as the 'learning curve'). Thus, 150 PTED patients will be registered as  
21 learning curve patients, and included in the study; however, they will not be included in the primary analysis.  
22 Additionally, competency in the use of PTED by the surgeons is to be evaluated using skill-based questions  
23 measured by a Likert-scale and the Objective Structured Assessment of Technical Skills (OSATS). These will be  
24 recorded and evaluated by both the teaching surgeon as well as the surgeons undergoing the training.<sup>21</sup>  
25  
26  
27  
28  
29  
30

### 31 **Prognostic factors**

32 The following potential prognostic factors are to be measured: 1) socio-demographic characteristics (e.g. age,  
33 gender); 2) characteristics of the complaint (e.g. duration and severity); 3) baseline pain and functional disability;  
34 4) lifestyle factors (e.g. smoking and alcohol use); 5) psychological factors (e.g. expectations of recovery,  
35 emotional well-being; 6) psychopathology as measured with the four dimensional symptom questionnaire  
36 (4DSQ; dimensions: distress, depression, anxiety and somatization);<sup>22</sup> 7) work-related factors (e.g. physical  
37 workload, job satisfaction); and 8) previously received treatment due to the same episode of back complaints  
38 (e.g. medication and physiotherapy).  
39  
40  
41  
42

### 43 **Outcome measurements**

44 The outcomes are to be measured by validated self-reported questionnaires and by physical examination. Data  
45 are to be collected prior to randomisation (baseline), the day following surgery, at 2, 4, and 6 weeks, and at 3, 6,  
46 9, 12, and 24 months following surgery (Table 1).  
47

48 All questionnaires will be sent automatically by e-mail with a personal link to the digital questionnaire. If  
49 necessary, a reminder will be sent after 3 days; after six days the research nurse will call the patient with the  
50 request to fill in the questionnaire. Patients who deviate from the original protocol will be registered and will be  
51 asked to continue filling in the self-reported questionnaires.  
52  
53  
54

### 55 **Primary outcomes**

56  
57  
58  
59  
60



1  
2  
3 The primary outcome, leg pain, is to be measured by the Visual Analogue Scale (VAS; scale 0-100 mm). This  
4 outcome measure has been identified in a systematic review to be one of the most commonly measured  
5 outcomes, and is specific and responsive to change in a population undergoing lumbar spine surgery.<sup>23</sup>  
6  
7

#### 8 Secondary outcomes

9  
10 *Functional status:* will be measured with the Oswestry Disability Index (ODI). The ODI<sup>24</sup> is one of the principal  
11 condition-specific outcome measures used in the management of spinal disorders. The current version of the  
12 ODI (2.1a) is to be used.<sup>25</sup> The ODI has been extensively tested and showed good psychometric properties.<sup>26</sup>  
13  
14

15 *Low-back pain:* will be measured with the VAS (scale ranging from 0mm (no pain) to 100mm (worst imaginable  
16 pain)).  
17  
18

19 *Generic quality of life:* will be measured with the Dutch version of the short form SF36. The SF-36 questionnaire  
20 has been validated and found reliable for low back pain.<sup>27</sup> The questions are divided into eight domains: 1)  
21 physical functioning, 2) physical role limitations, 3) emotional role limitations, 4) social functioning, 5) physical  
22 pain, 6) general mental health, 7) vitality, and 8) general health perception. Per domain the scores of the items  
23 are added up and transformed into a scale of 0 to 100. A higher score reflects a better health condition. In  
24 addition, these eight domains can be summarized in a physical and psychological main domain.  
25  
26  
27

28 *Self-perceived recovery of the patient:* will be measured with a seven-point Likert-scale. The score on this scale  
29 varies from 'completely recovered' to 'worse than ever'. We will dichotomize the outcome with 'completely  
30 recovered', 'moderately recovered' and 'a bit recovered' as 'recovered' and the other four categories as 'not  
31 recovered'.  
32  
33  
34

35 *Patient satisfaction:* will be measured using the Likert-scale, Body Image and the Cosmesis scale.<sup>28, 29</sup> Body  
36 satisfaction will be measured using a four-point Likert-scale (ranging from 'not at all', 'a little', 'quite', to 'yes,  
37 very much'). Satisfaction change of complaints and satisfaction treatment will be measured using a seven-point  
38 Likert-scale (ranging from 'completely satisfied with current symptoms' to 'completely dissatisfied with current  
39 symptoms'). The scales will be completed by the patients prior to and following surgery. Scar satisfaction will  
40 also be measured using the seven-point Likert-scale and with a 1-10 numeric rating scale (ranging from 1 = 'as  
41 ugly as conceivable' to 10 = 'almost no scar perceived').  
42  
43  
44

45 *Physical examination:* will be performed at 6 weeks, 3 and 12 months following surgery. This will include: scar  
46 size; patellar and achilles tendon reflexes; straight leg raising test; cross straight leg raising test; finger-floor  
47 distance; strength measurement of the quadriceps using the Medical Research Council (MRC); sensibility  
48 dermatomes L1-S1, abdominal muscle strength; and patients' weight. The patellar and tendon reflexes are to be  
49 measured in a sitting upright position with both feet dangling above the ground. Tendon reflexes are tapped up  
50 to a maximum of two times with the reflex hammer. Reflexes are distinguished into absent, reduced, normal,  
51 increased, and clonus reflexes. The straight leg raising test and cross straight leg raising test are both measured  
52 as negative when no shooting leg pain is perceived, and positive when shooting pain is perceived. Finger-floor  
53 distance is the distance between the longest finger and the floor when the patients perform a forward bent with  
54 the knees extended. Muscle strength of the quadriceps is measured from a sitting position. Patients will be  
55  
56  
57  
58  
59  
60

asked to extent their knee while the research nurse exerts counter-pressure just above the ankle. Muscle strengths are rated on the Dutch version of the MRC, ranging from 0= no contraction to 5 = normal muscle strength). For the sensibility the research nurse checks every dermatome area (L1-S1) by touching the patient with a sharp and blunt object. Patients indicate with their eyes closed when sensation is felt. Sensibility varies from decreased, normal or increased sensibility compared to the other leg. Abdominal strength is measured by counting the maximal number of abdominal crunches from the supine position. Patients are asked to reach the hands towards the bended knees and to lift the scapulae from the surface. At any time, the lumbar spine will be supported by the underlying surface to minimize the range of motion of the lumbar spine. Without an increase of pain the maximum is set at a cut-off point of 26 crunches.

Screening and operation case record forms are to be completed by the surgeons, while discharge forms, physical examination and baseline intake forms are to be completed by a trained research nurse.

Table 1. Flowchart visits and case report forms

Visits and Case report forms	Intake + baseline	Surgery	1-2 days following treatment	2 weeks following treatment	4 weeks following treatment	6 weeks following treatment	3 months following treatment	6 months following treatment	9 months following treatment	12 months following treatment	24 months following treatment
Surgeon visit	X					X*					
Informed consent	X										
Research nurse visit	X					X	X			X	
Randomisation	X										
Surgery		X									
Discharge		X**	X**								
Four-Dimensional Symptom Questionnaire (4DSQ)	X										
Oswestry Disability Index (ODI)	X		X	X	X	X	X	X	X	X	X
Cost questionnaires	X			X	X	X	X	X	X	X	
EuroQol (EQ-5D-5L)	X		X	X	X	X	X	X	X	X	X
VAS leg pain, VAS Back pain	X		X	X	X	X	X	X	X	X	X
Quality of Life VAS	X		X	X	X	X	X	X	X	X	X
Patient self-perceived recovery and satisfaction			X***	X	X	X	X	X	X	X	X
Short Form 36 (SF36)	X			X	X	X	X	X	X	X	X
Physical examination	X					X	X			X	
Revisit and complications											With occurrence

\* 6 weeks visit may be performed also by the research nurse depending on the normal protocol hospital.

\*\* Discharge form will be filled in depending on discharge moment.

\*\*\* Only self-perceived recovery is measured not self-perceived satisfaction.

### Complications, operative morbidity, and re-operations

1  
2  
3 Immediately following surgery and discharge, the surgeon and research nurse will perform a systematic  
4 assessment of complications (including urinary tract infection, secondary bleeding, and progressive neurological  
5 deficit). In addition, surgeons will record any perioperative complications like cerebrospinal fluid leakage, nerve  
6 root damage, and if the surgery was initiated at the wrong disc level. Re-operation at the initial site is to be  
7 considered a poor outcome. Re-operation in both groups will be recorded. Perioperative morbidity will be  
8 assessed with operation time, perioperative blood loss, hospital stay and re-operative rate as related to the  
9 primary condition (lumbar disc herniation).  
10  
11  
12

### 13 **Sample size calculation**

14 The mean difference and standard deviation (SD) for the VAS (leg pain) used in the sample size calculation was:  
15 mean 5; SD 14.9.<sup>30</sup> The margin of non-inferiority was set at 5, (one-sided) alpha at 0.05 and beta at 0.10 (power  
16 0.9). We estimated that in total 306 patients are needed to demonstrate non-inferiority on the primary  
17 outcome. Accounting for 20% attrition, the aim is to recruit 382 patients. As the Ministry of Health in the  
18 Netherlands has stipulated that PTED will only be reimbursed if patients participate in the randomised trial, an  
19 extra 300 patients will be necessary for the inclusion of 150 patients in the PTED learning curve. Consequently, a  
20 total of 682 patients will be recruited for this study. Patients are likely to participate in this study, because PTED  
21 will only be reimbursed by Dutch health care insurance for participants in this study. Therefore, reaching the  
22 target sample size is not likely to become a problem.  
23  
24  
25  
26  
27

### 28 **Data analysis**

29 Data analysis will be conducted by a researcher or statistician blinded for treatment allocation after follow-up is  
30 finished. No interim analysis will be performed.  
31  
32

33 All data handling (entry, coding, storage and analysis) is confidential and complies with the Dutch Personal Data  
34 Protection Act. The anonymous data are stored in a central warehouse for at least 15 years.  
35  
36

### 37 **Effect analysis**

38 Characteristics of the patients will be presented using descriptive statistics (mean (SD), median (range) or  
39 proportion) to assess if balanced groups were obtained after randomisation. The non-inferiority margins are set  
40 and listed in table 2.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 2. Non-inferiority margins

Outcome measurements	Expected differences	Non-inferiority margin
VAS leg pain (0-100 scale)	<5 **	5
ODI (0-100 scale)	<5 *	5
VAS low back pain (0-100 scale)	<5 *	5
SF36 (0-100)	<5*	5
Self-perceived recovery (% 1 and 2 on the 7-point Likert-scale)	<10%***	5
Patient satisfaction (% 4 on the 4-point Likert-scale)	<5%	5
Patient satisfaction (% 1 and 2 on the 7-points Likert-scale)	<5%	5
Scar satisfaction (1-10 scale)	<1	0.5
Patellar reflex (% normal reflexes)	<5%	5
Achilles reflex (% normal reflexes)	<5%	5
Straight leg raising test (% negative tests)	<5%	5
Cross straight leg raising (% negative tests)	<5%	5
Finger floor distance (cm)	<5	5
Muscle strength quadriceps (% normal muscular strength)	5%	5
Sensibility dermatomes L1-S1 (% normal sensibility)	5%	5
EQ-5D-5L	<0.05****	0.05
Costs (healthcare perspective)	<\$500,- ***	250
Costs (societal perspective)	<\$1500,- ***	500

\* Obtained from the literature <sup>13,30</sup>

\*\* Obtained from the literature <sup>13</sup>

\*\*\* Obtained from the literature <sup>30</sup>

\*\*\*\* Obtained from the literature <sup>31</sup>

The primary data analysis will examine the effects of PTED for leg pain for those patients not in the learning curve, and shall be conducted according to the intention-to-treat principle. If necessary, missing items will be imputed using multiple imputation techniques. Linear and generalized multi-level analyses will be used, accounting for dependency of measurements over time within patients and patients nested within the surgeons, thus representing a 3-level model: time, patient and surgeon in that order. The data are to be examined longitudinally and the primary analysis will be aimed at average differences in effectiveness between the two treatment modalities. We will also include treatment \* time interactions to explore whether these effects are different over time. In addition to the crude analyses, all analyses will be adjusted for potential confounders, such as age, gender, nature and severity of the presenting complaint. In a secondary analysis, a per-protocol analysis shall be conducted. The secondary continuous outcomes, such as low-back pain, functional status, will be analysed similar to the primary data analysis; however, recovery and some of the physical performance measures (Table 2.) are to be treated as a dichotomous variable and will be analysed in logistic regression analyses.

Complications will be summarized for the time period of the study, but also presented for those complications encountered before and after 6 weeks.

#### *Sensitivity analyses effect*

Sensitivity analysis will be conducted in those patients with a) a paramedian/median disc herniation and b) a foraminal/extra-foraminal disc herniation. The latter is much less common, representing for approximately 7% to 12% of all lumbar disc herniations.<sup>32</sup> The goal of this analysis is to test the robustness of the data to changes

1  
2  
3 in underlying assumptions regarding the type or location of the hernia, and in this particular case, to determine  
4 whether the same effect is found for the subgroup with paramedian/median lumbar disc herniation as for the  
5 entire group. Finally, a sensitivity analysis will be conducted for all patients, including those in the 'learning  
6 curve' in order to determine if these outcomes are different than the primary analyses.  
7  
8

9  
10 Results from all analyses will be expressed as mean effect estimate with 95% confidence intervals and these  
11 estimates will be subsequently compared to the margin of non-inferiority in order to make inferences about the  
12 non-inferiority of the intervention, PTED.  
13

### 14 **Economic evaluation**

15 Both cost-effectiveness and cost-utility analysis will be conducted from a societal perspective alongside the RCT.  
16 We will measure, value and analyse total costs of all patients and relate the difference in costs to the difference  
17 in effects between the two groups.  
18  
19

20  
21 Direct costs include costs of the interventions, hospitalisation after surgery, medication and other health care  
22 utilization. Patient costs and cost of productivity loss, absenteeism and presenteeism, will also be included.  
23 Health care utilization, patients cost and productivity loss will be measured using self-completed cost  
24 questionnaires. The cost of the interventions will be estimated using a bottom-up approach (micro-costing) and  
25 hospitalisation will be registered using case record forms. The Dutch tariff of the EQ-5D-5L<sup>33,34</sup> will be used to  
26 calculate the quality-adjusted life years (QALYs). The EuroQol measures five dimensions: mobility, self-care, daily  
27 activities, pain/discomfort, and anxiety/depression. Each dimension consists of one item, while five levels are  
28 distinguished ('no', 'slight', 'moderate', 'severe problems', 'unable to do').  
29  
30  
31  
32

33 Costs resulting from productivity loss are to be estimated using the friction cost method, which assumes that  
34 sick workers are replaced after a period of time (12 weeks).<sup>35</sup> Mean productivity costs per working hour are to  
35 be adjusted for age and gender and used to estimate the cost of absenteeism. Health care utilization is to be  
36 valued according to the guidelines published in the updated handbook for economic evaluation in The  
37 Netherlands.<sup>35</sup> Medication is to be valued using prices from the Royal Dutch Society for Pharmacy.<sup>36</sup>  
38  
39  
40

### 41 *Cost-effectiveness analysis*

42 Total costs will be related to the primary effect measure, leg pain. A cost-utility analysis will be performed with  
43 QALYs. From the EQ-5D-5L utilities will be obtained and QALYs will be calculated using linear interpolation  
44 between measurement points. The primary analysis will be conducted according to intention-to-treat. Missing  
45 data will be imputed using multiple imputation by chained equations.<sup>37</sup> Incremental Cost Effectiveness Ratios  
46 (ICERs) will be calculated by dividing the difference in costs by the difference in effects. We will perform a cost-  
47 effectiveness analysis with leg pain as outcome and a cost-utility analysis with QALYs as outcome. In order to  
48 account for the possible clustering of data, analyses will be performed using linear multilevel analyses.<sup>38</sup>  
49 Accounting for the possible clustering of data (e.g. at the hospital and surgeon level) is very important, as most  
50 economic evaluations fail to do so, whereas ignoring the possible clustering of data might lead to inaccurate  
51 levels of uncertainty and inaccurate point estimates.<sup>38</sup> Bias corrected and accelerated bootstrapping with 5,000  
52 replications will be performed in order to estimate 95% confidence intervals around cost differences and the  
53 uncertainty surrounding the ICERs. Uncertainty will be shown in cost- effectiveness planes and cost-  
54  
55  
56  
57  
58  
59  
60

effectiveness acceptability curves, and sensitivity analyses will be performed to test the robustness of the study results.<sup>39-41</sup>

### *Sensitivity analysis cost-effectiveness*

Sensitivity analyses will be conducted for the most important cost drivers in order to determine the robustness of the findings. In addition, the main analyses are to be repeated using only complete cases (i.e. complete clinical outcome data and complete cost data). Lastly, the impact of the Human Capital Approach will be compared to the friction cost method approach. The Human Capital Approach evaluates the total costs of productivity loss without considering the possibility of replacing the sick worker.

## **ETHICS AND DISSEMINATION**

This study has received approval of the Medical Ethical Committee of the VU Medical Centre Amsterdam and confirmation can be supplied upon request [corresponding number: NL50951.029.14; November 5, 2015]. Serious adverse events (SAE) and adverse events will be registered; SAE will be reported within 24 hours (see section 'complications, operative morbidity, and re-operations'). The sponsor (also) has an insurance, which is in accordance with the legal requirements in the Netherlands. The insurance applies to the damage that becomes apparent during the study or within four years after the end of the study. This study will be monitored according to a detailed monitoring plan adapted to the risk classification of the Dutch University Federation guidelines. Based on this guideline, the risk classification of this study is regarded negligible. Considerations in this assessment are that this is an investigator-initiated trial, not with vulnerable patients, and while side effects are known, such as nerve root damage, severe adverse events are extremely rare. Audits may be required by and will be granted to Medical Ethical Committee and regulatory authority inspections. Patients' permission for these audits is obtained with informed consent.

The final trial results will be communicated to participants, healthcare professionals, professional organisations and relevant guideline committees in the Netherlands. We will publish the results in an international peer reviewed open access scientific journal. There are no publication restrictions.

## **DISCUSSION**

This large, multi-centre, pragmatic study will be conducted to resolve the discussion regarding the effects and costs of PTED compared to open microdiscectomy for patients with lumbosacral radicular syndrome caused by a lumbar disc herniation. At the moment, in the Netherlands PTED does not comply with standards of practice, and therefore, is not included in the current Dutch public healthcare package. The Dutch Ministry of Health classified PTED as an important technique to examine, but placed conditions on reimbursement of the intervention. Namely, patients must be included in the RCT in order to receive reimbursement. This presented a unique challenge for us because this means that all patients are to be included from the beginning while the surgeons who are not experienced in PTED are still undergoing the training, hence, the reason for the learning curve and the reason why these participants will be included in the study. Additionally, in order to prevent discussion regarding the effects of PTED following this study, a document was signed by all participating parties (e.g. professional surgical organisations, insurance companies, Dutch healthcare institute) to agree upon the study design and the criteria for in- or exclusion of PTED from the Dutch public healthcare package.



Since the trial was published in a trial registry (clinicaltrials.gov; November 2013), the protocol has been modified, namely a physical examination has been added in order to obtain more objective information regarding the physical rehabilitation, and a numeric rating scale (NRS) has been added in order to measure back, leg pain and quality of life. The reason for the latter is, the VAS may be completed by participants using different digital apparatuses (i.e. PC, tablet or mobile phone), with the result that the lengths of the VAS scale may vary. The validity and reliability is, therefore, uncertain. Based upon the literature, it would appear that the NRS and VAS demonstrate comparable values for pain following surgery;<sup>42</sup> however, the aforementioned issue, namely the use of the VAS on different digital apparatuses has not previously been examined. In order to determine whether this has bearing on the outcomes, a sensitivity analysis shall be conducted, and we will examine the correlation between the VAS and NRS.

The trial is an ongoing study and runs from February 2016 to February 2020.

#### Acknowledging:

Dr. C.L.A.M. Vleggeert-Lankamp, Drs. P.R.A.M. Depauw, Drs. P.J. Schutte, Drs. A.W.J. Vreeling; the hospitals, Albert Schweitzer Ziekenhuis, Alrijne Ziekenhuis, Elisabeth-Tweesteden Ziekenhuis, Franciscus Gasthuis en Vlietland, Park Medisch Centrum, Rijnstate Ziekenhuis; and the Advisory Committee: Dr. J.L.P. Kuijpers, Zorgverzekeraars Nederland (ZN), patient organisation 'Nederlandse Vereniging van Rugpatiënten de "Wervelkolom"', Dutch Spine Society, Nederlandse Orthopaedische Vereniging.

#### REFERENCES

1. Heliövaara, M, Impivaara, O, Sievers, K et al. Lumbar disc syndrome in Finland. *J Epidemiol Community Health* 1987;41:251-8.
2. Konstantinou, K and Dunn, KM. Sciatica: review of epidemiological studies and prevalence estimates. *Spine (Phila Pa 1976)* 2008;33:2464-72.
3. Nielen, MMJ. Incidentie en prevalentie van gezondheidsproblemen in de Nederlandse huisartsenpraktijk in 2014. *NIVEL Zorgregistraties eerste lijn* 2015. ICPC L86. <https://www.nivel.nl/node/4309> (date accessed June 2017).
4. Spijker-Huiges, A, Groenhof, F, Winters, JC et al. Radiating low back pain in general practice: incidence, prevalence, diagnosis, and long-term clinical course of illness. *Scand J Prim Health Care* 2015;33:27-32.
5. Oosterhuis, T, Smaardijk, V, Kuijer, P et al. Actualisatie wetenschappelijke kennis en inzichten over het lumbosacraal radiculair syndroom. *Uitvoeringsinstituut werknemersverzekeringen UWV* 2016; 16-06. <https://www.uwv.nl/overuwv/Images/Actualisatie-lumbosacraal-radiculair-syndroom.pdf> (date assessed June 2017).
6. Schaafstra, A, Spinnewijn, WEM, Bons, S et al. NHG-Standaard Lumbosacraal radiculair syndroom (tweede herziening). *Huisarts & Wetenschap* 2015;58:308-20.
7. Koes, BW, van Tulder, MW, and Peul, WC. Diagnosis and treatment of sciatica. *BMJ* 2007;334:1313-7.
8. Kristman, VL, Hartvigsen, J, Leboeuf-Yde, C et al. Does Radiating Spinal Pain Determine Future Work Disability? A Retrospective Cohort Study of 22,952 Danish Twins. *Spine* 2012;37:1003-1013.
9. Konstantinou, K, Hider, SL, Jordan, JL et al. The impact of low back-related leg pain on outcomes as compared with low back pain alone: a systematic review of the literature. *Clin J Pain* 2013;29:644-54.
10. Gadraj, PS, van Tulder, MW, Dirven, CM et al. Clinical outcomes after percutaneous transforaminal endoscopic discectomy for lumbar disc herniation: a prospective case series. *Neurosurg Focus* 2016;40:E3.



11. Kamper, SJ, Ostelo, RW, Rubinstein, SM et al. Minimally invasive surgery for lumbar disc herniation: a systematic review and meta-analysis. *Eur Spine J*2014;23:1021-43.
12. Gu, YT, Cui, Z, Shao, HW et al. Percutaneous transforaminal endoscopic surgery (PTES) for symptomatic lumbar disc herniation: a surgical technique, outcome, and complications in 209 consecutive cases. *J Orthop Surg Res*2017;12:25.
13. Gibson, JNA, Subramanian, AS, and Scott, CEH. A randomised controlled trial of transforaminal endoscopic discectomy vs microdiscectomy. *European Spine Journal*2017;26:847-856.
14. Pan, Z, Ha, Y, Yi, S et al. Efficacy of Transforaminal Endoscopic Spine System (TESSYS) Technique in Treating Lumbar Disc Herniation. *Med Sci Monit*2016;22:530-9.
15. Hermantin, FU, Peters, T, Quartararo, L et al. A prospective, randomized study comparing the results of open discectomy with those of video-assisted arthroscopic microdiscectomy. *J Bone Joint Surg Am*1999;81:958-65.
16. Mayer, HM and Brock, M. Percutaneous endoscopic discectomy: surgical technique and preliminary results compared to microsurgical discectomy. *J Neurosurg*1993;78:216-25.
17. Krappel, FA, Schmitz, R, Bauer, E et al. Open or endoscopic nucleotomy? Results of a prospective, controlled clinical trial with independent follow-up, MRI and special reference to cost-effectiveness. *Orthopadische Prax*2001;37:164-169.
18. Scott, CEH and Gibson, AJN. A cost utility comparison of transforaminal endoscopic discectomy and microdiscectomy. *The Spine Journal*2016;16:S47.
19. NVN and CBO. Richtlijn lumbosacraal radiculair syndroom. *Nederlandse Vereniging voor Neurologie en Kwaliteitsinstituut voor de Gezondheidszorg CBO*2008.  
[http://www.anesthesiologie.nl/uploads/openbaar/RL\\_Lumbosacraal\\_Radiculair\\_Syndroom\\_2008.pdf](http://www.anesthesiologie.nl/uploads/openbaar/RL_Lumbosacraal_Radiculair_Syndroom_2008.pdf) (date assessed June 2017).
20. Gadjradj, PS and Harhangi, BS. Percutaneous Transforaminal Endoscopic Discectomy for Lumbar Disk Herniation. *Clin Spine Surg*2016;29:368-371.
21. Hiemstra, E and F.W., J. Richtlijn OSATS. *Tijdschrift voor Medisch Onderwijs*2010;29:135-141.
22. Terluin, B, van Marwijk, HW, Ader, HJ et al. The Four-Dimensional Symptom Questionnaire (4DSQ): a validation study of a multidimensional self-report questionnaire to assess distress, depression, anxiety and somatization. *BMC Psychiatry*2006;6:34.
23. DeVine, J, Norvell, DC, Ecker, E et al. Evaluating the correlation and responsiveness of patient-reported pain with function and quality-of-life outcomes after spine surgery. *Spine (Phila Pa 1976)*2011;36:S69-74.
24. Fairbank, JC, Couper, J, Davies, JB et al. The Oswestry low back pain disability questionnaire. *Physiotherapy*1980;66:271-3.
25. van Hooff, ML, Spruit, M, Fairbank, JC et al. The Oswestry Disability Index (version 2.1a): validation of a Dutch language version. *Spine (Phila Pa 1976)*2015;40:E83-90.
26. Smeets, R, Koke, A, Lin, CW et al. Measures of function in low back pain/disorders: Low Back Pain Rating Scale (LBPRS), Oswestry Disability Index (ODI), Progressive Isoinertial Lifting Evaluation (PILE), Quebec Back Pain Disability Scale (QBPDS), and Roland-Morris Disability Questionnaire (RDQ). *Arthritis Care Res (Hoboken)*2011;63 Suppl 11:S158-73.
27. Chapman, JR, Norvell, DC, Hermsmeyer, JT et al. Evaluating common outcomes for measuring treatment success for chronic low back pain. *Spine (Phila Pa 1976)*2011;36:S54-68.
28. Lind, MY, Hop, WC, Weimar, W et al. Body image after laparoscopic or open donor nephrectomy. *Surg Endosc*2004;18:1276-9.
29. Dunker, MS, Stiggelbout, AM, van Hogezaand, RA et al. Cosmesis and body image after laparoscopic-assisted and open ileocolic resection for Crohn's disease. *Surg Endosc*1998;12:1334-40.
30. Arts, MP, Brand, R, van den Akker, ME et al. Tubular discectomy vs conventional microdiscectomy for sciatica: a randomized controlled trial. *JAMA*2009;302:149-58.

- 1
- 2
- 3
- 4 31. van den Akker, ME, Arts, MP, van den Hout, WB et al. Tubular discectomy vs conventional
- 5 microdiscectomy for the treatment of lumbar disk-related sciatica: cost utility analysis alongside a
- 6 double-blind randomized controlled trial. *Neurosurgery*2011;69:829-35; discussion 835-6.
- 7 32. Lee, JH and Lee, SH. Clinical and Radiological Characteristics of Lumbosacral Lateral Disc Herniation in
- 8 Comparison With Those of Medial Disc Herniation. *Medicine*2016;95:
- 9 33. Herdman, M, Gudex, C, Lloyd, A et al. Development and preliminary testing of the new five-level version
- 10 of EQ-5D (EQ-5D-5L). *Qual Life Res*2011;20:1727-36.
- 11 34. Versteegh, MM, Vermeulen, KM, Evers, SMAA et al. Dutch Tariff for the Five-Level Version of EQ-5D.
- 12 *Value in Health*2016;19:343-352.
- 13 35. Hakkaart-Van Roijen, L, van der Linden, N, Bouwmans, C et al. Kostenhandleiding: Methodologie van
- 14 kostenonderzoek en referentieprijzen voor economische evaluaties in de gezondheidszorg [Manual for
- 15 economic evaluations. Methods and standard cost prices for economic evaluations in health care].
- 16 *Diemen, The Netherlands: Health Care Institute (ZiN)*2016. [https://www.zorginstituutnederland.nl/over-](https://www.zorginstituutnederland.nl/over-ons/publicaties/publicatie/2016/02/29/richtlijn-voor-het-uitvoeren-van-economische-evaluaties-in-de-gezondheidszorg)
- 17 [ons/publicaties/publicatie/2016/02/29/richtlijn-voor-het-uitvoeren-van-economische-evaluaties-in-de-](https://www.zorginstituutnederland.nl/over-ons/publicaties/publicatie/2016/02/29/richtlijn-voor-het-uitvoeren-van-economische-evaluaties-in-de-gezondheidszorg)
- 18 [gezondheidszorg](https://www.zorginstituutnederland.nl/over-ons/publicaties/publicatie/2016/02/29/richtlijn-voor-het-uitvoeren-van-economische-evaluaties-in-de-gezondheidszorg) (date assessed June 2017).
- 19 36. Z-Index. G-Standaard. *The Hague, The Netherlands: Z-Index BV*.2009. [https://www.z-index.nl/g-](https://www.z-index.nl/g-standaard)
- 20 [standaard](https://www.z-index.nl/g-standaard) (date assessed January 2017).
- 21 37. White, IR, Royston, P, and Wood, AM. Multiple imputation using chained equations: Issues and guidance
- 22 for practice. *Statistics in Medicine*2011;30:377-399.
- 23 38. Gomes, M, Grieve, R, Nixon, R et al. Methods for Covariate Adjustment in Cost-Effectiveness Analysis
- 24 That Use Cluster Randomised Trials. *Health Economics*2012;21:1101-1118.
- 25 39. Drummond, M and Drummond, M, *Methods for the economic evaluation of health care programmes*.
- 26 3rd ed. Oxford medical publications. 2005, Oxford ; New York: Oxford University Press. 379 p.
- 27 40. Black, WC. The CE plane: a graphic representation of cost-effectiveness. *Med Decis Making*1990;10:212-
- 28 4. 41. Fenwick, E, O'Brien, BJ, and Briggs, A. Cost-effectiveness acceptability curves--facts, fallacies and
- 29 frequently asked questions. *Health Econ*2004;13:405-15.
- 30 42. Breivik, H, Borchgrevink, PC, Allen, SM et al. Assessment of pain. *Br J Anaesth*2008;101:17-24.
- 31
- 32
- 33
- 34
- 35
- 36
- 37

## FOOTNOTES

### Contributorship statement:

SH, JvS, WP, MvT, SR contributed to writing the proposal and acquiring the grant from The Netherlands Organisation for Health and Care Research (ZonMw). AS prepared the first draft of the study protocol. All authors (AS, PG, SH, JvS, WP, MvT, MdB, SR) contributed to the final design of this study protocol, assisted with drafting the manuscript, and approved the final version of the manuscript. MdB developed the statistical analysis plan.

### Funding statement:

This study was funded by ZonMw, The Netherlands Organisation for Health Research and Development [projectnumber 837004013]. The funder did not play any role in the study design. This research received no specific grant from any funding agency in the public, commercial or not-profit sectors.

### Competing interest statement:

1  
2  
3 The professional judgment concerning the primary interest (e.g. validity of the research) will not be influenced  
4 by a secondary interest. All authors have read and understood the BMJ policy on declaration of interests. We all  
5 declare that we have no competing interests other than receiving grants. For this particular study we had  
6 financial support from The Netherlands Organisation for Health Research and Development. We do not have  
7 relationships with any organisations that might have an interest in the submitted work. All authors have  
8 completed the ICJE uniform at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf).  
9  
10  
11

12 **Provenance and peer review:** Not commissioned; externally peer reviewed.  
13  
14

15 **Data sharing statement:**

16 This article refers to a study protocol; data are not yet available. We will share all data through open access once  
17 the final results have been published.  
18  
19

20 **Ethical approval:** This study has received approval of the Medical Ethical Committee of the VU Medical Centre  
21 Amsterdam and confirmation can be supplied upon request [corresponding number: NL50951.029.14;  
22 November 5, 2015].  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1. Flowchart visits and case report forms

## Visits and Case report forms

	Intake + baseline	Surgery	1-2 days following treatment	2 weeks following treatment	4 weeks following treatment	6 weeks following treatment	3 months following treatment	6 months following treatment	9 months following treatment	12 months following treatment	24 months following treatment
Surgeon visit	X					X*					
Informed consent	X										
Research nurse visit	X					X	X			X	
Randomisation	X										
Surgery		X									
Discharge		X**	X**								
Four-Dimensional Symptom Questionnaire (4DSQ)	X										
Oswestry Disability Index (ODI)	X		X	X	X	X	X	X	X	X	X
Cost questionnaires	X			X	X	X	X	X	X	X	
EuroQol (EQ-5D-5L)	X		X	X	X	X	X	X	X	X	X
VAS leg pain, VAS Back pain	X		X	X	X	X	X	X	X	X	X
Quality of Life VAS	X		X	X	X	X	X	X	X	X	X
Patient self-perceived recovery and satisfaction			X***	X	X	X	X	X	X	X	X
Short Form 36 (SF36)	X			X	X	X	X	X	X	X	X
Physical examination	X					X	X			X	
Revisit and complications											With occurrence

\* 6 weeks visit may be performed also by the research nurse depending on the normal protocol hospital.

\*\* Discharge form will be filled in depending on discharge moment.

\*\*\* Only self-perceived recovery is measured not self-perceived satisfaction.

Table 2. Non-inferiority margins

Outcome measurements	Expected differences	Non-inferiority margin
VAS leg pain (0-100 scale)	<5 **	5
ODI (0-100 scale)	<5 *	5
VAS low back pain (0-100 scale)	<5 *	5
SF36 (0-100)	<5*	5
Self-perceived recovery (% 1 and 2 on the 7-point Likert-scale)	<10%***	5
Patient satisfaction (% 4 on the 4-point Likert-scale)	<5%	5
Patient satisfaction (% 1 and 2 on the 7-points Likert-scale)	<5%	5
Scar satisfaction (1-10 scale)	<1	0.5
Patellar reflex (% normal reflexes)	<5%	5
Achilles reflex (% normal reflexes)	<5%	5
Straight leg raising test (% negative tests)	<5%	5
Cross straight leg raising (% negative tests)	<5%	5
Finger floor distance (cm)	<5	5
Muscle strength quadriceps (% normal muscular strength)	5%	5
Sensibility dermatomes L1-S1 (% normal sensibility)	5%	5
EQ-5D-5L	<0.05****	0.05
Costs (healthcare perspective)	<\$500,- ***	250
Costs (societal perspective)	<\$1500,- ***	500

\* Obtained from the literature <sup>13,30</sup>

\*\* Obtained from the literature <sup>13</sup>

\*\*\* Obtained from the literature <sup>30</sup>

\*\*\*\* Obtained from the literature <sup>31</sup>



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents\*

Section/item	Item No	Description	Addressed on page number
<b>Administrative information</b>			
Title	1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	___ 1 ___
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	___ 2 ___
	2b	All items from the World Health Organization Trial Registration Data Set	___ 2 ___
Protocol version	3	Date and version identifier	___ 4 ___
Funding	4	Sources and types of financial, material, and other support	___ 15 ___
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	___ 1 and 15 ___
	5b	Name and contact information for the trial sponsor	___ 1 ___
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	___ 15 ___
	5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	___ 13 en 15 ___

1  
2  
3 **Introduction**  
4

5 Background and	6a	Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention	_____ 2 and 3 _____
6 rationale			
7			
8	6b	Explanation for choice of comparators	_____ 2 and 3 _____
9			
10 Objectives	7	Specific objectives or hypotheses	_____ 3 _____
11			
12 Trial design	8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory)	_____ 3 and 4 _____ (see also design section)_____
13			
14			
15			
16			
17			

18 **Methods: Participants, interventions, and outcomes**  
19

20 Study setting	9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will be collected. Reference to where list of study sites can be obtained	_____ 4 _____
21			
22			
23 Eligibility criteria	10	Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and individuals who will perform the interventions (eg, surgeons, psychotherapists)	_____ 4 _____
24			
25			
26 Interventions	11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be administered	_____ 5 and 6 _____
27			
28			
29	11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose change in response to harms, participant request, or improving/worsening disease)	_____ 6 en 9 _____
30			
31			
32	11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence (eg, drug tablet return, laboratory tests)	_____ n.a. _____
33			
34			
35	11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	_____ 6 _____
36			
37 Outcomes	12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen efficacy and harm outcomes is strongly recommended	_____ 6-8 _____
38			
39			
40			
41			
42			
43			
44			
45			
46			
47			
48			
49			



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Participant timeline	13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure)	_____8_____
Sample size	14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	_____9_____
Recruitment	15	Strategies for achieving adequate participant enrolment to reach target sample size	_____9_____

**Methods: Assignment of interventions (for controlled trials)**

Allocation:

Sequence generation	16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	_____5_____
Allocation concealment mechanism	16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	_____5_____
Implementation	16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	_____5_____
Blinding (masking)	17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	_____5_____
	17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	_____N.A._____

**Methods: Data collection, management, and analysis**

Data collection methods	18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	_____6-11_____
-------------------------	-----	--	----------------

1			
2			
3		18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols
4			_____6_____
5			
6	Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol
7			_____9_____
8			
9			
10	Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol
11			_____9-12_____
12			
13		20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)
14			__10 and 12__
15		20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)
16			_____9-10_____
17			
18			
19	<b>Methods: Monitoring</b>		
20			
21	Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed
22			_ 9, 12, 15 and 16__
23			
24			
25			
26		21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial
27			_____9_____
28			
29	Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct
30			_____9,12 and 16__
31			
32	Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor
33			_____12_____
34			
35			
36	<b>Ethics and dissemination</b>		
37			
38	Research ethics approval	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval
39			_____12 and 15__
40			
41			
42			
43			
44			
45			
46			
47			
48			
49			

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Protocol amendments	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial registries, journals, regulators)	_____4_____
Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	_____4_____
	26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	_____N.A._____
Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	_____9_____
Declaration of interests	28	Financial and other competing interests for principal investigators for the overall trial and each study site	_____15 en 16_____
Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	___15 and 6_____
Ancillary and post-trial care	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	_____12_____
Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	_____12_____
	31b	Authorship eligibility guidelines and any intended use of professional writers	_____15_____
	31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	_____n.a._____
<b>Appendices</b>			
Informed consent materials	32	Model consent form and other related documentation given to participants and authorised surrogates	appendix of bmj open online submission system__

1 2 3 4 5	Biological specimens	33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	_____N.A._____
-----------------------	-------------------------	----	--	----------------

---

6 \*It is strongly recommended that this checklist be read in conjunction with the SPIRIT 2013 Explanation & Elaboration for important clarification on the items.  
7 Amendments to the protocol should be tracked and dated. The SPIRIT checklist is copyrighted by the SPIRIT Group under the Creative Commons  
8 "[Attribution-NonCommercial-NoDerivs 3.0 Unported](https://creativecommons.org/licenses/by-nc-nd/3.0/)" license.  
9

For peer review only

# BMJ Open

**The PTED study: design of a non-inferiority, randomised controlled trial to compare the effectiveness and cost-effectiveness of percutaneous transforaminal endoscopic discectomy (PTED) vs. open microdiscectomy for patients with a symptomatic lumbar disc herniation.**

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-018230.R1
Article Type:	Protocol
Date Submitted by the Author:	20-Sep-2017
Complete List of Authors:	Seiger, Ankie; Vrije Universiteit Amsterdam, Health Sciences Gadjradj, Pravesh; Erasmusmc, Neurosurgery Harhangi, Sanjay; Erasmus MC, Neurosurgery van Susante, Job; Ziekenhuis Rijnstate, Orthopedics Peul, Wilco; Leiden University Medical Center, Neurosurgery Tulder, Maurits; University of Amsterdam, Health Sciences De Boer, Michiel; Vrije Universiteit Amsterdam, Health sciences Rubinstein, Sidney; Vrije Universiteit Amsterdam, Health Science
<b>Primary Subject Heading</b>:	Surgery
Secondary Subject Heading:	Health economics, Epidemiology, Neurology
Keywords:	HEALTH ECONOMICS, NEUROSURGERY, ORTHOPAEDIC & TRAUMA SURGERY

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**The PTED study: design of a non-inferiority, randomised controlled trial to compare the effectiveness and cost-effectiveness of percutaneous transforaminal endoscopic discectomy (PTED) vs. open microdiscectomy for patients with a symptomatic lumbar disc herniation.**

Seiger A<sup>1</sup>, Gadgradj P.S.<sup>2,3</sup>, Harhangi B.S.<sup>2</sup>, van Susante J.L.C.<sup>4</sup>, Peul W.C.<sup>3</sup>, van Tulder M.W.<sup>1</sup>, de Boer M.R.<sup>1</sup>, Rubinstein S.M.<sup>1</sup>

<sup>1</sup> Department of Health Sciences, Faculty of Science, Amsterdam Public Health research institute, Vrije Universiteit Amsterdam, The Netherlands

<sup>2</sup> Department of Neurosurgery, Erasmus MC: University Medical Centre Rotterdam, The Netherlands

<sup>3</sup> Department of Neurosurgery, Leiden University Medical Centre, LUMC, The Netherlands

<sup>4</sup> Department of Orthopedics, Rijnstate Hospital, The Netherlands

**Corresponding author:**

Ankie Seiger, MSc

Vrije Universiteit Amsterdam, Faculty of Science, Department of Health Sciences

De Boelelaan 1085

1081 HV Amsterdam

a.seiger@vu.nl

+31 20 59 82745

(Word count: 4803)

**Keywords:** Surgery, lumbar disc herniation, open microdiscectomy, percutaneous transforaminal endoscopic discectomy, effectiveness, economic evaluation.

## ABSTRACT

**INTRODUCTION:** Lumbosacral radicular syndrome is often caused by a disc herniation. The standard surgical technique to remove a disc herniation is open microdiscectomy. An alternative technique is Percutaneous Transforaminal Endoscopic Discectomy (PTED), which is less invasive. In The Netherlands, PTED is not currently considered standard care and therefore, not reimbursed within public health insurance. A pragmatic, multi-centre, non-inferiority, randomised controlled trial has been designed to determine the effectiveness and cost-effectiveness of PTED versus open microdiscectomy for the treatment of a lumbar disc herniation.

**METHOD AND ANALYSIS:** In total, 682 patients between 18-70 years of age with > 10 weeks of radiating pain or with > 6 weeks of excessive radiating pain are to be recruited from participating centres. Patients must have an indication for surgery based upon an MRI demonstrating compression of the nerve root from a lumbar disc herniation. Patients are to be randomised to PTED or open microdiscectomy. The primary outcome is self-reported leg pain measured by the 0-100 mm Visual Analogue Scale. Secondary outcomes include self-reported health and functional status; back pain; self-perceived recovery; and a physical examination. Outcomes will be measured the day following surgery, at 2, 4, and 6 weeks, and at 3, 6, 9, 12 and 24 months. Physical examination will be performed at 6 weeks, and 3 and 12 months. An economic evaluation will be performed from a societal perspective and cost-questionnaires will be used (e.g. EQ-5D-5L). The data will be analysed longitudinally; the non-inferiority margin for the primary outcome is 5. Bootstrapping techniques will be used for the economic evaluation.

**ETHICS AND DISSEMINATION:** This study has received approval of the Medical Ethical Committee of the VU Medical Centre Amsterdam: NL50951.029.14. The results will be published in an international peer reviewed scientific journal.

**REGISTRATION:** This trial is registered at ClinicalTrials.gov: NCT02602093.

### Strengths and Limitations

- Large, multi-centre, pragmatic, randomised controlled trial
- Use of standardized and validated outcomes instruments
- Longitudinal and multi-level analysis
- Inclusion of an economic evaluation
- Potential performance bias due to the lack of blinding of patients and care providers

## INTRODUCTION

Lumbosacral radicular syndrome is a common health problem with a lifetime prevalence that varies from 12.2% to 43% and has a point prevalence ranging from 1.6% to 13.4%.<sup>1-6</sup> Lumbosacral radicular syndrome is often caused by a lumbar herniated disc<sup>7</sup> and is associated with a greater incidence of sickness benefit,<sup>8</sup> increased pain and disability, and poorer quality of life<sup>9</sup> than those with non-specific low-back pain. In cases of a disc herniation, lumbosacral radicular syndrome can be treated either conservatively or surgically.<sup>7</sup>

To remove the disc herniation, the standard surgical technique is open microdiscectomy. A more recently developed technique is percutaneous transforaminal endoscopic discectomy (PTED). In short, open microdiscectomy is performed under general anaesthesia and surgeons operate with a direct vision on the herniated disc, while PTED is conducted transforaminally. These patients undergo local anaesthesia and surgeons operate through a working cannula with an indirect vision via an endoscope. Based upon the current



1  
2  
3 literature, PTED is a safe method for the removal of a lumbar disc herniation.<sup>10</sup> Possible benefits of PTED versus  
4 open microdiscectomy are: 1) Decreased medical costs because patients are treated on an outpatient basis; 2) It  
5 is easier to remove intra- and extra-foraminal herniated discs; 3) There is less chance of scar formation; and 4)  
6 The technique is potentially more effective for obese patients. However, too few, large prospective studies have  
7 examined this in detail, therefore, the benefits may be speculated.<sup>11-13</sup>  
8  
9

10  
11 Despite that PTED is becoming more commonly used, there are still questions regarding its effect and the  
12 associated costs.<sup>11, 13, 14</sup> A recent systematic review<sup>11</sup> identified three randomised controlled trials (RCTs), which  
13 examined the effect of PTED compared to open microdiscectomy.<sup>15-17</sup> Their results suggest that there is low to  
14 very low quality evidence that PTED is not more effective than open microdiscectomy for self-reported back  
15 pain, leg pain, functional status, recovery, return-to-work, and satisfaction with surgery. Importantly, all three  
16 studies were of poor methodological quality and examined relatively few patients (i.e. ranging from 40 to 60  
17 individuals). A more recent study concluded that PTED shows similar results compared to open  
18 microdiscectomy.<sup>13</sup> However, this was a single-centre study; it was conducted by a surgeon with a keen interest  
19 in the results of PTED; it included patients over a long period of time (i.e. the study started in 2006 and was  
20 published in 2017) suggesting possible selection of patients included in the trial; and the inclusion criteria  
21 published in the original protocol were different from those in the final publication. Additionally, the economic  
22 evaluation of this study has not yet been published.<sup>18</sup> This makes it difficult to assess the cost-effectiveness.  
23 Therefore, discussion regarding the effectiveness and cost-effectiveness of PTED remains.  
24  
25  
26  
27  
28  
29

30 In The Netherlands, the effectiveness of PTED has been heatedly debated. According to the Dutch Health Care  
31 Institute, a new surgical technique must meet certain requirements in order to be reimbursed by the public  
32 health insurance system. The Health Care Institute promotes the quality of Dutch Health Care and advises the  
33 Ministry of Health, Welfare and Sport on the content of the public health insurance. Based on a review,<sup>11</sup> the  
34 Health Care Institute claimed there is insufficient evidence for PTED to be included for reimbursement from the  
35 public health insurance package and as a result, patients are forced to pay the costs of the PTED treatment out-  
36 of-pocket. In 2017, the PTED and open microdiscectomy costs are approximately €5000 and €3000, respectively.  
37 In order to deal with this issue and to answer the remaining questions about PTED, this large, pragmatic,  
38 methodologically rigorous multi-centre study has been designed. The costs of PTED and open microdiscectomy  
39 will be fully reimbursed by the Dutch health insurance companies for patients participating in this study.  
40  
41  
42  
43

44 This study is expected to have a major societal impact because it will determine if PTED should be included in  
45 the Dutch health insurance package. Furthermore, this study will provide more insight in PTED internationally,  
46 resulting in improved care for patients with a lumbar disc herniation. The primary hypothesis of this study is that  
47 PTED is not less effective and not less cost-effective compared to standard care (i.e. open microdiscectomy) for  
48 patients with symptomatic, lumbosacral radicular syndrome as a result of a lumbar disc herniation. Therefore, a  
49 non-inferiority design will be used.  
50  
51  
52

## 53 **METHOD AND ANALYSIS**

### 54 **Study design**

55  
56  
57  
58  
59  
60

1  
2  
3 A pragmatic, multi-centre non-inferiority randomised controlled trial (RCT) will be used. Following the baseline  
4 measurements, wherein clinical and socio-demographic measurements will be collected, patients are to be  
5 randomised to one of the two groups: the control group will receive standard open microdiscectomy and the  
6 intervention group will receive PTED. Patients will be followed for two years, but the primary analysis will be  
7 conducted on the one-year data.  
8  
9

10  
11 Important protocol modifications will be registered at ClinicalTrials.gov and communicated to all relevant parties  
12 involved in this study (Medical Ethical Committee, ZonMw, included patients, participating surgeons, and  
13 members from the advisory board (listed in the acknowledgments)).  
14

### 15 16 17 **Study population**

18 In total, 682 patients with a magnetic resonance imaging (MRI) confirmed lumbosacral radicular syndrome, due  
19 to a lumbar disc herniation are to be recruited. Patients will be recruited from five hospitals and one private  
20 health clinic located in Arnhem/Zevenaar, Leiderdorp, Tilburg and Rotterdam (The Netherlands). Each patient is  
21 required to sign a written informed consent prior to participation.  
22

23  
24 In order to be eligible to participate and in accordance with the Dutch Guideline Lumbosacral Radicular  
25 Syndrome,<sup>19</sup> a subject must meet all of the following conditions:  
26  
27

### 28 29 **Inclusion criteria**

- 30 • 18-70 years of age;
- 31 • > 10 weeks of radiating pain with or without motor or sensory loss in the leg, or with > 6 weeks of  
32 excessive radiating pain and no tendency for any clinical improvement;
- 33 • Indication for surgery;
- 34 • MRI demonstrating a lumbar disc herniation with nerve compression with or without concomitant  
35 spinal or lateral recess stenosis or sequestration;
- 36 • Sufficient knowledge of the Dutch language in order to complete forms and follow instructions  
37 independently.  
38  
39  
40  
41

### 42 43 **Exclusion criteria**

- 44 • Previous surgery on the same or adjacent disc level;
- 45 • Cauda equina syndrome;
- 46 • Spondylitic- or degenerative spondylolisthesis;
- 47 • Pregnancy;
- 48 • Severe comorbid medical or psychiatric disorder (American Society of Anaesthesiologists >2);
- 49 • Severe caudal or cranial sequestration;
- 50 • Contra-indication for surgery;
- 51 • Moving abroad at short notice.  
52  
53  
54

### 55 56 **Study procedures** 57 58 59 60

1  
2  
3 Participating surgeons will screen all eligible patients with a lumbar disc herniation during the consultation  
4 (Table 1). If eligible for inclusion, patients will receive information relevant to the study by means of a letter.  
5 Dutch law requires that patients are given at least two days to consider participation. Following this initial  
6 screening, the patient is to be examined by a trained research nurse and the informed consent is obtained;  
7 baseline measurements will be performed; and patients will be randomised.  
8  
9

### 10 11 **Randomisation**

12 Patients will be randomised in a 1:1 ratio to PTED or open microdiscectomy. An experienced statistician will  
13 prepare computer-generated, random number tables. Treatment allocation will be concealed. The key will be  
14 withheld from all participants and researchers involved in this study. Variable block sizes of 4, 6 and 8 will be  
15 used and stratified by treatment centre. The random-number tables will be entered into a computer system by  
16 an independent software company, and allocation will be performed by the computer system once baseline  
17 data and physical examination are to be obtained from an independent research nurse responsible for the  
18 treatment allocation.  
19  
20  
21

### 22 23 **Blinding**

24 No attempt will be made to blind the patients. Blinding is considered impossible, because the procedures are  
25 fundamentally different. Furthermore, outcomes assessors cannot be blinded, given that the primary outcomes  
26 are all self-reported. The analysis will be performed blinded for treatment allocation.  
27  
28

### 29 30 **Treatment**

#### 31 *Intervention: Percutaneous Transforaminal Endoscopic Discectomy (PTED)*

32 PTED is to be conducted as follows:<sup>20</sup> local anaesthesia is to be administered and consists of light sedation with  
33 dexmedetomidine or a combination of propofol and remifentanyl for the convenience of the patient. The  
34 amount of administered sedation should still allow the patient to respond to nerve root manipulation.  
35 Verification of the site is to be performed by an image intensifier using fluoroscopy (anteroposterior and lateral  
36 view) and is depending upon the patient's posture. An incision just above the dorsolateral side of the pelvis is  
37 conducted, where a needle is to be set from the incision to the superior articular process of the lower involved  
38 vertebrae of the herniated disc. Position will be checked again under fluoroscopy. After the needle has reached  
39 the superior articular process, a guide wire is to be inserted. Following that, a series of conical rods are to be  
40 introduced, subsequently a drill/reamer is to be introduced through the cannula and rods. After drilling through  
41 the superior articular process is conducted in order to enlarge the neuroforamen, the instruments are to be  
42 removed, but the guidewire is left in place and the endoscope with the working channels are to be introduced  
43 via an 8 mm cannula. The image intensifier ensures that the position of the cannula is maintained. Following  
44 removal of the disc herniation with a rongeur, the cannula and endoscope are removed. The patient is to be  
45 treated on an outpatient basis. In order to decompress the nerve root, it is sometimes necessary to remove the  
46 superior articular process. With the outside-in technique this can be successfully performed.<sup>10, 20</sup>  
47  
48  
49  
50  
51  
52

#### 53 *Comparison: Open microdiscectomy*

54 Open microdiscectomy is to be conducted as follows: general or spinal anaesthesia is to be administered.  
55 Verification is to be performed using a fluoroscopy and the patient is to be positioned prone or in the salaam  
56 position. Loupe or microscope magnification may be used according to the surgeon's preference. A paramedian  
57  
58  
59  
60

1  
2  
3 incision is to be performed and the level is to be indicated. Following the identification of the lamina, the yellow  
4 ligament will be removed to identify the nerve root and disc herniation. Laminotomy as well as foraminotomy is  
5 to be performed, if necessary. For the foraminal herniated disc we will use a partial medial facetectomy and for  
6 the  
7  
8 extra-foraminal herniated disc a parafacetal approach. For all surgeries, the amount of degenerative disc  
9 material shall be removed at the discretion of the attending surgeon. Post-operative policy will be followed and  
10 it is expected that the duration of recovery in the hospital may vary from 1-2 days, but the patient will be  
11 discharged as soon as medically responsible.  
12

### 13 *Co-interventions*

14 Pain medication will be offered to patients, should this be necessary. In addition, use of co-interventions will be  
15 monitored by self-reported cost questionnaires, in which medication usage and any health care utilization is  
16 recorded throughout the follow-up period.  
17  
18

### 19 **Learning curve**

20 It will be necessary to train surgeons in the use of PTED. Prior to the start of this study only two surgeons in The  
21 Netherlands were proficient in this technique. One of these surgeons is participating in this study (BSH). This  
22 experienced surgeon will provide the training to the other surgeons, all of whom have more than ten years of  
23 surgical experience. The initial training will be first conducted on cadavers, and only once the surgeons are  
24 comfortable with the use of the procedure, they will then perform this technique on patients under the tutelage  
25 of the PTED-experienced surgeon. It is expected that 50 patients per surgeon will be necessary to become  
26 proficient in PTED (defined as the 'learning curve'). Thus, 150 PTED patients will be registered as learning curve  
27 patients. Additionally, competency in the use of PTED by the surgeons is to be evaluated using skill-based  
28 questions measured by a Likert-scale and the Objective Structured Assessment of Technical Skills (OSATS). These  
29 will be recorded and evaluated by both the teaching surgeon as well as the surgeons undergoing the training.<sup>21</sup>  
30  
31  
32  
33  
34  
35

### 36 **Prognostic factors**

37 The following potential prognostic factors are to be measured: 1) socio-demographic characteristics (e.g. age,  
38 gender); 2) characteristics of the complaint (e.g. duration and severity); 3) baseline pain and functional disability;  
39 4) lifestyle factors (e.g. smoking and alcohol use); 5) psychological factors (e.g. expectations of recovery,  
40 emotional well-being); 6) psychopathology as measured with the four dimensional symptom questionnaire  
41 (4DSQ; dimensions: distress, depression, anxiety and somatization);<sup>22</sup> 7) work-related factors (e.g. physical  
42 workload, job satisfaction); and 8) previously received treatment due to the same episode of back complaints  
43 (e.g. medication and physiotherapy).  
44  
45  
46  
47

### 48 **Outcome measurements**

49 The outcomes are to be measured by validated self-reported questionnaires and by physical examination. Data  
50 are to be collected prior to randomisation (baseline), the day following surgery, at 2, 4, and 6 weeks, and at 3, 6,  
51 9, 12, and 24 months following surgery (Table 1).  
52

53 All questionnaires will be sent automatically by e-mail with a personal link to the digital questionnaire. If  
54 necessary, a reminder will be sent after 3 days; after six days the research nurse will call the patient with the  
55 request to fill in the questionnaire. Deviations from the protocol (e.g. conversion from PTED to open  
56 microdiscectomy) will be registered and outcomes will continue to be measured.  
57  
58  
59  
60

1  
2  
3  
4  
5 Primary outcomes

6 The primary outcome, leg pain, is to be measured by the Visual Analogue Scale (VAS; scale 0-100 mm). This  
7 outcome measure has been identified in a systematic review to be one of the most commonly measured  
8 outcomes, and is specific and responsive to change in a population undergoing lumbar spine surgery.<sup>23</sup>  
9

10  
11 Secondary outcomes

12 *Functional status:* will be measured with the Oswestry Disability Index (ODI). The ODI<sup>24</sup> is one of the principal  
13 condition-specific outcome measures used in the management of spinal disorders. The ODI (2.1a) is to be used.<sup>25</sup>  
14 The ODI has been extensively tested and showed good psychometric properties.<sup>26</sup>  
15

16  
17  
18 *Low-back pain:* will be measured with the VAS (scale ranging from 0mm (no pain) to 100mm (worst imaginable  
19 pain)).  
20

21  
22 *Generic quality of life:* will be measured with the Dutch version of the short form SF36. The SF-36 questionnaire  
23 has been validated and found reliable for low back pain.<sup>27</sup> The questions are divided into eight domains: 1)  
24 physical functioning, 2) physical role limitations, 3) emotional role limitations, 4) social functioning, 5) physical  
25 pain, 6) general mental health, 7) vitality, and 8) general health perception. Per domain the scores of the items  
26 are added up and transformed into a scale of 0 to 100. A higher score reflects a better health condition. In  
27 addition, these eight domains can be summarized in a physical and psychological main domain.  
28  
29

30  
31 *Self-perceived recovery of the patient:* will be measured with a seven-point Likert-scale. The score on this scale  
32 varies from 'completely recovered' to 'worse than ever'. We will dichotomize the outcome with 'completely  
33 recovered', 'moderately recovered' and 'a bit recovered' as 'recovered' and the other four categories as 'not  
34 recovered'.  
35

36  
37  
38 *Patient satisfaction:* will be measured using the Likert-scale, Body Image and the Cosmesis scale.<sup>28, 29</sup> Body  
39 satisfaction will be measured using a four-point Likert-scale (ranging from 'not at all', 'a little', 'quite', to 'yes,  
40 very much'). Satisfaction change of complaints and satisfaction treatment will be measured using a seven-point  
41 Likert-scale (ranging from 'completely satisfied with current symptoms' to 'completely dissatisfied with current  
42 symptoms'). The scales will be completed by the patients prior to and following surgery. Scar satisfaction will  
43 also be measured using the seven-point Likert-scale and with a 1-10 numeric rating scale (ranging from 1 = 'as  
44 ugly as conceivable' to 10 = 'almost no scar perceived').  
45  
46  
47

48  
49 *Physical examination:* will be performed at 6 weeks, and at 3 and 12 months following surgery. This will include:  
50 scar size; patellar and Achilles tendon reflexes; straight leg raising test; cross straight leg raising test; finger-floor  
51 distance; strength measurement of the quadriceps using the Medical Research Council (MRC); sensibility  
52 dermatomes L1-S1, abdominal muscle strength; and patients' weight. The patellar and tendon reflexes are to be  
53 measured in a sitting upright position with both feet dangling above the ground. Tendon reflexes are tapped up  
54 to a maximum of two times with the reflex hammer. Reflexes are distinguished into absent, reduced, normal,  
55 increased, and clonus reflexes. The straight leg raising test and cross straight leg raising test are both measured  
56 as negative when no shooting leg pain is perceived, and positive when shooting pain is perceived. Finger-floor  
57  
58  
59  
60

distance is the distance between the longest finger and the floor when the patients perform a forward bent with the knees extended. Muscle strength of the quadriceps is measured from a sitting position. Patients will be asked to extent their knee while the research nurse exerts counter-pressure just above the ankle. Muscle strengths are rated on the Dutch version of the MRC, ranging from 0= no contraction to 5 = normal muscle strength. For the sensibility the research nurse checks every dermatome area (L1-S1) by touching the patient with a sharp and blunt object. Patients indicate with their eyes closed when sensation is felt. Sensibility varies from decreased, normal or increased sensibility compared to the other leg. Abdominal strength is measured by counting the maximal number of abdominal crunches from the supine position. Patients are asked to reach the hands towards the bended knees and to lift the scapulae from the surface. At any time, the lumbar spine will be supported by the underlying surface to minimize the range of motion of the lumbar spine. Without an increase of pain the maximum is set at a cut-off point of 26 crunches.

Screening and operation case record forms are to be completed by the surgeons, while discharge forms, physical examination and baseline intake forms are to be completed by a trained research nurse.

Table 1. Flowchart visits and case report forms

Visits and Case report forms

	Intake + baseline	Surgery	1-2 days following treatment	2 weeks following treatment	4 weeks following treatment	6 weeks following treatment	3 months following treatment	6 months following treatment	9 months following treatment	12 months following treatment	24 months following treatment
Surgeon visit	X					X*					
Informed consent	X										
Research nurse visit	X					X	X			X	
Randomisation	X										
Surgery		X									
Discharge		X**	X**								
Four-Dimensional Symptom Questionnaire (4DSQ)	X										
Oswestry Disability Index (ODI)	X		X	X	X	X	X	X	X	X	X
Cost questionnaires	X			X	X	X	X	X	X	X	
EuroQol (EQ-5D-5L)	X		X	X	X	X	X	X	X	X	X
VAS leg pain, VAS Back pain	X		X	X	X	X	X	X	X	X	X
Quality of Life VAS	X		X	X	X	X	X	X	X	X	X
Patient self-perceived recovery and satisfaction			X***	X	X	X	X	X	X	X	X
Short Form 36 (SF36)	X			X	X	X	X	X	X	X	X
Physical examination	X					X	X			X	
Revisit and complications											

\* 6 weeks visit may be performed also by the research nurse depending on the normal protocol hospital.

\*\* Discharge form will be filled in depending on discharge moment.



1  
2  
3 \*\*\* Only self-perceived recovery is measured not self-perceived satisfaction.  
4

### 5 **Complications, operative morbidity, and re-operations**

6 Immediately following surgery and discharge, the surgeon and research nurse will perform a systematic  
7 assessment of complications (including urinary tract infection, secondary bleeding, and progressive neurological  
8 deficit). In addition, surgeons will record any perioperative complications like cerebrospinal fluid leakage, nerve  
9 root damage, and if the surgery was initiated at the wrong disc level. Re-operation at the initial site is to be  
10 considered a poor outcome. Re-operation in both groups will be recorded. Perioperative morbidity will be  
11 assessed with operation time, perioperative blood loss, hospital stay and re-operative rate as related to the  
12 primary condition (lumbar disc herniation).  
13  
14  
15  
16

### 17 **Sample size calculation**

18 The mean difference and standard deviation (SD) for the VAS (leg pain) used in the sample size calculation was:  
19 mean 5; SD 14.9.<sup>30</sup> The margin of non-inferiority was set at 5, (one-sided) alpha at 0.05 and beta at 0.10 (power  
20 0.9). We estimated that in total 306 patients are needed to demonstrate non-inferiority on the primary  
21 outcome. Accounting for 20% attrition, the aim is to recruit 382 patients. As the Ministry of Health in the  
22 Netherlands has stipulated that PTED will only be reimbursed if patients participate in the randomised trial, an  
23 extra 300 patients will be necessary for the inclusion of 150 patients in the PTED learning curve. Consequently, a  
24 total of 682 patients will be recruited for this study. Patients are likely to participate in this study, because PTED  
25 will only be reimbursed by Dutch health care insurance for participants in this study. Therefore, reaching the  
26 target sample size is not likely to become a problem.  
27  
28  
29  
30  
31

### 32 **Data analysis**

33 Data analysis will be conducted by a researcher or statistician blinded for treatment allocation after follow-up is  
34 finished. No interim analysis will be performed.  
35

36 All data handling (entry, coding, storage and analysis) is confidential and complies with the Dutch Personal Data  
37 Protection Act. The anonymous data are stored in a central warehouse for at least 15 years.  
38

### 39 **Effect analysis**

40 Characteristics of the patients will be presented using descriptive statistics (mean (SD), median (range) or  
41 proportion) to assess if balanced groups were obtained after randomisation. The non-inferiority margins are set  
42 and listed in table 2.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Table 2. Non-inferiority margins

Outcome measurements	Expected differences	Non-inferiority margin
VAS leg pain (0-100 scale)	<5 **	5
ODI (0-100 scale)	<5 *	5
VAS low back pain (0-100 scale)	<5 *	5
SF36 (0-100)	<5*	5
Self-perceived recovery (% 1 and 2 on the 7-point Likert-scale)	<10%***	5
Patient satisfaction (% 4 on the 4-point Likert-scale)	<5%	5
Patient satisfaction (% 1 and 2 on the 7-points Likert-scale)	<5%	5
Scar satisfaction (1-10 scale)	<1	0.5
Patellar reflex (% normal reflexes)	<5%	5
Achilles reflex (% normal reflexes)	<5%	5
Straight leg raising test (% negative tests)	<5%	5
Cross straight leg raising (% negative tests)	<5%	5
Finger floor distance (cm)	<5	5
Muscle strength quadriceps (% normal muscular strength)	5%	5
Sensibility dermatomes L1-S1 (% normal sensibility)	5%	5
EQ-5D-5L	<0.05****	0.05
Costs (healthcare perspective)	<\$500,- ***	250
Costs (societal perspective)	<\$1500,- ***	500

\* Obtained from the literature <sup>13,30</sup>

\*\* Obtained from the literature <sup>13</sup>

\*\*\* Obtained from the literature <sup>30</sup>

\*\*\*\* Obtained from the literature <sup>31</sup>

The primary data analysis will examine the effects of PTED for leg pain for those patients not in the learning curve, and shall be conducted according to the intention-to-treat principle. If necessary, missing items will be imputed using multiple imputation techniques. Linear and generalized multi-level analyses will be used, accounting for dependency of measurements over time within patients and patients nested within the surgeons, representing a 3-level model: time, patient and surgeon. The data are to be examined longitudinally and the primary analysis will be aimed at average differences in effectiveness between the two treatment modalities. We will also include treatment \* time interactions to explore whether these effects are different over time. In addition to the crude analyses, all analyses will be adjusted for potential confounders, such as age, gender, nature and severity of the presenting complaint. In a secondary analysis, a per-protocol analysis shall be conducted. The secondary continuous outcomes, such as low-back pain, functional status, will be analysed similar to the primary data analysis; however, recovery and some of the physical performance measures (Table 2) are to be treated as a dichotomous variable and will be analysed in logistic regression analyses.

Complications will be summarized for the time period of the study, but also presented for those complications encountered before and after 6 weeks.

#### *Subgroup analyses effect*

Subgroup analysis will be conducted in those patients with a) paramedian/median disc herniation b) foraminal/extra-foraminal disc herniation, and c) L5-S1 disc herniation. The goal of the subgroup analyses is to test the robustness of the data to changes in underlying assumptions regarding the type or location of the

hernia. In addition, we will examine the effects for all patients, including those in the 'learning curve' in order to determine if these outcomes are different than the primary analyses.

Results from all analyses will be expressed as mean effect estimate with 95% confidence intervals and these estimates will be subsequently compared to the margin of non-inferiority in order to make inferences about the non-inferiority of the intervention, PTED.

### **Economic evaluation**

Both cost-effectiveness and cost-utility analysis will be conducted from a societal perspective alongside the RCT. We will measure, value, and analyse total costs of all patients and relate the difference in costs to the difference in effects between the two groups.

Direct costs include costs of the interventions, hospitalisation after surgery, medication and other health care utilization. Patient costs and cost of productivity loss, absenteeism and presenteeism, will also be included. Health care utilization, patients cost and productivity loss will be measured using self-completed cost questionnaires. The cost of the interventions will be estimated using a bottom-up approach (micro-costing) and hospitalisation will be registered using case record forms. The Dutch tariff of the EQ-5D-5L will be used to calculate the quality-adjusted life years (QALYs).<sup>32, 33</sup> The EuroQol measures the five dimensions: mobility, self-care, daily activities, pain/discomfort, and anxiety/depression. Each dimension consists of one item, while five levels are distinguished ('no', 'slight', 'moderate', 'severe problems', 'unable to do').

Costs resulting from productivity loss are to be estimated using the friction cost method, which assumes that sick workers are replaced after a period of time (i.e. 12 weeks).<sup>34</sup> Mean productivity costs per working hour are to be adjusted for age and gender and used to estimate the cost of absenteeism. Health care utilization is to be valued according to the guidelines published in the updated handbook for economic evaluation in The Netherlands.<sup>34</sup> Medication is to be valued using prices from the Royal Dutch Society for Pharmacy.<sup>35</sup>

#### *Cost-effectiveness analysis*

Total costs will be related to the primary effect measure, leg pain. A cost-utility analysis will be performed with QALYs. From the EQ-5D-5L utilities will be obtained and QALYs will be calculated using linear interpolation between measurement points. The primary analysis will be conducted according to intention-to-treat. Missing data will be imputed using multiple imputation by chained equations.<sup>36</sup> Incremental Cost Effectiveness Ratios (ICERs) will be calculated by dividing the difference in costs by the difference in effects. We will perform a cost-effectiveness analysis with leg pain and a cost-utility analysis with QALYs as outcome. In order to account for the possible clustering of data, analyses will be performed using linear multilevel analyses.<sup>37</sup> Bias corrected and accelerated bootstrapping with 5,000 replications will be performed in order to estimate 95% confidence intervals around cost differences and the uncertainty surrounding the ICERs. Uncertainty will be shown in cost-effectiveness planes and cost-effectiveness acceptability curves, and sensitivity analyses will be performed to test the robustness of the study results.<sup>38-40</sup>

#### *Sensitivity analysis cost-effectiveness*

1  
2  
3 Sensitivity analyses will be conducted for the most important cost drivers in order to determine the robustness  
4 of the findings. In addition, the main analyses are to be repeated using only complete cases (i.e. complete  
5 clinical outcome data and complete cost data). Lastly, the impact of the Human Capital Approach will be  
6 compared to the friction cost method approach. The Human Capital Approach evaluates the total costs of  
7 productivity loss without considering the possibility of replacing the sick worker.  
8  
9

## 10 11 **ETHICS AND DISSEMINATION**

12 This study has received approval of the Medical Ethical Committee of the VU Medical Centre Amsterdam and  
13 confirmation can be supplied upon request [corresponding number: NL50951.029.14; November 5, 2015].  
14 Serious adverse events (SAE) and adverse events will be registered; SAE will be reported within 24 hours (see  
15 section 'complications, operative morbidity, and re-operations'). The sponsor has an insurance, which is in  
16 accordance with the legal requirements in the Netherlands. The insurance applies to the damage that becomes  
17 apparent during the study or within four years after the end of the study. This study will be monitored according  
18 to a detailed monitoring plan adapted to the risk classification of the Dutch University Federation guidelines.  
19 Based on this guideline, the risk classification of this study is regarded negligible. Considerations in this  
20 assessment are that this is an investigator-initiated trial, not with vulnerable patients, and while side effects are  
21 known, such as nerve root damage, severe adverse events are extremely rare. Audits may be required by and  
22 will be granted to the Medical Ethical Committee and to the regulatory authority inspections. Patients'  
23 permission for these audits is obtained with informed consent.  
24  
25  
26  
27  
28  
29

30 The final trial results will be communicated to participants, healthcare professionals, professional organisations  
31 and relevant guideline committees in the Netherlands. We will publish the results in an international peer  
32 reviewed open access scientific journal. There are no publication restrictions.  
33  
34

## 35 **DISCUSSION**

36 This large, multi-centre, pragmatic study will be conducted to resolve the discussion regarding the effects and  
37 costs of PTED compared to open microdiscectomy for patients with a lumbosacral radicular syndrome caused by  
38 a lumbar disc herniation. Learning curve patients and foraminal/extraforaminal disc herniations will be included,  
39 because the Dutch Ministry of Health and the Netherlands Organisation for Health and Research Development  
40 (ZonMw) requested this. At the moment, in the Netherlands PTED does not comply with standards of practice  
41 and is not included in the Dutch public healthcare package. However, The Dutch Ministry of Health classified  
42 PTED as an important technique to examine and decided that PTED will be conditionally admitted to the Dutch  
43 public health insurance package for those patients participating in this study. In other words, insurance  
44 companies are obliged to reimburse PTED for patients participating in this study. This conditional  
45 reimbursement only applies during the four years of this study. After this study a decision will be made if PTED  
46 should or should not be included in the Dutch public health insurance package. Open microdiscectomy is already  
47 included in the Dutch public health insurance package and reimbursed for all patients. This advantage of  
48 reimbursement presented a unique challenge, because this means that all patients are to be included from the  
49 beginning of this agreement. Thus, patients will be included also when surgeons are still undergoing the PTED  
50 training (learning curve patients). The other requirement was that the inclusion and exclusion criteria had to be  
51 in accordance with the Dutch Guideline Lumbosacral Radicular Syndrome and similar to an earlier study  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 performed in this field.<sup>19, 41</sup> For this reason, also patients with foraminal and extraforaminal disc herniations will  
4 be included in this study. Extra subgroup analysis will be performed in order to assess possible differences in  
5 effect. In order to prevent discussion regarding the effects of PTED following this study, a document was signed  
6 by all participating parties (e.g. professional surgical organisations, insurance companies, Dutch healthcare  
7 institute) to agree upon the study design and the criteria for in- or exclusion of PTED from the Dutch public  
8 health insurance package.  
9  
10

11  
12 Since the trial was published in a trial registry (clinicaltrials.gov; November 2013), the protocol has been  
13 modified. A physical examination has been added in order to obtain more objective information regarding the  
14 physical rehabilitation and a numeric rating scale (NRS) has been added in order to measure back, leg pain and  
15 quality of life. The reason for the latter is, the VAS may be completed by participants using different digital  
16 apparatuses (i.e. PC, tablet or mobile phone), with the result that the lengths of the VAS scale may vary. The  
17 validity and reliability is, therefore, uncertain. Based upon the literature, it would appear that the NRS and VAS  
18 demonstrate comparable values for pain following surgery;<sup>42</sup> however, the aforementioned issue, namely the  
19 use of the VAS on different digital apparatuses has not previously been examined. In order to determine  
20 whether this has bearing on the outcomes, a sensitivity analysis shall be conducted, and we will examine the  
21 correlation between the VAS and NRS.  
22  
23  
24  
25  
26

27 The trial is an ongoing study and runs from February 2016 to February 2020.  
28  
29

### 30 Acknowledgements:

31 Dr. C.L.A.M. Vleggeert-Lankamp, Drs. P.R.A.M. Depauw, Drs. P.J. Schutte, Drs. A.W.J. Vreeling; the hospitals,  
32 Albert Schweitzer Ziekenhuis, Alrijne Ziekenhuis, Elisabeth-Tweesteden Ziekenhuis, Franciscus Gasthuis en  
33 Vlietland, Park Medisch Centrum, Rijnstate Ziekenhuis; and the Advisory Committee: Dr. J.L.P. Kuijpers,  
34 Zorgverzekeraars Nederland (ZN), patient organisation 'Nederlandse Vereniging van Rugpatiënten de  
35 "Wervelkolom"', Dutch Spine Society, Nederlandse Orthopaedische Vereniging.  
36  
37  
38

### 39 REFERENCES

- 40 1. Heliovaara, M, Impivaara, O, Sievers, K et al. Lumbar disc syndrome in Finland. *J Epidemiol Community*  
41 *Health*1987;41:251-8.
- 42 2. Konstantinou, K and Dunn, KM. Sciatica: review of epidemiological studies and prevalence estimates.  
43 *Spine (Phila Pa 1976)*2008;33:2464-72.
- 44 3. Nielen, MMJ. Incidentie en prevalentie van gezondheidsproblemen in de Nederlandse huisartsenpraktijk  
45 in 2014. *NIVEL Zorgregistraties eerste lijn*2015. ICPC L86. <https://www.nivel.nl/node/4309> (date  
46 accessed June 2017).
- 47 4. Spijker-Huiges, A, Groenhof, F, Winters, JC et al. Radiating low back pain in general practice: incidence,  
48 prevalence, diagnosis, and long-term clinical course of illness. *Scand J Prim Health Care*2015;33:27-32.
- 49 5. Oosterhuis, T, Smaardijk, V, Kuijer, P et al. Actualisatie wetenschappelijke kennis en inzichten over het  
50 lumbosacraal radiculair syndroom. *Uitvoeringsinstituut werknemersverzekeringen UWV*2016; 16-06.  
51 <https://www.uwv.nl/overuwv/Images/Actualisatie-lumbosacraal-radiculair-syndroom.pdf> (date  
52 assessed June 2017).
- 53 6. Schaafstra, A, Spinnewijn, WEM, Bons, S et al. NHG-Standaard Lumbosacraal radiculair syndroom  
54 (tweede herziening). *Huisarts & Wetenschap*2015;58:308-20.
- 55 7. Koes, BW, van Tulder, MW, and Peul, WC. Diagnosis and treatment of sciatica. *BMJ*2007;334:1313-7.  
56  
57  
58  
59  
60

- 1
- 2
- 3
- 4 8. Kristman, VL, Hartvigsen, J, Leboeuf-Yde, C et al. Does Radiating Spinal Pain Determine Future Work
- 5 Disability? A Retrospective Cohort Study of 22,952 Danish Twins. *Spine*2012;37:1003-1013.
- 6
- 7 9. Konstantinou, K, Hider, SL, Jordan, JL et al. The impact of low back-related leg pain on outcomes as
- 8 compared with low back pain alone: a systematic review of the literature. *Clin J Pain*2013;29:644-54.
- 9
- 10 10. Gadraj, PS, van Tulder, MW, Dirven, CM et al. Clinical outcomes after percutaneous transforaminal
- 11 endoscopic discectomy for lumbar disc herniation: a prospective case series. *Neurosurg*
- 12 *Focus*2016;40:E3.
- 13
- 14 11. Kamper, SJ, Ostelo, RW, Rubinstein, SM et al. Minimally invasive surgery for lumbar disc herniation: a
- 15 systematic review and meta-analysis. *Eur Spine J*2014;23:1021-43.
- 16
- 17 12. Gu, YT, Cui, Z, Shao, HW et al. Percutaneous transforaminal endoscopic surgery (PTES) for symptomatic
- 18 lumbar disc herniation: a surgical technique, outcome, and complications in 209 consecutive cases. *J*
- 19 *Orthop Surg Res*2017;12:25.
- 20
- 21 13. Gibson, JNA, Subramanian, AS, and Scott, CEH. A randomised controlled trial of transforaminal
- 22 endoscopic discectomy vs microdiscectomy. *European Spine Journal*2017;26:847-856.
- 23
- 24 14. Pan, Z, Ha, Y, Yi, S et al. Efficacy of Transforaminal Endoscopic Spine System (TESSYS) Technique in
- 25 Treating Lumbar Disc Herniation. *Med Sci Monit*2016;22:530-9.
- 26
- 27 15. Hermantin, FU, Peters, T, Quartararo, L et al. A prospective, randomized study comparing the results of
- 28 open discectomy with those of video-assisted arthroscopic microdiscectomy. *J Bone Joint Surg*
- 29 *Am*1999;81:958-65.
- 30
- 31 16. Mayer, HM and Brock, M. Percutaneous endoscopic discectomy: surgical technique and preliminary
- 32 results compared to microsurgical discectomy. *J Neurosurg*1993;78:216-25.
- 33
- 34 17. Krappel, FA, Schmitz, R, Bauer, E et al. Open or endoscopic nucleotomy? Results of a prospective,
- 35 controlled clinical trial with independent follow-up, MRI and special reference to cost-effectiveness.
- 36 *Orthopadische Prax*2001;37:164-169.
- 37
- 38 18. Scott, CEH and Gibson, AJN. A cost utility comparison of transforaminal endoscopic discectomy and
- 39 microdiscectomy. *The Spine Journal*2016;16:S47.
- 40
- 41 19. NVN and CBO. Richtlijn lumbosacraal radiculair syndroom. *Nederlandse Vereniging voor Neurologie en*
- 42 *Kwaliteitsinstituut voor de Gezondheidszorg CBO*2008.
- 43 [http://www.anesthesiologie.nl/uploads/openbaar/RL\\_Lumbosacraal\\_Radiculair\\_Syndroom\\_2008.pdf](http://www.anesthesiologie.nl/uploads/openbaar/RL_Lumbosacraal_Radiculair_Syndroom_2008.pdf)
- 44 (date assessed June 2017).
- 45
- 46 20. Gadraj, PS and Harhangi, BS. Percutaneous Transforaminal Endoscopic Discectomy for Lumbar Disk
- 47 Herniation. *Clin Spine Surg*2016;29:368-371.
- 48
- 49 21. Hiemstra, E and F.W., J. Richtlijn OSATS. *Tijdschrift voor Medisch Onderwijs*2010;29:135-141.
- 50
- 51 22. Terluin, B, van Marwijk, HW, Ader, HJ et al. The Four-Dimensional Symptom Questionnaire (4DSQ): a
- 52 validation study of a multidimensional self-report questionnaire to assess distress, depression, anxiety
- 53 and somatization. *BMC Psychiatry*2006;6:34.
- 54
- 55 23. DeVine, J, Norvell, DC, Ecker, E et al. Evaluating the correlation and responsiveness of patient-reported
- 56 pain with function and quality-of-life outcomes after spine surgery. *Spine (Phila Pa 1976)*2011;36:S69-
- 57 74.
- 58
- 59 24. Fairbank, JC, Couper, J, Davies, JB et al. The Oswestry low back pain disability questionnaire.
- 60 *Physiotherapy*1980;66:271-3.
25. van Hooff, ML, Spruit, M, Fairbank, JC et al. The Oswestry Disability Index (version 2.1a): validation of a Dutch language version. *Spine (Phila Pa 1976)*2015;40:E83-90.
26. Smeets, R, Koke, A, Lin, CW et al. Measures of function in low back pain/disorders: Low Back Pain Rating Scale (LBPRS), Oswestry Disability Index (ODI), Progressive Isoinertial Lifting Evaluation (PILE), Quebec Back Pain Disability Scale (QBPDS), and Roland-Morris Disability Questionnaire (RDQ). *Arthritis Care Res (Hoboken)*2011;63 Suppl 11:S158-73.



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46
27. Chapman, JR, Norvell, DC, Hermsmeyer, JT et al. Evaluating common outcomes for measuring treatment success for chronic low back pain. *Spine (Phila Pa 1976)*2011;36:S54-68.
  28. Lind, MY, Hop, WC, Weimar, W et al. Body image after laparoscopic or open donor nephrectomy. *Surg Endosc*2004;18:1276-9.
  29. Dunker, MS, Stiggelbout, AM, van Hogezaand, RA et al. Cosmesis and body image after laparoscopic-assisted and open ileocolic resection for Crohn's disease. *Surg Endosc*1998;12:1334-40.
  30. Arts, MP, Brand, R, van den Akker, ME et al. Tubular discectomy vs conventional microdiscectomy for sciatica: a randomized controlled trial. *JAMA*2009;302:149-58.
  31. van den Akker, ME, Arts, MP, van den Hout, WB et al. Tubular discectomy vs conventional microdiscectomy for the treatment of lumbar disk-related sciatica: cost utility analysis alongside a double-blind randomized controlled trial. *Neurosurgery*2011;69:829-35; discussion 835-6.
  32. Herdman, M, Gudex, C, Lloyd, A et al. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res*2011;20:1727-36.
  33. Versteegh, MM, Vermeulen, KM, Evers, SMAA et al. Dutch Tariff for the Five-Level Version of EQ-5D. *Value in Health*2016;19:343-352.
  34. Hakkaart-Van Roijen, L, van der Linden, N, Bouwmans, C et al. Kostenhandleiding: Methodologie van kostenonderzoek en referentieprijzen voor economische evaluaties in de gezondheidszorg [Manual for economic evaluations. Methods and standard cost prices for economic evaluations in health care]. Diemen, The Netherlands: Health Care Institute (ZiN)2016. <https://www.zorginstituutnederland.nl/over-ons/publicaties/publicatie/2016/02/29/richtlijn-voor-het-uitvoeren-van-economische-evaluaties-in-de-gezondheidszorg> (date assessed June 2017).
  35. Z-Index. G-Standaard. The Hague, The Netherlands: Z-Index BV.2009. <https://www.z-index.nl/g-standaard> (date assessed January 2017).
  36. White, IR, Royston, P, and Wood, AM. Multiple imputation using chained equations: Issues and guidance for practice. *Statistics in Medicine*2011;30:377-399.
  37. Gomes, M, Grieve, R, Nixon, R et al. Methods for Covariate Adjustment in Cost-Effectiveness Analysis That Use Cluster Randomised Trials. *Health Economics*2012;21:1101-1118.
  38. Drummond, M and Drummond, M, *Methods for the economic evaluation of health care programmes*. 3rd ed. Oxford medical publications. 2005, Oxford ; New York: Oxford University Press. 379 p.
  39. Black, WC. The CE plane: a graphic representation of cost-effectiveness. *Med Decis Making*1990;10:212-4.
  40. Fenwick, E, O'Brien, BJ, and Briggs, A. Cost-effectiveness acceptability curves--facts, fallacies and frequently asked questions. *Health Econ*2004;13:405-15.
  41. Peul, WC, van Houwelingen, HC, van den Hout, WB et al. Surgery versus prolonged conservative treatment for sciatica. *N Engl J Med*2007;356:2245-56.
  42. Breivik, H, Borchgrevink, PC, Allen, SM et al. Assessment of pain. *Br J Anaesth*2008;101:17-24.

## FOOTNOTES

### Contributorship statement:

SH, JvS, WP, MvT, SR contributed to writing the proposal and acquiring the grant from The Netherlands Organisation for Health and Care Research (ZonMw). AS prepared the first draft of the study protocol. All authors (AS, PG, SH, JvS, WP, MvT, MdB, SR) contributed to the final design of this study protocol, assisted with drafting the manuscript, and approved the final version of the manuscript. MdB developed the statistical analysis plan.

**Funding statement:**

This study was funded by ZonMw, The Netherlands Organisation for Health Research and Development [project number 837004013]. The funder did not play any role in the study design. This research received no specific grant from any funding agency in the public, commercial or not-profit sectors.

**Competing interest statement:**

The professional judgment concerning the primary interest (e.g. validity of the research) will not be influenced by a secondary interest. All authors have read and understood the BMJ policy on declaration of interests. We all declare that we have no competing interests other than receiving grants. For this particular study we had financial support from The Netherlands Organisation for Health Research and Development. We do not have relationships with any organisations that might have an interest in the submitted work. All authors have completed the ICJE uniform at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf).

**Provenance and peer review:** Not commissioned; externally peer reviewed.

**Data sharing statement:**

This article refers to a study protocol; data are not yet available. We will share all data through open access once the final results have been published.

**Ethical approval:** This study has received approval of the Medical Ethical Committee of the VU Medical Centre Amsterdam and confirmation can be supplied upon request [corresponding number: NL50951.029.14; November 5, 2015].



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents\*

Section/item	Item No	Description	Addressed on page number
<b>Administrative information</b>			
Title	1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	___ 1 ___
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	___ 2 ___
	2b	All items from the World Health Organization Trial Registration Data Set	___ 2 ___
Protocol version	3	Date and version identifier	___ 4 ___
Funding	4	Sources and types of financial, material, and other support	___ 15 ___
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	___ 1 and 15 ___
	5b	Name and contact information for the trial sponsor	___ 1 ___
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	___ 15 ___
	5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	___ 13 en 15 ___



1  
2  
3 **Introduction**  
4

5 Background and	6a	Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention	_____ 2 and 3 _____
6 rationale			
7			
8	6b	Explanation for choice of comparators	_____ 2 and 3 _____
9			
10 Objectives	7	Specific objectives or hypotheses	_____ 3 _____
11			
12 Trial design	8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory)	_____ 3 and 4 _____ (see also design section)_____
13			
14			
15			
16			
17			

18 **Methods: Participants, interventions, and outcomes**  
19

20 Study setting	9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will be collected. Reference to where list of study sites can be obtained	_____ 4 _____
21			
22			
23 Eligibility criteria	10	Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and individuals who will perform the interventions (eg, surgeons, psychotherapists)	_____ 4 _____
24			
25			
26 Interventions	11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be administered	_____ 5 and 6 _____
27			
28			
29	11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose change in response to harms, participant request, or improving/worsening disease)	_____ 6 en 9 _____
30			
31			
32	11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence (eg, drug tablet return, laboratory tests)	_____ n.a. _____
33			
34			
35	11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	_____ 6 _____
36			
37 Outcomes	12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen efficacy and harm outcomes is strongly recommended	_____ 6-8 _____
38			
39			
40			
41			
42			
43			
44			
45			
46			
47			
48			
49			

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Participant timeline	13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure)	_____8_____
Sample size	14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	_____9_____
Recruitment	15	Strategies for achieving adequate participant enrolment to reach target sample size	_____9_____

**Methods: Assignment of interventions (for controlled trials)**

Allocation:

Sequence generation	16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	_____5_____
Allocation concealment mechanism	16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	_____5_____
Implementation	16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	_____5_____
Blinding (masking)	17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	_____5_____
	17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	_____N.A._____

**Methods: Data collection, management, and analysis**

Data collection methods	18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	_____6-11_____
-------------------------	-----	--	----------------

1			
2			
3		18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols
4			_____6_____
5			
6	Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol
7			_____9_____
8			
9			
10	Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol
11			_____9-12_____
12			
13		20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)
14			__10 and 12__
15		20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)
16			_____9-10_____
17			
18			
19	<b>Methods: Monitoring</b>		
20			
21	Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed
22			_ 9, 12, 15 and 16__
23			
24			
25			
26		21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial
27			_____9_____
28			
29	Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct
30			__9,12 and 16__
31			
32	Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor
33			_____12_____
34			
35			
36	<b>Ethics and dissemination</b>		
37			
38	Research ethics approval	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval
39			_____12 and 15_____
40			
41			
42			
43			
44			
45			
46			
47			
48			
49			

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Protocol amendments	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial registries, journals, regulators)	4
Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	4
	26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	N.A.
Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	9
Declaration of interests	28	Financial and other competing interests for principal investigators for the overall trial and each study site	15 en 16
Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	15 and 6
Ancillary and post-trial care	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	12
Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	12
	31b	Authorship eligibility guidelines and any intended use of professional writers	15
	31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	n.a.
<b>Appendices</b>			
Informed consent materials	32	Model consent form and other related documentation given to participants and authorised surrogates	appendix of bmj open online submission system

1  
2  
3 Biological 33 Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular \_\_\_\_\_ N.A. \_\_\_\_\_  
4 specimens analysis in the current trial and for future use in ancillary studies, if applicable  
5

---

6 \*It is strongly recommended that this checklist be read in conjunction with the SPIRIT 2013 Explanation & Elaboration for important clarification on the items.  
7 Amendments to the protocol should be tracked and dated. The SPIRIT checklist is copyrighted by the SPIRIT Group under the Creative Commons  
8 "[Attribution-NonCommercial-NoDerivs 3.0 Unported](https://creativecommons.org/licenses/by-nc-nd/3.0/)" license.  
9

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>