## Supplemental Table 1: Crude and adjusted estimates of effect of intimate partner violence and alcohol use on SGA and preterm births. (exposures measured during third trimester)

	Small for gestational age					Preterm birth	
	No of pregnancy	No of SGA	Risk Ratio [95% CI]		No of	Risk Ratio [95% CI]	
			crude	Adjusted <sup>1</sup>	preterm	Crude	Adjusted <sup>1</sup>
Intimate partne	r violence						
No violence	531 21	63	1.00 [Reference]	1.00 [Reference]	70 2	1.00 [Reference]	1.00 [Reference]
Physical	21	1	0.39 [0.05, 2.93]	0.39 [0.05, 3.30]	2	0.69 [0.16, 3.03]	0.72 [0.15, 3.31]
Sexual	1	0	*	*	0	*	*
Psychological	27	2	0.59 [0.14, 2.54]	0.58 [0.12, 2.75]	2	0.52 [0.12, 2.24]	0.49 [0.11, 2.24]
Alcohol use							
Low	517	57	1.00 [Reference]	1.00 [Reference]	67	1.00 [Reference]	1.00 [Reference]
Hazardous drinking in women	54	9	1.55 [0.72, 3.33]	1.62 [0.68, 3.84]	8	1.16 [0.52, 2.57]	1.13 [0.48, 2.65]