

Supplementary File 3: Unweighted descriptive statistics for outcome variables by dosage group

Outcome	Group	Baseline	24-month
Ecstasy intentions, % (n)	FULL	1.06 (4)	3.88 (9)
	INCOMPLETE	3.88 (10) ^b	8.18 (13)
	CONTROL	0.82 (4)	3.48 (10)
NPS intentions, % (n)	FULL	1.32 (5)	1.73 (4)
	INCOMPLETE	2.33 (6)	5.66 (9)
	CONTROL	0.82 (4)	2.79 (8)
Synthetic cannabis intentions, % (n)	FULL	5.82 (22)	6.03 (14)
	INCOMPLETE	8.53 (22) ^b	9.43 (15)
	CONTROL	4.5 (22)	8.7 (25)
Synthetic stimulants intentions, % (n)	FULL	1.9 (7)	1.72 (4)
	INCOMPLETE	2.7 (7) ^b	5.03 (8)
	CONTROL	0.8 (4)	1.05 (3)
Ever used ecstasy, % (n)	FULL	1.59 (6)	3.88 (9)
	INCOMPLETE	4.25 (11) ^b	12.35 (20)
	CONTROL	1.23 (6)	2.7 (8)
Ever used NPS, % (n)	FULL	1.06 (4)	2.16 (5)
	INCOMPLETE	6.18 (16) ^b	8.07 (13)
	CONTROL	2.86 (14)	2.44 (7)
Ever used synthetic cannabis, % (n)	FULL	0.79 (3)	1.72 (4)
	INCOMPLETE	4.63 (12)	8.02 (13)
	CONTROL	2.45 (12)	1.38 (4)
Ever used synthetic stimulants, % (n)	FULL	0.00 (0)	1.69 (4)
	INCOMPLETE	2.06 (5) ^b	4.88 (8)
	CONTROL	0.0 (0)	1.02 (3)
Ecstasy knowledge (M, SD)	FULL	10.51 (0.13) ^a	11.63 (.22)
	INCOMPLETE	10.67 (0.16) ^b	10.59 (.28)
	CONTROL	9.18 (2.87)	9.99 (.21)
NPS knowledge (M, SD)	FULL	3.33 (.07) ^a	3.92 (.09)
	INCOMPLETE	3.32 (.08) ^b	3.40 (.12)
	CONTROL	2.71 (.06)	3.07 (.09)

^a Significant baseline difference full dose vs. control

^b Significant baseline difference incomplete dose vs. control