

BMJ Open Smoking behaviours of Hong Kong Chinese hospitalised patients and predictors of smoking abstinence after discharge: a cross-sectional study

Ka Yan Ho,¹ William Ho Cheung Li,¹ Katherine Ka Wai Lam,¹ Man Ping Wang,¹ Wei Xia,¹ Lok Yin Ho,¹ Kathryn Choon Beng Tan,² Hubert Kit Man Sin,³ Elaine Cheung,⁴ Maysi Pik Hung Mok,⁴ Tai Hing Lam⁵

To cite: Ho KY, Li WHC, Lam KKW, *et al.* Smoking behaviours of Hong Kong Chinese hospitalised patients and predictors of smoking abstinence after discharge: a cross-sectional study. *BMJ Open* 2018;**8**:e023965. doi:10.1136/bmjopen-2018-023965

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-023965>).

Received 3 May 2018

Revised 11 September 2018

Accepted 19 October 2018



© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹School of Nursing, The University of Hong Kong, Hong Kong, China

²Department of Medicine, The University of Hong Kong, Hong Kong, China

³Department of Medicine and Geriatrics, Tuen Mun Hospital, Hong Kong, China

⁴Department of Medicine and Geriatrics, United Christian Hospital, Hong Kong, China

⁵School of Public Health, The University of Hong Kong, Hong Kong, China

Correspondence to

Dr William Ho Cheung Li; william3@hku.hk

ABSTRACT

Objectives Patients admitted to hospitals represent an excellent teachable moment for smoking cessation, as they are required to abstain from tobacco use during hospitalisation. Nevertheless, smoking behaviours of hospitalised patients, and factors that lead to smoking abstinence thereafter, remain relatively underexplored, particularly in a Hong Kong Chinese context. This study aimed to examine the smoking behaviours of hospitalised patients and explore factors leading to their abstaining from cigarette use after being hospitalised.

Design A cross-sectional design was employed.

Setting This study was conducted in three outpatient clinics in different regions in Hong Kong.

Participants A total of 382 recruited Chinese patients.

Primary and secondary outcome measures The patients were asked to complete a structured questionnaire that assessed their smoking behaviours before, during and after hospitalisation.

Results The results indicated 23.6% of smokers smoked secretly during their hospital stay, and about 76.1% of smokers resumed smoking after discharge. Multivariate logistic regression analysis found that number of days of hospitalisation admission in the preceding year (OR 1.02; 95% CI 1.01 to 1.27; $p=0.036$), patients' perceived correlation between smoking and their illness (OR 1.08; 95% CI 1.01 to 1.17; $p=0.032$), withdrawal symptoms experienced during hospitalisation (OR 0.75; 95% CI 0.58 to 0.97; $p=0.027$) and smoking cessation support from healthcare professionals (OR 1.18; 95% CI 1.07 to 1.36; $p=0.014$) were significant predictors of smoking abstinence after discharge.

Conclusions The results of this study will aid development of appropriate and innovative smoking cessation interventions that can help patients achieve more successful smoking abstinence and less relapse.

Trial registration number NCT02866760.

INTRODUCTION

Cigarette smoking, responsible for around 7 million deaths annually worldwide, is the single greatest preventable cause of death.¹ It harms nearly every organ in the body and is

Strengths and limitations of this study

- This is the first study to investigate the smoking behaviours of hospitalised patients in a Chinese context.
- This is the first study to identify contributing factors that lead to smoking abstinence after patients are discharged.
- Biochemical validation was not conducted to verify self-reported abstinence; therefore, results could be biased by social desirability.
- Participants were asked to provide details on their smoking behaviour during their hospital stay and after discharge, which may have resulted in recall bias.

associated with numerous diseases, including stroke, diabetes, cancer, coronary heart disease and respiratory disease,² all of which contribute to substantial amounts of hospitalisation and healthcare expenditure, posing a serious challenge to medical systems.³ Although the prevalence of daily cigarette smoking in Hong Kong has decreased from 23.3% in 1982 to 10.5% in 2015, 641 300 everyday smokers remain,⁴ and 400 000 hospitalisations per year are attributable to smoking.⁵ Such compelling numbers cannot be overlooked or neglected.

Hospitalisation represents an excellent teachable moment for smoking cessation.⁶ This is because being hospitalised with a smoking-related disease may impel change in smokers' perceptions of their personal vulnerability, which in turn can greatly enhance their motivation to quit.⁷ Furthermore, hospitalised smokers may have more available time to receive intensive smoking cessation interventions, which can remarkably increase their chances of successful abstinence after discharge.⁸ Additionally, a smoke-free policy

has been implemented in all public hospitals in Hong Kong since 2007. Violating the policy results in a fixed fine of HK\$1500 (~US\$192).⁹ Implementation of such a policy may promote smoking cessation, as it prompts smokers, on hospital admission, to undergo an abrupt stop from their habitual use of tobacco. It also creates a smoke-free environment that temporarily facilitates abstinence by separating hospitalised smokers from smoking cues.¹⁰

Given that habitual smoking in patients who have diseases can reduce the efficacy of clinical and medical treatments, and increase the risk of treatment-related side effects,¹¹ and that smoking abstinence is enforced during hospitalisation, it is of paramount importance for healthcare professionals to seize this golden opportunity to promote cessation and help patients quit. The clinical guidelines for treating tobacco use and nicotine dependence also emphasise the importance of targeting smoking cessation interventions at hospitalised smokers.¹² Nevertheless, cigarette smoking is addictive, quitting is extremely difficult and the rate of relapse is high, especially among patients with diseases.¹³ It is therefore crucial that healthcare professionals first understand how hospitalised smokers perceive the risks of smoking, and their behaviour towards smoking, before any effective, appropriate smoking cessation intervention can be planned, developed and evaluated.

Numerous studies in the West have investigated smoking behaviours of hospitalised patients.^{10 14 15} In a secondary analysis of data from a randomised control trial on 650 adult smokers admitted to an urban teaching hospital, 4% self-reported violating policy by smoking indoors.¹⁰ Another cross-sectional study of 229 hospitalised smokers found 60%–70% complied with the smoke-free policy.¹⁴ A study of 79 hospitalised smokers revealed about 75% intended to quit after discharge.¹⁵ However, the majority perceived their symptoms associated with nicotine cravings would be a primary barrier towards maintaining abstinence.¹⁵ Indeed, a study revealed smokers with nicotine cravings were more likely to smoke while hospitalised than those who did not report cravings.¹⁰ Additionally, withdrawal symptoms and patients' perceived correlation between smoking and their illness were found to be factors significantly associated with compliance with the smoke-free policy during hospitalisation.¹⁴

There is scarce evidence on key factors that lead to post-hospitalisation smoking abstinence. While some studies have examined smoking behaviours of hospitalised patients, they primarily focused on understanding patients' compliance with smoke-free policies.^{14 15} One study explored predictors of continued abstinence after discharge, though it was a secondary analysis of data from a randomised controlled trial in which hospitalised smokers had already received some degree of intensive counselling on smoking cessation; that might confound the results, especially those regarding smoking behaviours after discharge.¹⁰ The present study aimed to address the gap in existing literature by examining the

smoking behaviours of hospitalised patients in hospitals with smoke-free policies. It also sought to identify factors leading to smoking abstinence after discharge.

METHODS

Study design

A cross-sectional study was conducted in outpatient clinics of three acute hospitals in different regions in Hong Kong. All participating hospitals had been awarded full accreditation status by the Australian Council on Healthcare Standards.¹⁶ These three hospitals were chosen because they are the largest acute regional hospitals and patients were normally referred back for medical follow-up in the outpatient clinics after discharge.

Participants

Patients were eligible for this study, if they were (1) aged ≥ 18 years, (2) able to speak Cantonese, (3) current smokers who resumed smoking or ex-smokers who had quit smoking after hospitalisation, and (4) hospitalised in either a medical or surgical unit for ≥ 48 hours in the previous 3 months. 'Ex-smokers' in this study referred to patients who reported not having smoked in the preceding 7 days. We excluded patients with mental illnesses or cognitive and learning problems noted in their medical records because previous studies suggest such people possess smoking characteristics different from those of smokers in general.¹⁷ Including them in the study might therefore bias the results, impeding generalisability to all hospitalised smokers.

The sample size was calculated based on the results of previous literature that indicated 46%–56.8% of smokers were willing to quit during hospitalisation.^{2 18} We assumed a similar proportion of our participants were willing and adopted a 95% confidence level. The sample size could be obtained using the following formula:

$$N = [Z^2 \times p(1 - p)] / e^2$$

where Z represents the number of SD from the mean, which is 1.96; p refers to the expected proportion, which is 0.46 in this case; and e is the margin of error, which is 0.05. The sample size required for the study was therefore 382.

Measure

Smoking behaviours

An expert panel developed a structured questionnaire to explore smoking behaviours during and after hospitalisation. The panel included a chair professor, associate professor, assistant professor and research assistant professor from a local university, all of whom had extensive knowledge on smoking cessation. The questionnaire collected data on experience of withdrawal symptoms and nicotine cravings during and after hospitalisation. Withdrawal symptoms were assessed using the Minnesota Nicotine Withdrawal Scale (MNWS). This scale includes 15 items evaluated on a 5-point Likert scale (0, not present; 1, slight; 2, mild; 3, moderate; 4, severe). The possible

range of scores was 0–60, with higher scores indicating more severe withdrawal symptoms.¹⁹ Intensity of physical addiction to nicotine was assessed using the Fagerström Test for Nicotine Dependence (FTND). This test contains six items, with scores of 0–3, 4–5 and 6–10, representing mild, moderate and high levels of nicotine dependence, respectively.²⁰ The questionnaire also covered four other main areas: (1) whether participants received smoking cessation advice from healthcare professionals during hospitalisation; (2) smoking and quitting behaviours before, during and after hospitalisation; (3) intention to quit; (4) risk perception towards smoking by a binary scale (ie, yes or no). All questions in the questionnaire were written in Chinese.

Sociodemographic and clinical characteristics

A sociodemographic sheet was administered to collect participants' background information, including their age, sex, educational attainment and marital status. A research assistant also accessed the participants' medical records and extracted number of admissions in the previous year, diagnosis of the most recent admission, comorbidities and length of stay of the most recent admission.

Data collection

Data were collected from March to August 2017. Convenience sampling was used. A research assistant approached patients who had a medical follow-up in the outpatient clinics, and clearly explained the study in detail. The patients were told their participation was completely voluntary, and refusal to join or answer would in no way influence the care they received. Those who were eligible and willing to participate were asked to provide written consent and complete the questionnaire with a demographic sheet. The entire consent process took 10–15 min, causing only minimal disturbance to the clinical routine.

Data analysis

Data analysis was performed using IBM SPSS Statistics for Windows, V.23.0 (IBM). Descriptive statistics were used to detail participants' sociodemographic and clinical characteristics, and their smoking profiles. Continuous variables were presented in either mean with SD or median with IQR, while categorical variables were presented in frequency and proportion. Multivariate logistic regression analysis was conducted to identify predictors of post-hospitalisation smoking abstinence, with smokers as the reference group. The selection of variables in multivariate analysis was based on a theory-driven approach. Despite there was a little understanding of smoking behaviours of hospitalised patients in the Hong Kong Chinese context, variables that were potentially associated with quitting according to previous literature were entered into the regression analyses to build the model.^{10 14 15}

Table 1 Demographic, socioeconomic and clinical characteristics of participants (n=382)

Sex, n (%)	
Male	370 (96.9)
Female	12 (3.1)
Age, mean (SD)	
	54.7 (12.7)
Marital status, n (%)	
Married	260 (68.0)
Unmarried	60 (15.7)
Divorced, widowed, separated	49 (12.8)
Missing	13 (3.5)
Highest educational attainment, n (%)	
Primary school or below	99 (25.9)
Lower secondary school	144 (37.7)
Upper secondary school	104 (27.2)
Tertiary education	26 (6.8)
Missing	9 (2.4)
Employment status, n (%)	
Employed	204 (53.4)
Unemployed	62 (16.2)
Retired	110 (28.8)
Missing	6 (1.6)
Reason for hospital admission, n (%)	
Diabetes	100 (26.2)
Hypertension	156 (40.8)
Heart disease	84 (22.0)
Respiratory disease	22 (5.8)
Cancer	14 (3.6)
Missing	6 (1.6)
Length of stay in hospital (days), median (IQR)	
	5.0 (2.0–10.0)
No of admissions in the preceding year, median (IQR)	
	1.0 (1.0–2.0)

Patient and public involvement

Patients and public were not included in the development, design, implementation or dissemination of the research. The results would be disseminated to the participants through patient forums.

RESULTS

A total of 13848 patients were screened in the three outpatient clinics; of them, 468 were eligible and 382 subsequently agreed to participate in this study.

Table 1 shows the mean age of participants was 54.7 (SD 12.7) years. About 96.9% (370/382) were men, 68.0% (260/382) were married, 37.7% (144/382) had received lower secondary education and 53.4% (204/382) were employed. Concerning clinical characteristics, participants were admitted to hospitals for 1.0

Table 2 Smoking profiles of participants before and during hospitalisation (n=382)

Before hospitalisation	
Daily cigarette consumption, median (IQR)	20.0 (10.0–20.0)
Previous quit attempt(s), n (%)	
Yes	216 (56.5)
No	166 (43.5)
Previous reduction attempt(s), n (%)	
Yes	151 (39.5)
No	231 (60.5)
Nicotine dependency by Fagerström Test for Nicotine Dependence, n (%)	
Low	165 (43.2)
Moderate	102 (26.7)
High	115 (30.1)
During hospitalisation	
Complied with smoke-free policy, n (%)	
Yes	292 (76.4)
No	90 (23.6)
Withdrawal symptoms by Minnesota Nicotine Withdrawal Scale, mean (SD)	4.2 (5.4)
Made committed attempt(s) to reduce cigarette consumption*, n (%)	
Yes	77 (85.7)
No	13 (14.3)
Daily cigarette consumption*, mean (SD)	1.3 (5.0)
Received smoking cessation support from healthcare professionals, n (%)	
Yes	254 (66.5)
No	128 (33.5)

*Calculation based on participants who smoked during hospitalisation (n=90).

(IQR 1.0–2.0) times on average in the preceding year. For the most recent admission, 40.8% (156/382) were for hypertension (156/382), followed by 26.2% (100/382) for diabetes, 22.0% (84/382) for heart disease, 5.8% for respiratory disease and 3.6% (14/382) for cancer. The average length of period of hospitalisation was 5.0 (IQR 2.0–10.0) days.

Table 2 shows participants smoked 20.0 (IQR 10.0–20.0) cigarettes per day on average before hospitalisation. Approximately 43.5% (166/382), 26.7% (102/382) and 30.1% (115/382) had mild, moderate or high nicotine dependency, respectively, as gauged by the FTND.

Around 56.5% (216/382) and 39.5% (151/382) had previously attempted to quit or reduce smoking, respectively. Regarding smoking behaviours in hospitals, 76.4% (292/382) of the participants complied with the smoke-free policy, while 23.6% (90/382) admitted to having smoked. Of those who continued smoking during

Table 3 Smoking profile of participants after hospitalisation (n=382)

Quit smoking	
Yes	93 (24.3)
No	289 (75.7)
Daily cigarette consumption after hospitalisation*, mean (SD)	10.1 (11.4)
Intended to quit smoking after hospitalisation*, n (%)	
Yes	143 (49.5)
No	146 (50.5)

*Calculation based on participants who were smoking after discharge (n=289).

hospitalisation, 85.7% (77/90) reduced their smoking amount, with mean daily cigarette consumption of 1.3 (SD 5.0). The mean score of withdrawal symptoms as gauged by the MNWS was 4.2 (SD 5.4). Additionally, 66.5% (254/382) of the participants received smoking cessation support from healthcare professionals.

Table 3 shows the smoking behaviours of participants after hospital discharge. About 24.3% (93/382) had successfully quit smoking. Among those who did not, 49.5% (143/289) reported intention to quit; their mean daily cigarette consumption was 11.6 (SD 11.5).

Predictors of smoking abstinence following hospitalisation

Table 4 shows the results of univariate logistic regression analysis. Married (OR 2.46; 95% CI 1.03 to 5.87; p=0.043), retired (OR 2.08; 95% CI 1.06 to 4.00; p=0.032), daily cigarette consumption before hospitalisation (OR 0.98; 95% CI 0.97 to 1.00; p=0.036), number of days of hospitalisation (OR 1.03; 95% CI 1.00 to 1.05; p=0.040), number of hospital admissions in the preceding year (OR 1.02; 95% CI 1.01 to 1.03; p=0.005), diabetes (OR 0.46; 95% CI 0.21 to 0.99; p=0.046), hypertension (OR 0.52; 95% CI 0.32 to 0.87; p=0.012), patients' perceived correlation between smoking and their illness (OR 1.11; 95% CI 1.04 to 1.18; p=0.002), withdrawal symptoms experienced during hospitalisation (OR 0.72; 95% CI 0.58 to 0.89; p=0.003) and whether patients received smoking cessation support from healthcare professional (OR 1.19; 95% CI 1.07 to 1.33; p<0.001) were shown to be statistically significantly associated with post-discharge smoking abstinence. **Table 5** presents the results of that multivariate logistic regression analysis. Participants' sociodemographic and smoking characteristics—including age, sex, educational attainment, and marital and employment status—were controlled. Only number of days of hospitalisation admission in the preceding year (OR 1.02; 95% CI 1.01 to 1.27; p=0.036), patients' perceived correlation between smoking and their illness (OR 1.08; 95% CI 1.01 to 1.17; p=0.032), withdrawal symptoms experienced during hospitalisation (OR 0.75; 95% CI 0.58 to 0.97; p=0.027) and whether patients received smoking cessation support from healthcare professional during

Table 4 Univariate logistic regression analyses on predictors of smoking abstinence after discharge (n=382)

Unadjusted model		
	OR (95% CI)	P value
Age	0.99 (0.96 to 1.02)	0.596
Sex	0.28 (0.03 to 2.34)	0.239
Highest educational attainment		
Primary school or below	0.51 (0.19 to 1.38)	0.184
Lower secondary school	0.45 (0.17 to 1.19)	0.107
Upper secondary school	0.48 (0.18 to 1.29)	0.147
Tertiary education	Ref	
Marital status		
Married	2.46 (1.03 to 5.87)	0.043*
Unmarried	0.89 (0.27 to 2.91)	0.842
Divorced, widowed, separated	Ref	
Employment status		
Employed	Ref	
Awaiting employment/unemployed	0.97 (0.48 to 1.94)	0.920
Retired	2.08 (1.06 to 4.00)	0.032*
Daily cigarette consumption before hospitalisation	0.98 (0.97 to 1.00)	0.036*
Quit attempts before hospitalisation	0.63 (0.34 to 1.20)	0.162
Reduction attempts before hospitalisation	0.83 (0.34 to 2.01)	0.828
No of days of hospitalisation	1.03 (1.00 to 1.05)	0.040*
No of hospital admissions in the preceding year	1.02 (1.01 to 1.03)	0.005*
Reason for hospital admission		
Diabetes	0.46 (0.21 to 0.99)	0.046*
Hypertension	0.52 (0.32 to 0.87)	0.012*
Heart disease	0.86 (0.50 to 1.49)	0.600
Respiratory disease	0.81 (0.25 to 2.64)	0.730
Cancer	Ref	
Perceived correlation between smoking and illness	1.11 (1.04 to 1.18)	0.002*
Received smoking cessation support from healthcare professionals	1.19 (1.07 to 1.33)	<0.001**
Withdrawal symptoms experienced during hospitalisation	0.72 (0.58 to 0.89)	0.003*

*Significant at $p < 0.05$; **significant at $p < 0.001$.

hospitalisation (OR 1.18; 95% CI 1.07 to 1.36; $p = 0.014$) were continuously found to be significant predictors of post-discharge smoking abstinence.

DISCUSSION

To the best of our knowledge, this is the first study to investigate the smoking behaviours of hospitalised patients in a Chinese context. It is also the first study to identify contributing factors that lead to smoking abstinence after patients are discharged. In this study, almost

Table 5 Multivariate logistic regression analyses on predictors of smoking abstinence after discharge (n=382)

Adjusted model		
	OR (95% CI)	P value
Age	0.99 (0.97 to 1.02)	0.701
Sex	0.30 (0.03 to 2.69)	0.282
Highest educational attainment		
Primary school or below	0.53 (0.17 to 1.52)	0.225
Lower secondary school	0.49 (0.17 to 1.39)	0.177
Upper secondary school	0.49 (0.15 to 1.37)	0.162
Tertiary education	Ref	
Marital status		
Married	2.12 (0.88 to 5.82)	0.092
Unmarried	0.92 (0.28 to 3.03)	0.894
Divorced, widowed, separated	Ref	
Employment status		
Employed	Ref	
Awaiting employment/unemployed	0.89 (0.40 to 1.97)	0.772
Retired	1.85 (0.92 to 3.71)	0.084
Daily cigarette consumption before hospitalisation	0.98 (0.96 to 1.00)	0.050
Quit attempts before hospitalisation	0.70 (0.41 to 1.21)	0.202
Reduction attempts before hospitalisation	0.81 (0.49 to 1.51)	0.603
No of days of hospitalisation admission in the preceding year	1.02 (1.01 to 1.27)	0.036*
Reason for hospital admission		
Diabetes	0.51 (0.24 to 1.81)	0.297
Hypertension	0.55 (0.36 to 1.37)	0.147
Heart disease	0.65 (0.18 to 2.32)	0.507
Respiratory disease	0.52 (0.11 to 2.41)	0.400
Cancer	Ref	
Perceived correlation between smoking and illness	1.08 (1.01 to 1.17)	0.032*
Received smoking cessation support from healthcare professionals	1.18 (1.07 to 1.36)	0.014*
Withdrawal symptoms experienced during hospitalisation	0.75 (0.58 to 0.97)	0.027*

The model was adjusted by sex, age, highest educational attainment, marital status and employment status.

*Significant at $p < 0.05$.

all the participants were men. This indeed reflected a usual phenomenon in Hong Kong. According to the latest statistics from the Census and Statistics Department, Hong Kong, about 86% of daily cigarette smokers were men.²¹

The results show most participants complied with the hospital's policy to stop smoking; however, 23.6% did not, and smoked secretly during hospitalisation. Most had either moderate or high nicotine dependency before admission, which aggravated their ability to abstain, despite the hospital's policy.

The study showed 23.9% of the participants quit smoking after being discharged. Multivariate logistic regression analyses were conducted to aid in understanding the contributing factors leading to smoking abstinence of the patients after hospitalisation. The results revealed patients' perception of the relationship between smoking and disease was one of the most significant predictors of smoking abstinence. This finding is in line with that in a previous study showing that medically ill smokers who regarded themselves as more vulnerable to health consequences of smoking were more likely to quit.⁷ In addition, the number of hospital admissions in the preceding year was found to be a significant predictor of smoking abstinence after hospitalisation. This was probably due to the fact that smokers were required to temporarily abstain in hospitals, which provided a golden opportunity to initiate smoking cessation. However, as observed in the regression models, reasons for hospital admission could not be used to predict post-hospitalisation smoking abstinence. A possible explanation is that a majority of the participants had misconceptions and believed smoking was not a cause of their disease, and they resumed smoking after discharge. However, evidence shows smoking poses serious adverse effects to nearly every organ in the body.² Quitting smoking can also help slow disease progression²² and improve treatment efficacy.² In this regard, dispelling misconceptions about smoking and increasing smokers' perception of the risks of continued smoking, and the benefits of quitting, is a potential strategy for promoting smoking cessation for hospitalised smokers.

The multivariate analysis results revealed that withdrawal symptoms experienced during hospitalisation serve as a significant barrier to continued abstinence thereafter. These results are consistent with previous studies indicating withdrawal symptoms induced by nicotine dependence are a common reason for smokers continue to smoking after discharge.²³ Rigotti *et al*¹⁰ suggested withdrawal symptoms are less prominent in hospital settings, as smoking was completely banned there, resulting in a lack of environmental cues. However, it was expected that smoking patients may suffer more severe withdrawal symptoms when they were discharged and returned to an environment with different smoking cues. Therefore, equipping smokers with the skills to overcome cravings and withdrawal symptoms during hospitalisation is another crucial strategy for assisting them in continuing abstinence after discharge.

The multivariate logistic regression analyses indicated patients' receiving smoking cessation support during hospitalisation was a significant predictor of smoking abstinence. This finding further supports the WHO recommendation that healthcare professionals bear a

continued responsibility for promoting smoking cessation, as their advice can strongly affect motivating of hospitalised smokers to quit.²⁴ Nevertheless, inadequate cessation support of hospitalised smokers by healthcare professionals in Hong Kong was observed. About 30% of participants in the present study received no support for smoking cessation during their hospitalisation. This could owe to the fact many healthcare professionals in Hong Kong have never received formal training on smoking cessation and therefore lacked competence to provide adequate support or concrete methods for hospitalised smokers to quit smoking. Another potential reason is the problematic healthcare professional shortage highly prevalent in Hong Kong. The nurse:patient ratio in most public hospitals is 1:12, which is far below the international standard of 1:6 for developed economies.²⁵ Providing comprehensive smoking cessation advice for hospitalised smokers therefore appears infeasible in this study setting that has inadequate healthcare human resources.

Limitations

This study did have some limitations. First, we did not conduct biochemical validation to verify self-reported abstinence. This was because the participating hospitals implemented strict infection control policies. When we prepared this study, we discussed with our clinical partners to see whether we could conduct such validation. However, they expressed concern about the possibility of spreading infection in their facilities, especially when the procedures involved patients' saliva. Despite our suggestion to verify patients' smoking status by means of exhaled carbon monoxide, the necessary equipment needed to be shared, which might also have presented chances of infection. As biochemical validation was not conducted to verify self-reported abstinence, therefore, results could be biased by social desirability. Notably, there was potential for under-reporting of smoking during hospitalisation and over-reporting of abstinence after discharge. Biochemical validation was suggested for future research to minimise such bias. The participants were also asked to provide details on their smoking behaviours during their hospital stay and after discharge, which may have resulted in recall bias. Additionally, the study sample size was small compared with those from previous studies on hospitalised smokers,^{26 27} which in turn might have lowered the power to predict abstinence. We also were unable to obtain any data for comparison between our participants and non-respondents with regard to their demographics and clinical and smoking characteristics, thus working against representativeness among participants. Furthermore, because of the length of the questionnaire, we were not able to include other correlates of quitting, including attitudes towards tobacco use. Finally, although the participating hospitals had similar attributes and settings, certain potentially unobservable differences may have made our results less clear.

Implications for clinical practice and research

Despite the above limitations, the present findings have important implications for research and clinical practice. As provision of smoking cessation support is inadequate in clinical settings, hospitals should take the initiative to change their systems. This would motivate more healthcare professionals to assess health behaviours of hospitalised smokers and implement evidence-based interventions to help them quit. In particular, healthcare professionals should be given relevant training to enhance their self-efficacy and confidence in promoting smoking cessation to patients. A systematic review indicated numerous barriers to provision of smoking cessation interventions in hospital settings.²⁸ More studies should be conducted to identify barriers specific to the Hong Kong Chinese context.

The results also provide useful recommendations for guiding development of smoking cessation interventions for hospitalised smokers. Our previous studies demonstrated the success of using brief advice, based on the AWARD (Ask, Warn, Advise, Refer and Do-it-again) model, in helping community smokers achieve abstinence.^{29 30} This aids smokers in quitting by warning them about the high risk of premature death resulting from tobacco use, while referring those who require more counselling to existing smoking cessation services.³¹ Compared with intervention using the ‘5A’s’ (Ask, Advise, Assess, Assist and Arrange) approach, that based on the AWARD model is shorter, deliverable within 1 min³¹ and therefore feasible for adoption by healthcare professionals, and even by healthcare assistants, in busy hospital environments. Our study also found >80% of the participants who continued to smoke during hospitalisation reduced their tobacco use. This shows a decrease in daily cigarette consumption from 19.5 to 10.1 among those who continued to smoke after discharge. In fact, in other smoking cessation projects, we have found smokers not interested in quitting may be interested in reducing the number of cigarettes they smoke.^{12 32} Those findings suggest smoking reduction can be an acceptable and feasible alternative approach for starting cessation, especially for hospitalised smokers who do not want to achieve abstinence through abrupt quitting. Future studies are recommended to compare the effectiveness of using two approaches—immediate and progressive—in achieving smoking abstinence among hospitalised smokers.

CONCLUSION

Some smokers continued to smoke during hospitalisation despite smoke-free policies having been implemented in hospitals. A majority of smokers even resumed their smoking habits after discharge. While there is room for improvement in coverage of cessation support among hospitalised smokers, in particular for those in hospitals with a high nurse:patient ratio, a more innovative intervention therefore needs to be developed and evaluated. This should aim for greater effectiveness and feasibility

in healthcare professionals’ efforts to promote smoking cessation for this population.

Acknowledgements We thank the patients who took part in this study. We also thank Adam Goulston, MS, ELS, from Edanz Group (www.edanzediting.com/ac) for editing a draft of this manuscript.

Contributors HCWL, KYH, KKWL, MPW and THL conceived and designed the study and monitored the whole research process. HCWL, KYH, KKWL, WX and LYH searched the literature, reviewed the literature and extracted data. HCWL, KYH, KKWL, MPW, LYH and THL analysed and interpreted the data. HCWL and KYH drafted the manuscript. HCWL, KYH, KKWL, MPW, LYH, KCBT, HKMS, EC, MPH and THL critically revised the manuscript for important intellectual content. All authors approved the final version of the manuscript. We all agree to be accountable for all aspects of the work.

Funding This study was supported by Health and Medical Research Fund, Food and Health Bureau, Hong Kong SAR Government (grant no. 13143141).

Disclaimer The funder had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript or the decision to submit the paper for publication.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Ethical approval was attained from three different Research Ethics Committees, including the Institutional Review Board of the University of Hong Kong/Hospital Authority Hong Kong West Cluster (UW 15-651), the New Territories West Cluster Clinical and Research Ethics Committee (NTWC/CREC/16005) and the Kowloon Central/Kowloon East Cluster Research Ethics Committee (KC/KE-15-0221/ER-1).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement We do not have ethics permission to share the data.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. World Health Organization. Tobacco. 2018 <http://www.who.int/mediacentre/factsheets/fs339/en/> (accessed 1 Mar 2018).
2. U.S. Department of Health and Human Services. *The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General*. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014.
3. Rice DP, Hodgson TA, Sinsheimer P, et al. The economic costs of the health effects of smoking, 1984. *Milbank Q* 1986;64:489–547.
4. Census & Statistics Department. *Pattern of Smoking. Thematic Household Survey Report No. 59: Pattern of Smoking*. Hong Kong: Hong Kong Census & Statistics Department, 2016.
5. McGhee SM, Ho LM, Lapsley HM, et al. Cost of tobacco-related diseases, including passive smoking, in Hong Kong. *Tob Control* 2006;15:125–30.
6. Lando H, Hennrikus D, McCarty M, et al. Predictors of quitting in hospitalized smokers. *Nicotine Tob Res* 2003;5:215–22.
7. Gregor K, Borrelli B. Barriers to quitting smoking among medically ill smokers. *J Behav Med* 2012;35:484–91.
8. Kim SH, Lee JA, Kim KU, et al. Results of an inpatient smoking cessation program: 3-month cessation rate and predictors of success. *Korean J Fam Med* 2015;36:50–9.
9. The Government of the Hong Kong Special Administrative Region. Quitting smoking & tobacco control. 2017 <https://www.gov.hk/en/residents/health/addictions/smoking/tobaccocontrol.htm> (accessed 1 Mar 2018).
10. Rigotti NA, Arnsten JH, McKool KM, et al. Smoking by patients in a smoke-free hospital: prevalence, predictors, and implications. *Prev Med* 2000;31:159–66.
11. Lucas C, Martin J. Smoking and drug interactions. *Aust Prescr* 2013;36:102–4.

12. Fiore MC, Jaén CR, Baker TB, *et al.* *Treating Tobacco Use and Dependence: 2008 Update*. Rockville, MD: US Department of Health and Human Services, 2008.
13. Ho KY, Li WHC, Wang MP, *et al.* Comparison of two approaches in achieving smoking abstinence among patients in an outpatient clinic: a phase 2 randomized controlled trial. *Patient Educ Couns* 2018;101:885–93.
14. Sabidó M, Sunyer J, Masuet C, *et al.* Hospitalized smokers: compliance with a nonsmoking policy and its predictors. *Prev Med* 2006;43:113–6.
15. York NL, Kane C, Beaton K, *et al.* Identifying barriers to hospitalized patients' participation in a free smoking cessation support program. *Medsurg Nurs* 2017;26:25–32.
16. Quality and Safety Division, Hospital Authority. Report on hospital accreditation in hospital authority 2013–2015. 2016 https://accred.ha.org.hk/hosp_accred/overview/160126_Report%20on%20Hospital%20Accreditation%202013-2015.pdf (accessed 14 Jul 2018).
17. Gelenberg AJ, de Leon J, Evins AE, *et al.* Smoking cessation in patients with psychiatric disorders. *Prim Care Companion J Clin Psychiatry* 2008;10:52–8.
18. Doll R, Peto R, Wheatley K, *et al.* Mortality in relation to smoking: 40 years' observations on male British doctors. *BMJ* 1994;309:901–11.
19. Toll BA, O'Malley SS, McKee SA, *et al.* Confirmatory factor analysis of the Minnesota Nicotine Withdrawal Scale. *Psychol Addict Behav* 2007;21:216–25.
20. Huang CL, Lin HH, Wang HH. The psychometric properties of the Chinese version of the Fagerstrom Test for Nicotine Dependence. *Addict Behav* 2006;31:2324–7.
21. Census and Statistics Department. *Thematic Household Survey Report No. 64: Pattern of Smoking*. Hong Kong, 2018. <https://www.statistics.gov.hk/pub/B11302642018XXXB0100.pdf>. (accessed 15 Aug 2018).
22. Wu J, Sin DD. Improved patient outcome with smoking cessation: when is it too late? *Int J Chron Obstruct Pulmon Dis* 2011;6:259–67.
23. Greenhalgh EM, Stillman S, Ford C. Factors that predict success or failure in quit attempts. In: Scollo MM, Winstanley MH, eds. *Tobacco in Australia: facts and issues*. Melbourne: Cancer Council Victoria, 2016.
24. World Health Organization. *Framework Convention on Tobacco Control*. Switzerland, Geneva: the WHO Document Production Services, 2003.
25. The Government of the Hong Kong Special Administrative Region. Nursing manpower in public hospitals. 2016 <http://www.info.gov.hk/gia/general/201605/18/P201605180556.htm> (accessed 1 Mar 2018).
26. Cummins SE, Gamst AC, Brandstein K, *et al.* Helping hospitalized smokers: a factorial RCT of nicotine patches and counseling. *Am J Prev Med* 2016;51:578–86.
27. Richter KP, Faseru B, Shireman TI, *et al.* Warm handoff versus fax referral for linking hospitalized smokers to quitlines. *Am J Prev Med* 2016;51:587–96.
28. Sharpe T, Alshahane A, Ward KD, *et al.* Systematic review of clinician-reported barriers to provision of smoking cessation interventions in hospital inpatient settings. *J Smok Cessat* 2018;1–11.
29. Chan SS, Wong DC, Cheung YT, *et al.* A block randomized controlled trial of a brief smoking cessation counselling and advice through short message service on participants who joined the Quit to Win Contest in Hong Kong. *Health Educ Res* 2015;30:609–21.
30. Wang MP, Li WH, Cheung YT, *et al.* Brief advice on smoking reduction versus abrupt quitting for smoking cessation in chinese smokers: a cluster randomized controlled trial. *Nicotine Tob Res* 2017;20:67–72.
31. Suen YN, Wang MP, Li WH, *et al.* Brief advice and active referral for smoking cessation services among community smokers: a study protocol for randomized controlled trial. *BMC Public Health* 2016;16:387.
32. Lam TH, Cheung YT, Leung DY, *et al.* Effectiveness of smoking reduction intervention for hardcore smokers. *Tob Induc Dis* 2015;13:9.