

Supplementary information

Trends in hepatitis C treatment initiation among HIV/hepatitis C virus-coinfected men engaged in primary care in a multisite community health center in Maryland:
a retrospective cohort study

Yun-Chi Chen, et al.

Supplementary Table

Table S1. Characteristics of the HCV/HIV-coinfected men based on the educational attainment

Educational attainment	High school or less (N = 277)		More than high school (N = 83)		
Characteristics	No.	(%)	No.	(%)	<i>P</i> value
Age					.038
<40	41	(15)	22	(27)	
40 – 49	136	(49)	32	(39)	
≥50	100	(36)	29	(35)	
Race					.002
Black	245	(88)	60	(72)	
White	29	(11)	21	(25)	
Others	3	(1)	2	(2)	
Clinic sites					.81
Baltimore	244	(88)	74	(89)	
Eastern Shore	33	(12)	9	(11)	
Time of entering care					.44
Before 2008	130	(47)	43	(52)	
2008–2013	147	(53)	40	(48)	
Employment					< .001
Unemployed	217	(78)	39	(47)	
Employed	60	(22)	44	(53)	
Type of insurance					< .001
Private	30	(11)	27	(33)	
Medicare	97	(35)	23	(28)	
Medicaid	122	(44)	21	(25)	
Other	28	(10)	12	(14)	
Ever illicit drug use					.005
Yes	260	(94)	69	(83)	
No	17	(6)	14	(17)	
Ever IDU					< .001
Yes	215	(78)	44	(53)	
No	62	(22)	39	(47)	
Sexual behavior					< .001
Non-MSM	220	(79)	32	(39)	
MSM	57	(21)	51	(61)	
HIV RNA at initial visit					.75
≥400 copies/ml	168	(61)	52	(63)	
<400 copies/ml	109	(39)	31	(37)	
Baseline CD4 count					.025
<200 cell/mm ³	82	(30)	19	(23)	
200–499 cell/mm ³	127	(46)	31	(37)	
≥500 cell/mm ³	68	(25)	33	(40)	
Nadir CD4 count					.85
<200 cell/mm ³	160	(58)	47	(57)	
≥200 cell/mm ³	117	(42)	36	(43)	

Abbreviation: HCV, hepatitis C virus; IDU, injection drug use; MSM, men who have sex with men.

Boldface type indicates $P < 0.05$.

Supplementary Table

Table S2. Characteristics of the HCV/HIV-coinfected men entering care before 2008 or between 2008-2013

Characteristics	<2008 enrollees	2008–2013 enrollees	<i>P</i> value
Age (median, inter-quartile range)	N = 182	N = 189	
Age at clinic entry	45.9 (41.7–50.5)	49.4 (44.7–53.6)	<.001
Age as of 1/1/2008	49.2 (44.8–53.2)	46.4 (42.1–50.7)	<.001
Peak fibrosis stage (N, %)	N = 132	N = 143	.96
F0	13 (10%)	17 (12%)	
F1	15 (11%)	16 (11%)	
F2	41 (31%)	46 (32%)	
F3	22 (17%)	20 (14%)	
F4	41 (31%)	44 (31%)	
HIV RNA at initial visit (N, %)	N = 182	N = 189	.002
<400 copies/ml	56 (31%)	88 (47%)	
≥400 copies/ml	126 (69%)	101 (53%)	
HCV genotype (N, %)	N = 172	N = 183	.34
Genotype 1	164 (95%)	170 (93%)	
Other genotype	8 (5%)	13 (7%)	
Type of insurance (N, %)	N = 182	N = 189	.12
Private	34 (19%)	24 (13%)	
Public or other	148 (81%)	165 (87%)	
Educational attainment (N, %)	N = 173	N = 187	.44
>12 years	43 (25%)	40 (21%)	
≤12 years	130 (75%)	147 (79%)	
Being employed (N, %)	N = 182	N = 189	
Yes	62 (34%)	46 (24%)	.040
No	120 (66%)	143 (76%)	