

## SUPPLEMENTARY MATERIAL

*Supplementary File 2. e-Delphi participant demographics for Rounds 1, 2 and 3<sup>1</sup>*

VARIABLE		ROUND 1 (N=38)	ROUND 2 (N=23)	ROUND 3 (N=21)
<b>Age, years</b>	18-24	0 (0%)	0 (0%)	0 (0%)
	25-34	5 (13.2%)	2 (8.7%)	3 (14.3%)
	35-44	13 (34.2%)	10 (43.5%)	7 (33.3%)
	45-54	12 (31.6%)	6 (26.1%)	7 (33.3%)
	55-64	8 (21.1%)	5 (21.7%)	4 (19.1%)
	65+	0 (0%)	0 (0%)	0 (0%)
<b>Gender</b>	Female	32 (84.2%)	19 (82.6%)	18 (85.7%)
	Male	5 (13.1%)	4 (17.4%)	3 (14.3%)
	Other	1 (2.6%)	0 (0%)	0 (0%)
<b>Stakeholder group<sup>2</sup></b>	Clinician	20 (52.6%)	12 (52.2%)	10 (43.4%)
	Researcher	18 (47.4%)	11 (47.8%)	9 (42.8%)
	Decision Maker	12 (31.6%)	5 (21.7%)	7 (33.3%)
<b>Discipline background<sup>2</sup></b>	Nursing	5 (12.5%)	2 (8.7%)	5 (23.8%)
	Occupational therapy	5 (12.5%)	3 (13.0%)	2 (9.5%)
	Physiotherapy	5 (12.5%)	2 (8.7%)	3 (14.3%)
	Medicine	4 (10%)	2 (8.7%)	2 (9.5%)
	Speech pathology	3 (7.5%)	2 (8.7%)	2 (9.5%)
	Psychology	3 (7.5%)	1 (13.0%)	0 (0%)
	Dietetics	3 (7.5%)	2 (8.7%)	2 (9.5%)
	Pharmacy	3 (7.5%)	1 (4.3%)	1 (4.8%)
	Audiology	3 (7.5%)	2 (8.7%)	1 (4.8%)
	Social work	2 (5%)	2 (8.7%)	1 (4.8%)
	Other <sup>3</sup>	4 (10%)	3 (13.0%)	2 (9.5%)
<b>Area of specialty</b>	Multiple clinical specialties	9 (15.5%)	6 (26.1%)	2 (9.5%)
	Geriatrics	13 (22.4)	6 (26.1%)	4 (19.1%)
	Stroke	10 (17.2%)	4 (17.4%)	3 (14.3%)

## SUPPLEMENTARY MATERIAL

<b>(clinicians and researchers)</b>	Acquired brain injury	7 (12.1%)	2 (8.7%)	3 (14.3%)
	Progressive neurological conditions	7 (12.1%)	1 (4.3%)	2 (9.5%)
	Spinal cord injury	2 (3.4%)	0 (0%)	0 (0%)
	Other <sup>4</sup>	10 (17.2%)	1 (4.3%)	1 (4.8%)
<b>Years of experience (clinicians)</b>	1-3	2 (10%)	1 (8.3%)	1 (10.0%)
	3-5	1 (5%)	2 (16.7%)	0 (0%)
	5+	17 (85%)	9 (75.0%)	9 (90.0%)
<b>Career stage (researchers)</b>	Pre-doctoral	7 (38.9%)	3 (27.3%)	4 (44.4%)
	Post-doctoral: Early-mid career researcher ( $\leq 15$ years post-PhD)	7 (39.9%)	5 (45.4%)	2 (22.2%)
	Post-doctoral: Experienced researcher ( $>16$ years post-PhD)	4 (22.2%)	3 (27.3%)	3 (33.3%)
<b>Years of managerial experience (decision makers)</b>	1-3	1 (8.3%)	0 (0%)	0 (0%)
	3-5	0 (0%)	0 (0%)	0 (0%)
	5+	11 (91.7%)	5 (100%)	5 (100%)

<sup>1</sup>Whilst there were 38, 32 and 28 participants in Rounds 1, 2, and 3 (respectively), descriptive data was only available for 23 participants for Round 2 (72%) and 21 participants for Round 3 (75%).

<sup>2</sup>Participants were able to select more than one response.

<sup>3</sup>Includes "Health services research", "Biomedical science", "Rehabilitation Engineering" and "Sociology".

<sup>4</sup>Includes "Major trauma/burns", "Amputation", "Deconditioning", "Scope of practice", "Hearing impairment and rehabilitation", "mental health", "Ankle fracture", "trauma data analysis", "health economics" and "orthopaedic conditions".