

---

## Appendix 1. Systematic search strategies

Medline (PubMed website)

---

((("Guideline"[Publication Type] OR Guideline as Topic"[Mesh] OR 'Practice Guideline'[Publication Type] OR 'Health Planes Guidelines'[Mesh]) OR 'Clinical Protocols'[Mesh])) OR ('Consensus Development Conference, NIH"[Publication Type] OR 'Consensus Development Conference, NIH"[Publication Type] OR ('Consensus'[Mesh]))) OR Standard of Care[Mesh] "Guideline" Publication Type] OR 'Guideline as Topic'[Mesh] OR 'Practice Guideline'[Publication Type] OR 'Health Planes Guidelines'[Mesh] OR 'Clinical Protocols'[Mesh])) OR ('Consensus Development Conference, NIH"[Publication Type] OR 'Consensus Development Conference, NIH"[Publication Type] OR ('Consensus'[Mesh]))) OR Standard of Care[Mesh])) AND ((Depressive Disorder [Mesh] AND Depressive Disorder, Major [Mesh] OR Depressive Disorders OR disorder, Depressive OR Disorders, Depressive OR Neurosis, Depressives OR Depressive Neuroses OR Depressive Neurosis OR Neuroses, Depressive OR Depressions, Endogenous OR Endogenous Depression OR Endogenous Depressions OR Depressive Syndrome OR Depressive Syndromes OR Syndrome, Depressive OR Syndromes, Depressive OR depression, Neurotic OR Depressions, Neurotic OR Neurotic depression OR Neurotic Depressions OR Melancholia OR Melancholias OR Unipolar Depression OR Depression, Unipolar OR Depressions, Unipolar OR Unipolar Depressions)))

---

Cochrane

- # 1 MeSH descriptor: [Guideline] explode all trees
- # 2 MeSH descriptor: [Consensus] explode all trees
- # 3 MeSH descriptor: [Clinical Protocols] explode all trees
- # 4 #1 OR #2 OR #3
- # 5 MeSH descriptor: [Depression] explode all trees
- # 6 #4 AND #5

Embase

---

((practice guideline/mj OR 'consensus development'/exp/mj OR 'clinical protocol'/mj AND ('depression'/exp) AND (2011:py OR 2012:py OR 2013:py OR 2014:py OR 2015:py OR 2016:py AND [embase]/lim)

---

## Appendix 2. Reasons for excluding clinical practice guideline (CPG)

Reference of the excluded CPG	Reasons for exclusion
Institute for Clinical Systems Improvement. Adult Depression in Primary Care. Bloomington, MN: ICSI; 2016.	Duplicate
Austin M-P, Hight N, The Expert Working Group. Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline. Melbourne, Australia: Centre of Perinatal Excellence; 2017.	Duplicate
Grinspun, D, Bajnok I, Rey M. Delirium, Dementia, and Depression in Older Adults: Assessment and Care. Toronto, Canada: Registered Nurses' Association of Ontario; 2016.	Duplicate
National Guideline Clearinghouse. Delirium, Dementia, and Depression in Older Adults: Assessment and Care. Rockville, MD: Agency for Healthcare Research and Quality (AHRQ); 2016. Available from: <a href="https://www.guideline.gov">https://www.guideline.gov</a> . Accessed January 19, 2017.	Duplicate
National Institute for Health and Care Excellence. Depression in Adults: Recognition and Management. 2009. Available from: <a href="https://www.nice.org.uk/guidance/cg90/evidence">https://www.nice.org.uk/guidance/cg90/evidence</a> . Accessed June 30, 2017.	Duplicate
Boltz M (Ed.). Evidence-based Geriatric Nursing Protocols for Best Practice. New York, NY: Springer; 2012.	Duplicate
Depression. University of Michigan Health System. NGC:008672.	Duplicate
Álvarez Ariza M, Atienza Merino G, Ávila González MJ, et al. GPC sobre el Manejo de la Depresión en el Adulto. Madrid, Spain: Ministerio de Sanidad, Servicios Sociales e Igualdad; 2014.	Duplicate
National Guideline Clearinghouse. Depression (Singapore). Rockville, MD: Agency for Healthcare Research and Quality (AHRQ); 2012. Available from <a href="https://www.guideline.gov/summaries/summary/39324">https://www.guideline.gov/summaries/summary/39324</a> . Accessed October 19, 2016.	Duplicate
Austin M-P, Hight N, The Expert Working Group. Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline. Melbourne, Australia: Centre of Perinatal Excellence; 2017.	Duplicate

McDermott B, Baigent M, Chanen A, et al. Clinical Practice Guidelines: Depression in Adolescents and Young Adults. Melbourne, Australia: Agency for Healthcare Research and Quality; 2010.	Duplicate
Michigan Quality Improvement Consortium Guideline. Primary Care Diagnosis and Management of Adults with Depression. Detroit, MI: MQIC; 2018. Available from: <a href="http://mqic.org/guidelines.htm">http://mqic.org/guidelines.htm</a> . Accessed October 19, 2016.	Duplicate
National Institute for Health and Clinical Excellence. Depression in Children and Young People: Identification and Management in Primary, Community and Secondary Care. Leicester, UK: British Psychological Society; 2005.	Duplicate
Michigan Quality Improvement Consortium Guideline. Primary Care Diagnosis and Management of Adults with Depression. Detroit, MI: MQIC; 2018. Available from: <a href="http://mqic.org/guidelines.htm">http://mqic.org/guidelines.htm</a> . Accessed October 19, 2016.	Duplicate
Connolly KR, Thase ME. If at first you don't succeed: a review of the evidence for antidepressant augmentation, combination and switching strategies. <i>Drugs</i> 2011;71(1):43-64. doi:10.2165/11587620-000000000-00000.	Duplicate
National Institute for Health and Care Excellence. Common Mental Health Disorders. Identification and Pathways to Care. London, UK: NICE; 2011.	Without pharmacological treatment
Grinspun, D, Bajnok I, Rey M. Delirium, Dementia, and Depression in Older Adults: Assessment and Care. Toronto, Canada: Registered Nurses' Association of Ontario; 2016.	Duplicate
Joffres M, Jaramillo A, Dickinson J, et al. Recommendations on screening for depression in adults. <i>CMAJ</i> 2013;185(9):775-82. doi:10.1503/cmaj.130403.	Without pharmacological treatment
Boltz M (Ed.). Evidence-based Geriatric Nursing Protocols for Best Practice. New York, NY: Springer; 2012.	Duplicate
Patten SB. Updated CANMAT guidelines for treatment of major depressive disorder. <i>Can J Psychiatry</i> 2016; 61(9):504-5. doi:10.1177/0706743716660034.	Without pharmacological treatment
Rush AJ, Aaronson ST, Demyttenaere K. Difficult-to-treat depression: a clinical and research roadmap for when remission is elusive. <i>Aust N Z J Psychiatry</i> 2019;53(2):109-18. doi:10.1177/0004867418808585.	Without pharmacological treatment
Frye MA. Clinical practice: bipolar disorder--a focus on depression. <i>N Engl J Med</i> 2011;364(1):51-9.	Focused on special groups

Malhi GS, Bassett D, Boyce P, et al. Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for mood disorders. <i>Aust N Z J Psychiatry</i> 2015;49(12):1087-206. doi:10.1177/0004867415617657.	Duplicate
Andersen BL, DeRubeis RJ, Berman BS, et al. Screening, assessment, and care of anxiety and depressive symptoms in adults with cancer: an American Society of Clinical Oncology guideline adaptation. <i>J Clin Oncol</i> 2014;32(15):1605-19. doi:10.1200/JCO.2013.52.4611.	Subject matter
American Medical Directors Association. Guideline Summary: Depression in the Long Term Care Setting. Columbia, MD: AMDA; 2011.	Subject matter
Joffres M, Jaramillo A, Dickinson J, et al. Recommendations on screening for depression in adults. <i>CMAJ</i> 2013;185(9):775-82. doi:10.1503/cmaj.130403.	Subject matter
Li M, Kennedy EB, Byrne N, et al. The management of depression in patients with cancer: a clinical practice guide. <i>J OncolPract</i> 2016;12(8):747-56. doi:10.1200/JOP.2016.011072.	Subject matter
Ostacher MJ, Tandon R, Suppes T. Florida best practice psychotherapeutic medication guidelines for adults with bipolar disorder: a novel, practical, patient-centered guide for clinicians. <i>J Clin Psychiatry</i> 2016;77(7):920-6. doi:10.4088/JCP.15cs09841. Available in: <a href="http://www.embase.com/search/results?subaction=viewrecord&amp;from=export&amp;id=L611538719">http://www.embase.com/search/results?subaction=viewrecord&amp;from=export&amp;id=L611538719</a> .	Subject matter
Pfennig A, Bschor T, Falkai P, et al. The diagnosis and treatment of bipolar disorder: recommendations from the current S3 guideline. <i>Dtsch Arztebl Int</i> 110(6):92-100. doi:10.3238/arztebl.2013.0092.	Subject matter
Rosenblat JD, McIntyre RS. Treatment recommendations for DSM-5-defined mixed features. <i>CNS Spectr</i> 2017;22(2):147-54. doi:10.1017/S1092852916000432.	Subject matter
Alexopoulos GS. Pharmacotherapy for late-life depression. <i>J Clin Psychiatry</i> 2011;72(1):e04. doi:10.4088/JCP.7085tx2cj.	Not a CPG
Andresescu C, Reynolds CF III. Late-life depression: evidencebased treatment and promising new directions for research and clinical practice. <i>Psychiatr Clin North Am</i> 2011;34(2):335-55. doi:10.1016/j.psc.2011.02.005.	Not a CPG
Arnow BA, Steidtmann D, Blasey C, et al. The relationship between the therapeutic alliance and treatment outcome in	Not a CPG

two distinct psychotherapies for chronic depression. <i>J Consult Clin Psychol</i> 2013; 81(4):627-38. doi:10.1037/a0031530.	
Ayub-Dargél A, Masson M, Henry C. The RANZCP guidelines: managing mood disorders in the real world. <i>Aust N Z J Psychiatry</i> 2016;50(12): 1198-9. doi:10.1177/0004867416676373.	Not a CPG
Bland P. Raising standards of care for patients with depression. <i>Practitioner</i> 2011;255(1740):21-5.	Not a CPG
Boffin N, Bossuyt N, Declercq T, et al. Incidence, patient characteristics and treatment initiated for GP-diagnosed depression in general practice: results of a 1-year nationwide surveillance study. <i>Fam Prac</i> 2012;29(6):678-87. doi:10.1093/fampra/cms024.	Not a CPG
Bohra HM, Novak M. Depression in patients with chronic kidney disease. <i>CANNT J</i> 2015;25(3):34-8.	Not a CPG
Busch FN, Sandberg LS. Combined treatment of depression. <i>Psychiatr Clin North Am</i> 2012;35(1):165-79. doi:10.1016/j.psc.2011.10.002.	Not a CPG
Cohen A. The QOF, NICE, and depression. <i>Br J Gen Pract</i> 2011;61(590):549. doi:10.3399/bjgp11X593785.	Not a CPG
Baumann S, Benson-Martin J, Cossie Q, et al. SASOP treatment guidelines for psychiatric disorders: eminence or evidence based? <i>S Afr J Psychiatr</i> 2019;20(2):a529. doi:10.4102/sajpsychiatry.v20i2.529.	Not a CPG
Cosgrove L, Shaughnessy AF, Wheeler EE, et al. The American Psychiatric Association's guideline for major depressive disorder: a commentary. <i>Psychother Psychosom</i> 2012;81(3):186-8. doi:10.1159/000335523.	Not a CPG
Cuijpers P. Effective therapies or effective mechanisms in treatment guidelines for depression? <i>Depress Anxiety</i> 2013;30(11):1055-7. doi:10.1002/da.22205.	Not a CPG
Cuijpers P. Combined pharmacotherapy and psychotherapy in the treatment of mild to moderate major depression? <i>JAMA Psychiatry</i> 2014;71(7):747-8. doi:10.1001/jamapsychiatry.2014.277.	Not a CPG
De Coteau PA, Byrne CD, Russell, V. The HSE/ICGP guidelines on the management of depression and anxiety disorders in primary care. <i>Ir Med J</i> 2012;105(7):251.	Not a CPG
Desseilles M, Witte J, Chang TE, et al. Assessing the adequacy of past antidepressant trials: a clinician's guide to	Not a CPG

the antidepressant treatment response questionnaire. <i>J Clin Psychiatry</i> 2011;72(8):1152-4. doi:10.4088/JCP.11ac07225.	
Canadian Agency for Drugs and Technologies in Health. Diagnosing, Screening, and Monitoring Depression in the Elderly: A Review of Guidelines. Ottawa, ON: CADTH; 2015.	Not a CPG
Canadian Agency for Drugs and Technologies in Health. Antidepressants in Elderly Patients with Major and Minor Depression: A Review of Clinical Effectiveness and Guidelines. Ottawa, ON: CADTH; 2015.	Not a CPG
Gensichen J, Harter M, Klesse C, et al. Germany's national clinical practice guideline (S3) for unipolar depression—What is important for family practice? <i>ZFA</i> 2011;87(5):223-30. doi:10.3238/zfa.2011.0223.	Not a CPG
Geoffroy PA, Bellivier F. The RANZCP mood disorders guidelines: an easy step-by-step toolbox for daily practice. <i>Aust N Z J Psychiatry</i> 2016;50(10):1014-5. doi:10.1177/0004867416667829.	Not a CPG
Gitlin M. The Royal Australian and New Zealand College of Psychiatrists clinical guidelines for mood disorders: kudos and quarrels. <i>Aust NZ Psychiatry</i> 2016;50(10):937-8. doi:10.1177/0004867416668038	Not a CPG
Grobler, G. An overview of depression treatment guidelines. Abstracts 2nd African College of Neuropsychopharmacology Congress 30-31 July 2016 Stellenbosch, Western Cape, South Africa. <i>Acta Neuropsychiatr</i> 2016;28(s3):1-15. doi:10.1017/neu.2016.37.	Not a CPG
Heilmann KE, Wagner M, Riedel-Heller S, et al. Treating late life depression with antidepressants: a summary of recommendations in international guidelines. <i>Fortschr Neurol Psychiatr</i> 2015;83(7):381-91. doi:10.1055/s-0035-1553315.	Not a CPG
Horgan D, Dodd S. Combination antidepressants—use by GPs and psychiatrists. <i>Aust Fam Physician</i> 2011;40(6):397-400.	Not a CPG
Kasper S. Editorial. <i>World J Biol Psychiatry</i> 2013;14(5):333. doi:10.3109/15622975.2013.819703.	Not a CPG
Kendall T, Pilling S, Glover N, et al. Guidelines in mental health - national and international perspectives. <i>Int Rev Psychiatry</i> 2011;23(4):314-7. doi:10.3109/09540261.2011.607431.	Not a CPG

Kongsuk T. Clinical practice guideline major depressive disorder for general practitioners. <i>Value Health</i> 2013;16(7):A695. doi:10.1016/j.jval.2013.08.2091.	Not a CPG
Kurian BT, Grannemann B, Trivedi MH. Feasible evidencebased strategies to manage depression in primary care. <i>Curr Psychiatr Rep</i> 2012;14(4):370-5. doi:10.1007/s11920-0120290-y.	Not a CPG
Laux G. Update treatment of depression—S3 Guideline, internet- based psychotherapy, antidepressants and driving ability. <i>Nervenheilkunde</i> 2016;35(10):691-6. doi:10.1055/s-0037- 1616433.	Not a CPG
Leadholm AKK, Rothschild AJ, Nolen WA et al. The treatment of psychotic depression: is there consensus among guidelines and psychiatrists? <i>J Affect Disord</i> 145(2): 214-20. doi:10.1016/j.jad.2012.07.036.	Not a CPG
Malhi G, Oakley-Browne M, Hay P. Clinical practice guidelines project (CPG project) overview. <i>Aust NZ J Psychiatr</i> 2015;49(Supp 1):30. doi:10.1177/0004867415578344.	Not a CPG
Manning JS, Jackson WC. Providing guideline-concordant assessment and monitoring for major depression in primary care. <i>J Clin Psychiatry</i> 2015;76(1):e3. doi:10.4088/JCP.13013tx7c.	Not a CPG
Mathys M, Mitchell BG. Targeting treatment-resistant depression. <i>J Pharm Pract</i> 2011;24(6):520-33. doi:10.1177/0897190011426972.	Not a CPG
Morris DW, Trivedi MH. Measurement-based care for unipolar depression. <i>Curr Psychiatry Rep</i> 2011;13(6):446-58. doi:10.1007/s11920-011-0237-8.	Not a CPG
Nelson JC. Foreword. <i>CNS Drugs</i> 2013;27:3-4. doi:10.1007/s40263-012-0027-9.	Not a CPG
Nutt DJ. Highlights of the international consensus statement on major depressive disorder. <i>J Clin Psychiatry</i> 2011;72(6):e21. doi:10.4088/JCP.9058tx2c.	Not a CPG
Ogasawara K, Ozaki N. Review of the new treatment guideline for major depressive disorder by the Japanese Society of Mood Disorders. <i>Brain Nerve</i> 2012;64(10):1159-65. In Japanese.	Not a CPG
Oldham J. Fine-tuning our treatment strategies. <i>J Psychiatr Pract</i> 2011;17(3):157.	Not a CPG

doi:10.1097/01.pra.0000398408.13750.8d.	
Pai N. Are the Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for mood disorders meeting the needs of clinicians? <i>Aust N Z J Psychiatry</i> 2016;50(10):1015-6. doi:10.1177/0004867416667828.	Not a CPG
Patkar AA, Pae C-U. Atypical antipsychotic augmentation strategies in the context of guideline-based care for the treatment of major depressive disorder. <i>CNS Drugs</i> 2013;27(Suppl 1):S29-S37. doi:10.1007/s40263-012-0031-0.	Not a CPG
Rawlins M. Ten years of NICE mental health guidelines. <i>Int Rev Psychiatry</i> 2011;23(4):311-3. doi:10.3109/09540261.2011.606804.	Not a CPG
Reisdorf S. Revision of the national health care guidelines on unipolar depression. <i>MedMonatss Pharm</i> 2016;39(4):171-2.	Not a CPG
Roberge P, Fournier L, Brouillet H, et al. A provincial adaptation of clinical practice guidelines for depression in primary care: a case illustration of the ADAPTE method. <i>J Eval Clin Pract</i> 2015;21(6):1190-8. doi:10.1111/jep.12404.	Not a CPG
Schuklenk U, Vathorst SVD. Treatment-resistant major depressive disorder and assisted dying. <i>J Med Ethics</i> 2015;41(8), 577-83. doi:10.1136/medethics-2014-102458.	Not a CPG
Schulte-Korne G, Krick K. In reply. <i>Dtsch Arztebl Int</i> 2014;111(18):330. doi:10.3238/arztebl.2014.0330c.	Not a CPG
Tomba E, Fava GA. Treatment selection in depression: the role of clinical judgment. <i>Psychiatr Clin North Am</i> 2012;35(1):87-98. doi:10.1016/j.psc.2011.11.003.	Not a CPG
Treuer T, Liu C-Y, Salazar G, et al. Use of antidepressants in the treatment of depression in Asia: guidelines, clinical evidence, and experience revisited. <i>Asia Pac Psychiatry</i> 2013;5(4):219-30. doi:10.1111/appy.12090.	Not a CPG
Tundo A, Calabrese JR, Proietti L, et al. Short-term antidepressant treatment of bipolar depression: are ISBD recommendations useful in clinical practice? <i>J Affect Disord</i> 2015;171:155-60. doi:10.1016/j.jad.2014.09.019.	Not a CPG
Van Avendonk M, van Weel-Baumgarten E, van der Weele G, et al. Summary of the Dutch College of General Practitioners' practice guideline "Depression". <i>Ned Tijdschr Geneeskd</i> 2012;156(38):A5101.	Not a CPG
Vinberg M, Levinsen MF, Kessing LV. Treatment-resistant	Not a CPG



depression is treatable. <i>Ugeskr Laeger</i> 2011;173(9):651-4.	
Wang HR, Bahk WM, Park YM, et al. Korean medication algorithm for depressive disorder: comparisons with other treatment guidelines. <i>Psychiatry Investig</i> 2014;11(1):1-11. doi:10.4306/pi.2014.11.1.1.	Not a CPG
Won E, Park SC, Han KM, et al. Evidence-based, pharmacological treatment guideline for depression in Korea, revised edition. <i>J Korean Med Sci</i> 2014;29(4):468-84. doi:10.3346/jkms.2014.29.4.468.	Not a CPG
Xiang YT, Hu C, Wang G, et al. Prescribing patterns of antidepressants, antipsychotics and mood stabilizers in bipolar patients misdiagnosed with major depressive disorder in China. <i>Hum Psychopharmacol</i> 2012;27(6):626-31. doi:10.1002/hup.2262.	Not a CPG
Yang J, Han C, Yoon HK, et al. Experiences and barriers to implementation of clinical practice guideline for depression in Korea. <i>BMC Psychiatry</i> 2013;13:150. doi:10.1186/1471244X-13 - 150.	Not a CPG
Zimmerman M. Symptom severity and guideline-based treatment recommendations for depressed patients: implications of DSM-5's potential recommendation of the PHQ-9 as the measure of choice for depression severity. <i>Psychother Psychosom</i> 2012;81(6):329-32. doi:10.1159/000342262.	Not a CPG
Zimmerman M, Martinez JH, Friedman M, et al. How can we use depression severity to guide treatment selection when measures of depression categorize patients differently? <i>J Clin Psychiatry</i> 2012;73(10):1287-91. doi:10.4088/JCP.12m07775.	Not a CPG
Malhi GS, Outhred T, Hamilton A, et al. Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for mood disorders: major depression summary. <i>Med J Aust</i> 2018;208(4):175-80. doi:10.5694/mja17.00659.	Not a CPG
Evidence-Based Clinical Practice Guidelines for Depression in Adults on Traditional Korean Medicine; 2016.	Not found
S3-Leitlinie/Nationale Versorgungs Leitlinie Unipolare Depression; 2015. In German.	Language
Depression. Adapted Evidence-based Guideline. Ucraina; 2014 - ^enpecia. AganTOBaHaKaH́nHaHacTaHOBa. In Ukrainian.	Language

[Depressive Disorder. <i>Depressie (M44)</i> ]; 2012. In Dutch.	Language
Herrmann M, Gensichen J. Guidelines in general practice: the example depression. <i>Psychiatrie</i> 2015;66:119-24.	Language
Kuroki T, Tanaka T. [Threshold of application of antidepressant drugs for treatment of depressive disorder]. <i>Seishin Shinkeigaku Zasshi</i> 2015;117(4):269-76. In Japanese.	Language
Semba J. [Proper use of novel antidepressants in psychiatric clinical practice]. <i>Seishin Shinkeigaku Zasshi</i> . 2014;116(2):138-143. In Japanese.	Language
Yamada K. [Evidence of treatment for depressive episodes of bipolar disorder]. <i>Seishin Shinkeigaku Zasshi</i> 2011;113(9):873-9. In Japanese.	Language
Siu AL, US Preventive Services Task Force (USPSTF). Screening for depression in adults: US Preventive Services Task Force recommendation statement. <i>JAMA</i> 2016;315(4):380-7. doi:10.1001/jama.2015.18392.	Language
Thase ME. Recommendations for screening for depression in adults. <i>JAMA</i> 2016;315(4):349-50. doi:10.1001/jama.2015.18406.	Language
Patejuk-Mazurek I. Selected antidepressants and anxiolytics: practical guidelines and case studies. <i>Psychiatria</i> 2017;14(3):135-42.	Language
Clinical Practice Guidelines: Depression in Adolescents and Young Adults. Melbourne, Australia: NLMCRC; 2015.	Focused on special groups
Rayner L, Price A, Hotopf M, et al. The development of evidence-based European guidelines on the management of depression in palliative cancer care. <i>Eur J Cancer</i> 2011;47(5):702-12. doi:10.1016/j.ejca.2010.11.027.	Focused on special groups
World Health Organization. Pharmacological Interventions (Antidepressants) for 433 People with Dementia Who Have Associated Depression. Geneva, Switzerland: WHO; 2015. Available from: <a href="https://www.who.int/mental_health/mhgap/evidence/dementia/q4/en/">https://www.who.int/mental_health/mhgap/evidence/dementia/q4/en/</a> . Accessed August 1, 2017.	Focused on special groups
Colquhoun DM, Bunker SJ, Clarke DM, et al. Screening, referral and treatment for depression in patients with coronary heart disease. <i>Med J Aust</i> 2013;198(9):483-4. doi:10.5694/mja13.10153.	Focused on special groups
Galician Health Technology Assessment Agency. Clinical	Outdated

Practice Guideline on the Management of Major Depression in Adults. Santiago de Compostela, Spain: Galician Health Technology Assessment Agency; 2008.	
National Collaborating Centre for Mental Health (UK). Depression. The Treatment and Management of Depression in Adults. Leicester, UK: British Psychological Society; 2010.	Outdated
Lam RW, Parikh SV, Michalak EE, et al. Canadian Network for Mood and Anxiety Treatments (CANMAT) consensus recommendations for functional outcomes in major depressive disorder. <i>Ann Clin Psychiatr</i> 2015;27(2):142-9.	Outdated
Agency for Healthcare Research and Quality. Detecting Depression in Older Adults with Dementia. Ann Arbor, MI: University of Michigan Health System; [s.d].	Outdated
Adult Depression in Primary Care. Bloomington, MN: ICSI; 2013.	Outdated
Galletly C, Castle D, Dark F, et al. Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for the management of schizophrenia and related disorders. <i>Aust N Z J Psychiatry</i> 2016;50(5):410-72. doi:10.1177/0004867416641195.	Outdated
Karyotaki E, Smit Y, Cuijpers P, et al. The Long-term Efficacy of Psychotherapy, Alone or in Combination with Antidepressants, in the Treatment of Adult Major Depression. Good Clinical Practice (GCP). Brussels, Belgium: Belgian psychotherapy Health Care Knowledge Centre (KCE); 2014. Available from: <a href="#">pression_Report_0.pdf</a> . Accessed June 30, 2017.	Focused on psychotherapy
Goracci A, Forgione RN, De Giorgi R, Coluccia A, Cuomo S, Fagiolini A. Practical guidance for prescribing trazodone extended-release in major depression. <i>Expert Opin Pharmacother</i> . 2016;17(3):433-41. doi:10.1517/14656566.2016.1133587	Subject matter