

Appendix 1

Search terms

“isoniazid”, “rifampicin”, “pyrazinamide”, “ethambutol”, “streptomycin”, “rifabutin”, “kanamycin”, “amikacin”, “capreomycin”, “levofloxacin”, “moxifloxacin”, “ofloxacin”, “para-aminosalicylic acid”, “cycloserine”, “terizidone”, “ethionamide”, “protionamide”, “clofazimine”, “linezolid”, “amoxicillin/clavulanate”, “thioacetazone”, “imipenem/cilastatin”, “high-dose isoniazid”, “clarithromycin”, “bedaquiline”, “delamanid”, “meropenem”.

Coding of variables

Type of intervention
Drug therapy
Behavioral
Dots
Supplementation
Rehabilitation
Surgery

Age of participants	
Children	birth to 12 months
	1 to 5 years
	6 to 12 years
Adolescents only (<18)	13 to 17
Adults only	18 and over
Children and Adolescents	
Adolescents and adults	
All ages	
Not Reported	

Ethics
Local
International
Local and International
Not reported

Funding sources
Local Organisation
International organisation
Multiple funding
Industry
University
Hospital
Self funding
Unclear

Co-morbidities
HIV/AIDS
Diabetes
Rheumatoid disease
Multi-morbidity
Not applicable

Type of drug treatment
New drug
New combination
New treatment duration
New dosage

