

<b>Supplementary table 1.</b> Characteristics of 1011 FLS patients that participated and not-participated in this study.			
	<b>Participants</b> (n=500)	<b>Non-participants</b> (n=511)	<b>P-value</b>
Age in years	64.6 ± 8.6	68.3 ± 9.8	<.001
Female sex	357 (71.4)	396 (77.5)	.026
Baseline fracture			
- Finger or toe	58 (11.6)	53 (10.4)	<.001
- Minor	311 (62.2)	259 (50.7)	
- Major	105 (21.0)	157 (30.7)	
- Hip	26 (5.2)	42 (8.2)	
- Fall-related *	431 (86.2)	441 (86.3)	.963
Fall previous year §			
- 0	356 (71.2)	359 (70.3)	.741
- ≥ 1	144 (28.8)	152 (29.7)	
- ≥ 2	72 (14.4)	87 (17.0)	.252
BMD			
- Normal BMD	135 (27.0)	90 (17.6)	<.001
- Osteopenia	255 (51.0)	258 (50.5)	
- Osteoporosis	110 (22.0)	163 (31.9)	
Prevalent vertebral fracture			
- None	366 (73.2)	349 (68.3)	.010
- Grade 1	63 (12.6)	53 (10.4)	
- Grade 2-3	71 (14.2)	109 (21.3)	
At least one fall past year	143 (29.3)	152 (29.9)	.704

Continuous variables are presented as mean ± SD, categorical variables are presented as number of patients (%). § Fall resulting in baseline fracture not included. # According to Genant et al. § According to most severe prevalent vertebral fracture. Abbreviations: BMD, bone mineral density