

Suppl. table 3. Differences in clinical characteristics between the patients with years 0-1 analysis and years 1-2 analysis for all-cause mortality

	Years 0-1 n = 659	Years 1-2 n = 344	<i>p</i> -value
All-cause mortality for 1 year, n (%)	71 (11)	24 (7)	0.051
Age, years	81 ± 9	81 ± 9	0.780
Male sex, n (%)	296 (45)	157 (46)	0.827
Systolic blood pressure, mmHg	121 ± 19	131 ± 21	<0.001
Diastolic blood pressure, mmHg	65 ± 12	68 ± 12	<0.001
Heart rate, bpm	70 ± 15	70 ± 14	0.619
Atrial fibrillation, n (%)	290 (44)	152 (44)	0.956
Coronary artery disease, n (%)	122 (19)	63 (18)	0.938
Diabetes mellitus, n (%)	233 (35)	126 (37)	0.690
Dyslipidemia, n (%)	270 (41)	156 (45)	0.183
Hypertension, n (%)	565 (86)	300 (87)	0.520
<i>Echocardiographic data</i>			
LAD, mm	44 ± 8	44 ± 8	0.452
LAVI, mL/m ²	54 ± 25	53 ± 25	0.949
SV, mL	50 ± 20	48 ± 17	0.309
LVESV, mL	33 ± 17	30 ± 16	0.083
LVEDV, mL	82 ± 34	78 ± 29	0.024
LVEF, %	61 ± 8	62 ± 8	0.013
LVMI, g/m ²	102 ± 32	102 ± 31	0.840
E, m/sec	0.84 ± 0.30	0.86 ± 0.29	0.233
mean e', cm/sec	6.5 ± 2.1	6.6 ± 1.9	0.501
DcT, sec	0.21 ± 0.07	0.23 ± 0.07	0.030
E/e'	13.8 ± 5.7	14.0 ± 5.9	0.677
Ed/Ea	0.130 ± 0.056	0.121 ± 0.051	0.011

Values are presented as means ± standard deviations

or numbers (%).

DcT, deceleration time of E wave; Ea, arterial elastance;

Ed, diastolic elastance; LAD, left atrial diameter;

LAVI, left atrial volume index; SV, stroke volume;

LAV, left atrial volume; LVEF, left ventricular

ejection fraction; LVESV, left ventricular end-systolic

volume; LVEDV, left ventricular end-diastolic volume;

LVMI, left ventricular mass index