

Parents' questionnaire on the suitability for daily use and practical implementation of the new "Diagnosis and Treatment Standard for Neonatal Hypoglycaemia ($\geq 35+0$ weeks of gestation)"

Dear parents,

we would like to thank you again for participating in the ProBrain study.

We would like to ask you a few questions about how you experienced the use of the new treatment standard for hypoglycaemia in neonates.

Who completes this questionnaire: mother father both parents

Please indicate to what extent the statements apply to you.

Example:

Statement	strongly disagree	disagree	slightly disagree	slightly agree	agree	strongly agree
	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5

1. The blood glucose measurements gave me a sense of reassurance.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
2. I understand the rationale for the blood glucose measurements.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
3. I consider regular blood glucose measurements to detect hypoglycaemia in my baby to be useful.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
4. The measurements made me more concerned and worried about my baby's health.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
5. My baby's blood glucose measurements put additional burden on me.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
6. The blood glucose measurements stressed my baby.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
7. I consider the burden of the measures to be well founded and justified.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
8. The flyer on hypoglycaemia in babies gave me enough information and educated me well on the subject.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
9. The information on the flyer worried me more than it helped me.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
10. Before I participated in the ProBrain study, I already knew that hypoglycaemia could occur in neonates.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
11. I was very worried that my baby would be given formula because of hypoglycaemia and would develop breastfeeding problems.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5

12. I think avoiding supplemental feeding of even small amounts of formula is important.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
13. I would like to get more information in person by a doctor and/or nurses.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
14. Following a precise scheme for the detection and treatment of hypoglycaemia in neonates makes me feel that hypoglycaemia in my baby can be detected early.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
15. The standard for detection and treatment of hypoglycaemia in neonates makes me feel that hypoglycaemia in my baby can be treated well.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
16. I would recommend the Department of Obstetrics and Gynaecology at the University Hospital Düsseldorf because babies are cared for very carefully there with regard to hypoglycaemia.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
17. I will be happy to attend the scheduled follow-up visit when my child is two years old.	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5

Do you have any remarks/comments or suggestions for improvement?

Thank you for your participation!